Introduction

Objekttyp: Chapter

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech.

Hochschule, Stiftung Rübel, in Zürich

Band (Jahr): 71 (1986)

PDF erstellt am: 30.04.2024

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek* ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

1. INTRODUCTION

At the time of Hieronymus Bock (1498-1554) Lemna was supposed to be a first development stage of watercress. VALLISNERI (1714 according to MARIE-VICTORIN 1931) was the first to describe flowers of Lemna. 1729, MICHELI observed flowers of Lemna again. He named the flowering Lemna plants Lenticula and the non-flowering ones Lenticularia. (1753, 1771) distinguished five species (Lemna polyrhiza, L. gibba, L. minor, L. trisulca, L. arrhiza) within the Lemnaceae family all of which he attributed to the genus Lemna. The first monographic treatment of Lemnaceae was published 1839 by SCHLEIDEN. In 1868, HEGELMAIER completed his extensive monograph. Twenty-seven years later (1895) he wrote a new survey of the family with supplementary information. His monograph was so competent and extensive that it took nearly a hundred years until a new monograph appeared (DAUBS 1965). The emphasis in the work of HEGELMAIER was put on comparative morphology and histology. He studied the development of members of Lemnaceae with such an accuracy and such a technical skill that his work was not matched since, though the methods have been improved considerably in the last hundred years. The present monographic treatment does not contain all the details of HEGELMAIER's ontogenetic descriptions of the Lemnacean organs. Scientists interested in the ontogeny and histology of Lemnaceae still must consult the monograph of HEGELMAIER. Due to his absolutely reliable but sometimes pedantic perfection he wrote in a rather complex language with frequent involved passages which makes the reading of his papers difficult, for people of German mother tongue. DAUBS (1965) has the merit of having investigated much living material especially from the New World. He had access to new literature particularly from the English speaking countries. However, since his monograph was written as a dissertation, it is not astonishing that his achievements were rather incomplete. The work of HEGELMAIER was evaluated only in part, probably due to his very complex and not easily understandable language. Also, DAUBS reviewed only herbarium specimens from nine North American herbaria. For comparison, the present author consulted 120 herbaria from all over the world. incompleteness of DAUBS' monograph inspired DEN HARTOG and VAN DER PLAS (1970) to examine critically the taxonomic status of the different

members of <u>Lemnaceae</u> in a "Synopsis of <u>Lemnaceae</u>". Partial reviews of taxonomy and ecology of the family were done by THOMPSON (1898) for North America, and by LUDWIG (1909) and KANDELER (1979) for Europe. A modern synecological treatment of the family (mostly for Europe) originates from SCHWABE-BRAUN and TUEXEN (1981b).

Within the last sixty years the family of Lemnaceae became more and more important as suitable research material for plant physiology and phytochemistry. Some of the advantages of Lemnaceae are: fast and predominantly vegetative reproduction (genetically uniform clones), only small laboratory space requirement, and aseptic culture. In addition, the family aroused attention as a food source due to its high productivity and its high protein content. Furthermore, Lemnaceae are used as multipurpose test objects and for nutrient and mineral removal from sewage water. The literature on Lemnaceae has been growing exponentially for the last 30 years. HILLMAN (1961) cites 250 titles in his review of the descriptive and experimental literature of the family Lemnaceae. In 1980, the list of titles on Lemnaceae had increased to about 1300 titles (LANDOLT 1980d). The bibliography of the present work which will be published in volume 2 (LANDOLT and KANDELER 1987) contains more than 3000 titles.

The present author began his studies on Lemnaceae 1953 in California with comparative physiological and ecological investigations LANDOLT 1957). Since then, work on Lemnaceae was continued slowly with some supplementary physiological work and with the building up of a representative collection of more than 1000 living clones of Lemnaceae from all over the world. Some fieldtrips in Europe, North America, Argentina, Malaysia, and Australia completed the knowledge on the family. An original intention to write a monograph on Lemnaceae always had to be postponed due to many other duties and obligations. Ten years ago, I began to invest more time to complete the monograph, but sometimes the new literature grew faster than the available time to study the papers. Also, some of the new papers were so specialized that I no longer felt able to handle them competently. Fortunately, with Prof. Dr. Riklef Kandeler from Vienna (Austria) I found not only a very competent coauthor for the second volume of the monograph but also a very cooperative and generous friend. Prof. Kandeler has been familiar with the family of Lemnaceae for more than 30 years and he has the true perspective of modern plant physiology and phytochemistry.

The aim of this monograph is to give a survey of the present status of knowledge on <u>Lemnaceae</u>. All characteristics and all possible aspects should be presented and discussed. It appeared to be impossible to evaluate every paper dealing with <u>Lemnaceae</u> in detail, but we hope that a representative selection is given and that at least the bibliography in volume 2 is more or less complete.

The names of the species used in literature have been adapted to the present nomenclature (e.g. <u>Lemna aequinoctialis</u> instead of <u>L. perpusilla</u> or <u>L. paucicostata</u>, or <u>W. borealis</u> for <u>W. punctata</u>). Where the identity of the species used for experiments was not correct according to the knowledge of the present author, the name has been changed but the name used originally is put in brackets ("named as").

The first volume contains the morphology, ecology, geographical distribution, taxonomy, and nomenclature of the family. The second volume deals with phytochemistry, physiology, and application. It also includes the bibliography. In order to give as complete as possible a survey it could not be avoided that some findings and results appear in more than one chapter.

It is hoped that the present study may help to further the better understanding of the special characteristics and metabolic processes within the family of <u>Lemnaceae</u>. Also it is intended to stimulate work with <u>Lemnaceae</u> and the following up of the many problems which are still to be solved.

ACKNOWLEDGEMENTS

The final results of such an extensive and long lasting work could not have been achieved without the active and unselfish cooperation of collaborators which is very much appreciated. I am afraid that it is not possible to duly acknowledge here the valuable work of all the contributors. I received much advice, information, and help from colleagues from all parts of the world. Many living and dried samples of Lemnaceae arrived here within the last five years. The many collectors of living samples sent before 1980 are presented on page 9 to 11 of the first Lemnaceae volume (LANDOLT 1980a). To all the new contributors listed below, I am very grateful: W.R. Armstrong (San Marcos, Cal., U.S.A.), M. Baltisberger (Zürich, Switzerland), E.O. Beal+ (Bowling Green, Ky., U.S.A.), H.W. Beams (Iowa City, Iowa, U.S.A.), T. Beppu (Kyoto, Japan), B. Beyer (Zürich, Switzerland), J. Bogner (München, FRG), D.E. Boufford (Pittsburgh, Pa., U.S.A.), R.E. Brooks (Lawrence, Kansas, U.S.A.), R. Buchecker (Zürich, Switzerland), B.M.H. Burdet (Geneva, Switzerland), W. Büttiker (Jedda, South Arabia), K.P. Buttler (Frankfurt FRG), P.M. Catling (Ottawa, Canada), C.A. Ceska (Victoria, Canada), H.L. Clark (New Orleans, La., U.S.A.), Ch.F. Cleland (Rockville, Md., U.S.A.), C.D.K. Cook (Zürich, Switzerland), D.D. Culley (Baton Rouge, La., U.S.A.), O. Degener (Volcano, Hawaii, U.S.A.), H. Diekjobst (Iserloh, FRG), J.L. Dudley (Bethlehem, Pa., U.S.A.), G. Dussaussois (Pessac, France), Eskuche (Corrientes, Argentina), C.H. Eugster (Zürich, Switzerland), C. Eyster (Mobile, Al., U.S.A.), C. Farney (Rio de Janeiro, Brazil), O.A. Fernandez (Bahia Blanca, Argentina), G. Geber (Vienna, Austria), Gigon (Zürich, Switzerland), B. Giles (Cambridge, U.K.), D. Goepfert (Toronto, Canada), M.H. Grayum (St. Louis, Mo., U.S.A.), A. Gröber (Zürich, Switzerland), Ch.W. Heekman (Hamburg, FRG), H. Heine (Paris, France), F.N. Hepper (Kew, U.K.), P. Hiepko (Berlin, FRG), W.S. Hillman+ (New Haven, Co., U.S.A.), S.I. Honda (Dayton, Ohio, U.S.A.), A. Hussain P. and S. Jovet-Ast (Karachi, Pakistan), H. Jänichen (Berlin, GDR), (Paris, France), Ch. Kasselmann (Frankfurt, FRG), R.G. Kessel (Iowa City, Iowa, U.S.A.), F.K. Keneally (South Perth, Australia), A. Klein (Liestal, Switzerland), F. Klötzli (Zürich, Switzerland), L. Koumpli-Sovanthi (Athens, Greece), B. Krajncic (Maribor, Yugoslawia), B. Krüsi (Zürich, Switzerland), I. Kukkonen (Helsinki, Finland), J.B. Lachavanne (Geneva, Switzerland), A. Lawalrée (Meise, Belgium), E. Leupi (Aarau, Switzerland), R.A. Lewin (La Jolla, Cal., U.S.A.), R.M. Lowden (Santiago, Dom. Rep.), A. Lüönd (Zürich, Switzerland), B.W. Lüönd (Zürich, Switzerland), M.U. Marchant (South Perth, Australia), J.W. McClure (Oxford, Ohio, U.S.A.), M.E. Medley (Louisville, Ky., U.S.A.), M. Meyer (Davos, Switzerland), L.J. Musselman (Khartoum, Sudan), B. Nordenstam (Stockholm, Sweden), S. Omer (Karachi, Pakistan), S. Otaki (Tokyo, F.D. Ott (Memphis, Te., U.S.A.), T.M. Pedersen (Mburucuya, Argentina), F. Pedrotti (Camerino, Italia), A.H. Pieterse (Amsterdam, Netherlands), E. Polar (Istanbul, Turkey), D. Porath (Beer-Sheva, Israel), G.R. Proctor (Cambridge, Mass., U.S.A.), C. Ramirez (Valdivia, Chile), M. Ratnasabapathy (Kuala Lumpur, Malaysia), A. Raynal-Roques (Paris, France), E. Rejmankova (Davis, Cal., U.S.A.), H. Scholz (Berlin, FRG), D. Schotsman (Paris, France), A. Schwabe-Braun (Freiburg, GRD), Ch. Scotter (Cardiff, U.K.), C.Y. Shih (Laguna Hills, Cal., U.S.A.), H. Sigg (Winterthur, Switzerland), Ch. Simon (Basel, Switzerland), J.R. de Sloover (Louvain, Belgium), L. Steubing (Giessen, FRG), A. Takimoto

(Kyoto, Japan), J.W. Thieret (Highland Heights, Ky., U.S.A.), K. Tobolski (Kierownik, Poland), P. Uotila (Helsinki, Finland), K. Urbanska (Zürich, Switzerland), J. Vivant (Orthez, France), J. Waddington (Swift Current, Canada), S.M. Walters (Cambridge, U.K.), J. Wolek (Krakow, Poland), R. Yusuf (Karachi, Pakistan), K. Zarzycki (Krakow, Poland), E. Zogg (Walenstadt, Switzerland), E. Zuber (Grüningen, Switzerland).

The ready cooperation of all the directors and curators of the more than hundred herbaria which are listed in chapter 6 is thankfully acknowledged. The following colleagues kindly permitted the use of their photographs: J.L. Anderson, Logan, Utah, U.S.A. (figs. 2.15, 2.17, 2.22), W.R. Armstrong, San Marcos, Cal., U.S.A. (figs. 2.24, 2.43, 2.61, plates Xa,b,c, XIIa,b,c,d,e, XIIId, XIVa, XVIa). H.W. Beams, Iowa City, Iowa, U.S.A. (fig. 2.19), T. Beppu, Kyoto, Japan (fig. 3.2), G. Geber, Vienna, Austria (fig. 3.3), M.H. Grayum, St. Louis, Mo., U.S.A. (figs. 2.53, 2.54), R. Kandeler, Vienna, Austria (figs. 2.48, 2.66, 2.67), F. Klötzli, Zürich, Switzerland (fig. 4.1), C. Le Pabic, Paris, France (figs. 2.16, 2.18), C.Y. Shih, Laguna Hills, Cal., U.S.A. (figs. 2.32, 2.46, 2.57), K. Urbanska, Zürich, Switzerland (fig. 3.1), E. Wehrli, Zürich, Switzerland (fig. 2.64, 2.65). The technical help and the tremendous and varied work which is needed to produce such a monograph is especially appreciated. First of all I would like to thank Mrs Anita Hegi who curates the collection of living samples of Lemnaceae. She did splendid work in transforming the newly arrived clones into aseptic cultures and to keep more than a thousand clones safe and sound for many years. She also oversaw the cultivation experiments and put all the data together. She made the drawings with accuracy and mounted the plates. In addition, she was in charge of finding the literature which was not available in our own library. Mrs Erika Wohlmann-Bräm also did excellent work as predecessor of Mrs Anita Hegi as curator of the Lemnaceae collection. She is also responsible for a few graphs in the present work. Hans Sigg and René Graf did the photographic work with great skill and patience. Dr. Martin Wehrli kindly reproduced the SEM figures of seeds, and Dr. M.H. Grayum (St. Louis, Mo., U.S.A.) of the pollen grains. Erwin Schaeffer analyzed the water samples. The laborious work of copying the indications from the herbaria and of putting together and classifying the geographical information was done by Werner Engetschwiler. Mrs Anny Honegger typed and prepared carefully for printing and dealt with the voluminous Lemnaceae correspondence. Additional typing was done by Mrs Berti Brun and Mrs Renata Flubacher. Claude Fornallaz, Monique Honegger, and Dr. Georg Zumbühl read proofs. The bibliography was supplemented first by Dr. Nora Winterhalter; in the last years the library service of the Swiss Federal Institute of Technology delivered all information on new literature on Lemnaceae. A very welcome supplementary service was received from the Aquatic Weed Program Bibliography, University of Florida, Gainesville, U.S.A. The manuscript was carefully and critically checked by Prof. Dr. R. Kandeler, Vienna, Austria, and partly by Dr. H.L. Clark, New Orleans, La., U.S.A. I am very thankful for their constructive and very valuable cooperation. The English was kindly checked by Lee Ann Clark (up to 1984) and Sonia Türler.

Many sincere thanks to all of these persons and to all others who contributed to the present work.