

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübél, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübél (Zürich)

Band: 71 (1986)

Artikel: Biosystematic investigation in the family of duckweeds ("Lemnaceae"). Vol. 2 : the family of "Lemnaceae" : a monographic study. Volume 1

Autor: Landolt, Elias

Register: List of the figures and plants in volume 1

DOI: <https://doi.org/10.5169/seals-308748>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LIST OF THE FIGURES AND PLATES IN VOLUME 1

| | |
|---|----|
| 2. 1. Groups of fronds of various <u>Lemnaceae</u> | 14 |
| 2. 2. Young fronds of <u>Spirodela polyrrhiza</u> | 16 |
| 2. 3. Longitudinal section of <u>Wolffiella lingulata</u> | 17 |
| 2. 4. Longitudinal section of <u>Wolffia arrhiza</u> | 18 |
| 2. 5. Light micrographs of <u>Wolffia arrhiza</u> | 19 |
| 2. 6. Interpretation of the <u>Spirodela</u> frond, according to BRUNAUD (1974a,b) | 21 |
| 2. 7. Interpretation of the <u>Lemnaceae</u> frond, according to BROOKS (1940) and DAUBS (1965) | 22 |
| 2. 8. Interpretation of the <u>Lemna</u> frond, according to ASHBY et al. (1949). | 23 |
| 2. 9. Homology in <u>Pistia</u> and <u>Spirodela</u> , according to MEUSEL (1951) | 24 |
| 2.10. Interpretation of the <u>Lemnaceae</u> frond in the present work | 25 |
| 2.11. Changes of frond area of <u>Lemna minor</u> grown at 20°C and 5000 lux light intensity in the course of generations | 30 |
| 2.12. Diagram of the vegetative growth cycle of <u>Lemna aequinoctialis</u> | 32 |
| 2.13. Frond chains in <u>Lemna trisulca</u> | 33 |
| 2.14. <u>Lemna trisulca</u> producing spiral chains in the half-shade | 34 |
| 2.15. Young frond primordia of <u>Wolffia arrhiza</u> arising in the reproductive pouch of a mother frond | 40 |
| 2.16. Ultrastructure of a parenchymatic cell of <u>Spirodela polyrrhiza</u> | 41 |
| 2.17. Ultrastructure of a guard cell of a mature frond of <u>Wolffia arrhiza</u> | 42 |
| 2.18. Chloroplast of <u>Spirodela polyrrhiza</u> | 43 |
| 2.19. Chloroplasts of <u>Lemna</u> sp. | 44 |
| 2.20. Ultrastructure of young daughter frond cells of <u>Wolffia arrhiza</u> | 45 |
| 2.21. Upper epidermis of <u>Lemna aequinoctialis</u> and <u>Wolffia brasiliensis</u> | 49 |
| 2.22. Comparison of chloroplasts of <u>Wolffia arrhiza</u> in epidermal and in palisade-like cells | 50 |
| 2.23. Longitudinal section of stomata cells of <u>Lemna minor</u> | 51 |
| 2.24. <u>Wolffia brasiliensis</u> with a prominent papilla | 53 |
| 2.25. Transverse section of the distal part of <u>Lemna gibba</u> , showing the aerenchymatous tissue | 55 |
| 2.26. <u>Lemna gibba</u> . Gibbous fronds from below showing air spaces and roots | 55 |
| 2.27. Transverse sections of <u>Lemna gibba</u> fronds | 57 |
| 2.28. Basal part of the <u>Lemna trisulca</u> frond | 59 |
| 2.29. Nerves of <u>Lemna gibba</u> and <u>Lemna minor</u> | 60 |
| 2.30. Section of a frond segment of <u>Spirodela punctata</u> | 62 |
| 2.31. Raphide cell from the root of <u>Lemna minor</u> | 62 |
| 2.32. A developing young frond of <u>Lemna aequinoctialis</u> | 65 |
| 2.33. Transverse section of a young root sheath of <u>Lemna aequinoctialis</u> | 71 |
| 2.34. Rootcap of <u>Lemna gibba</u> | 72 |
| 2.35. Longitudinal section of the root tip of <u>Lemna minor</u> | 73 |
| 2.36. Transverse section of the root tip of <u>Spirodela polyrrhiza</u> and <u>Lemna minor</u> | 75 |
| 2.37. Transverse section of the root tip of <u>Lemna minor</u> | 76 |

| | |
|---|-----|
| 2.38. Organs of stabilization | 78 |
| 2.39. Turion of <u>Spirodela polyrrhiza</u> from below | 82 |
| 2.40. Flowering <u>Wolffiella Welwitschii</u> | 90 |
| 2.41. Two flowering fronds of <u>Wolffia columbiana</u> | 90 |
| 2.42. Flower of <u>Lemna disperma</u> | 93 |
| 2.43. Flower of <u>Wolffia angusta</u> | 93 |
| 2.44. Flowers of <u>Spirodela polyrrhiza</u> and <u>Lemna trisulca</u> | 94 |
| 2.45. Flower diagram of <u>Spirodela polyrrhiza</u> | 94 |
| 2.46. Flower of <u>Lemna aequinoctialis</u> | 95 |
| 2.47. Longitudinal section of a flowering frond of <u>Wolffia microscopica</u> | 95 |
| 2.48. Frond and flower primordia of <u>Lemna gibba</u> | 96 |
| 2.49. Transverse section of an anther of <u>Spirodela polyrrhiza</u> | 98 |
| 2.50. Longitudinal section of an anther of <u>Lemna aequinoctialis</u> during meiosis | 98 |
| 2.51. Size of pollen grains of two clones each of <u>Lemna aequinoctialis</u> and <u>Lemna turionifera</u> with varying chromosome numbers ($2n=40$ and 80) | 99 |
| 2.52. Pollen grain of <u>Lemna minor</u> | 99 |
| 2.53. Pollen grains of different <u>Lemnaceae</u> | 100 |
| 2.54. Surface of a pollen grain of <u>Lemna trisulca</u> | 102 |
| 2.55. Ovule with mature embryo sac of <u>Spirodela polyrrhiza</u> | 107 |
| 2.56. Ovule of <u>Lemna aequinoctialis</u> | 108 |
| 2.57. Pollen grains of <u>Lemna aequinoctialis</u> germinating on the stigma | 109 |
| 2.58. Fertilized embryo sac of <u>Lemna aequinoctialis</u> showing proembryo and 8-celled endosperm | 110 |
| 2.59. Longitudinal section of a seed of <u>Wolffia microscopica</u> | 111 |
| 2.60. Longitudinal section of the distal part of a seed of <u>Lemna aequinoctialis</u> | 112 |
| 2.61. Spherical droplet on the stigma of <u>Wolffiella lingulata</u> | 113 |
| 2.62. Fruits of different <u>Lemnaceae</u> | 116 |
| 2.63. Seeds of some <u>Lemnaceae</u> | 118 |
| 2.64. Seeds of different <u>Lemnaceae</u> | 120 |
| 2.65. Seed structures of different <u>Lemnaceae</u> | 121 |
| 2.66. Transverse sections of the fruits of <u>Lemna aequinoctialis</u> and <u>Lemna perpusilla</u> | 122 |
| 2.67. Transverse section through a germinating seed of <u>Lemna gibba</u> | 124 |
| 3.1. Chromosome preparations by URBANSKA | 129 |
| 3.2. Chromosome preparations by BEPPU | 131 |
| 3.3. Chromosome preparations by GEBER | 133 |
| 4.1. <u>Lemna minor</u> growing in the drip of a waterfall in Semien (Ethiopia) | 141 |
| 5.1. Mass and energy flow in a duckweed system | 187 |
| 5.2. Schematic presentation of a <u>Lemnetea</u> community | 207 |
| 5.3. <u>Lemna minuscula</u> as a pioneer species in an outlet of a spring (Argentina) | 211 |
| 5.4. Association of <u>Lemna gibba</u> and <u>Wolffiella oblonga</u> (Argentina) | 223 |
| 6.1. Geographic divisions of the world | 234 |
| 6.2. Distribution of the family of <u>Lemnaceae</u> | 235 |
| 6.3. Number of <u>Lemnaceae</u> species within the different regions of the world | 236 |
| 6.4. Distribution of <u>Spirodela intermedia</u> | 242 |

| | | |
|-------|---|-----|
| 6.5. | Distribution of <u>Spirodela polyrrhiza</u> | 254 |
| 6.6. | Distribution of <u>Spirodela punctata</u> | 261 |
| 6.7. | Distribution of <u>Lemna gibba</u> and <u>L. disperma</u> | 272 |
| 6.8. | Distribution of <u>Lemna minor</u> and <u>L. japonica</u> | 282 |
| 6.9. | Distribution of <u>Lemna ecuadoriensis</u> and <u>L. obscura</u> | 286 |
| 6.10. | Distribution of <u>Lemna turionifera</u> | 291 |
| 6.11. | Distribution of <u>Lemna trisulca</u> | 300 |
| 6.12. | Distribution of <u>Lemna perpusilla</u> and <u>L. tenera</u> | 302 |
| 6.13. | Distribution of <u>Lemna aequinoctialis</u> | 319 |
| 6.14. | Distribution of <u>Lemna valdiviana</u> | 326 |
| 6.15. | Distribution of <u>Lemna minuscula</u> | 334 |
| 6.16. | Distribution of <u>Wolffiella hyalina</u> , <u>W. repanda</u> , <u>W. rotunda</u> and <u>W. neotropica</u> | 338 |
| 6.17. | Distribution of <u>Wolffiella Welwitschii</u> | 342 |
| 6.18. | Distribution of <u>Wolffiella lingulata</u> | 346 |
| 6.19. | Distribution of <u>Wolffiella oblonga</u> | 351 |
| 6.20. | Distribution of <u>Wolffiella gladiata</u> and <u>W. denticulata</u> | 354 |
| 6.21. | Distribution of <u>Wolffia microscopica</u> and <u>W. elongata</u> | 356 |
| 6.22. | Distribution of <u>Wolffia brasiliensis</u> | 361 |
| 6.23. | Distribution of <u>Wolffia borealis</u> and <u>W. australiana</u> | 364 |
| 6.24. | Distribution of <u>Wolffia angusta</u> and <u>W. arrhiza</u> | 370 |
| 6.25. | Distribution of <u>Wolffia columbiana</u> | 376 |
| 6.26. | Distribution of <u>Wolffia globosa</u> | 380 |
| 6.27. | Highest known stations of <u>Spirodela polyrrhiza</u> in relation to the latitude | 385 |
| 6.28. | Highest known stations of <u>Lemna gibba</u> in relation to the latitude | 385 |
| 6.29. | Highest known stations of <u>Lemna minor</u> in relation to the latitude | 387 |
| 6.30. | Highest known stations of <u>Lemna trisulca</u> in relation to the latitude | 387 |
| 6.31. | Highest known stations of <u>Lemna aequinoctialis</u> in relation to the latitude | 389 |
| 6.32. | Highest known stations of <u>Lemna minuscula</u> in relation to the latitude | 389 |
| 6.33. | Highest known stations of <u>Wolffiella oblonga</u> in relation to the latitude | 390 |
| 6.34. | Climate zones of the world | 405 |
| 6.35. | Distribution of <u>Spirodela punctata</u> with the year of the first record | 407 |
| 7.1. | Correlation between the DNA content and the degree of primitivity | 419 |
| 7.2. | Supposed relationship between the genera of the <u>Lemnaceae</u> | 420 |
| 7.3. | Genetic distance between some species of <u>Lemna</u> | 423 |
| 7.4. | Supposed taxonomical relationship within the genus <u>Lemna</u> | 424 |
| 7.5. | Supposed taxonomical relationship within the section <u>Lemna</u> | 425 |
| 7.6. | Supposed taxonomical relationship within the genus <u>Wolffiella</u> | 426 |
| 7.7. | Supposed taxonomical relationship within the genus <u>Wolffia</u> | 427 |
| 7.8. | Supposed taxonomical relationship within the section <u>Wolffia</u> | 428 |
| 9.1. | Transparent preparations of <u>Spirodela</u> species | 497 |
| 9.2. | Transparent preparations of <u>Lemna</u> species | 498 |
| 9.3. | Transparent preparations of <u>Wolffiella</u> species | 503 |
| 9.4. | Transparent preparations of <u>Wolffia</u> species | 508 |

Plates I to XVI: Colour photographs of Lemnaceae species and
typical habitats

513-528

LIST OF THE TABLES IN VOLUME 1

| | | |
|------|--|-----|
| 3.1. | Chromosome numbers of <u>Lemnaceae</u> | 134 |
| 3.2. | DNA content of the chromosomes of <u>Lemnaceae</u> and <u>Pistia</u> | 137 |
| 4.1. | Range of nutrient content in waters with <u>Lemnaceae</u> | 150 |
| 4.2. | Lowest yearly mean of some nutrients in waters of Central Europe with <u>Lemnaceae</u> species | 150 |
| 4.3. | Lowest nutrient demand of some <u>Lemnaceae</u> species | 165 |
| 4.4. | Three levels of approximate minimum nutrient requirements in <u>Lemnaceae</u> | 165 |
| 4.5. | Flowering and fruiting percentage of <u>Lemnaceae</u> species in nature | 169 |
| 4.6. | Flowering and fruiting time of <u>Lemnaceae</u> species | 172 |
| 4.7. | Chemical composition of waters with flowering <u>Lemnaceae</u> | 176 |
| 5.1. | Competition for nitrogen and phosphorus between <u>Lemna turionifera</u> , <u>Spirodela polyrrhiza</u> , and <u>Wolffia borealis</u> | 206 |
| 5.2. | Seasonal changes of the <u>Lemnaceae</u> vegetation in a pond in California | 209 |
| 6.1. | Number of species of each genus of the <u>Lemnaceae</u> within main regions of the world | 237 |
| 6.2. | The highest known localities of some species of <u>Lemnaceae</u> | 382 |
| 6.3. | Ecological limits of <u>Lemnaceae</u> distribution | 393 |
| 6.4. | Minimal duration of days with mean temperatures higher than 10°C for <u>Lemnaceae</u> species | 395 |
| 6.5. | Temperature limits of the three coolest months for <u>Lemnaceae</u> species | 397 |
| 6.6. | Temperature limits of the three warmest months for <u>Lemnaceae</u> species | 399 |
| 6.7. | Lowest mean temperature of the three warmest months needed for <u>Lemnaceae</u> species | 400 |
| 6.8. | Limits of the occurrence of <u>Lemnaceae</u> according to the aridity factor of Martonne | 401 |
| 6.9. | Occurrence of <u>Lemnaceae</u> species within climate zones | 403 |
| 7.1. | Different characteristics of <u>Spirodela</u> and <u>Pistia</u> | 415 |
| 7.2. | Primitive and derived characteristics within the <u>Lemnaceae</u> family | 418 |
| 7.3. | Degree of primitivity of the <u>Lemnaceae</u> species | 422 |
| 9.1. | Characteristics of the species of section <u>Lemna</u> | 478 |
| 9.2. | Occurrence of the species of section <u>Lemna</u> | 479 |
| 9.3. | Morphological characteristics of <u>Lemna aoukikusa</u> | 488 |
| 9.4. | Characteristics typical for species of the <u>Wolffiella oblonga</u> group | 539 |