

**Zeitschrift:** Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

**Herausgeber:** Geobotanisches Institut, Stiftung Rübel (Zürich)

**Band:** 93 (1987)

**Artikel:** Vegetation of stabilizing and eroding slopes in eastern Nepal = Die Vegetation stabilisierender und erodierender Hänge in Ostnepal

**Autor:** Schaffner, Ruth

**Kapitel:** 2: The investigated area

**DOI:** <https://doi.org/10.5169/seals-308864>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 04.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## **2. THE INVESTIGATION AREA**

### **2.1. PHYSIOGRAPHY**

The present study covers an area in eastern Sindhupalchok and western Dolakha District between the Sunkosi and Tamakosi Rivers in the central midlands of Nepal (Figs. 2.1, 2.2). The relief is hilly and steep and there are hardly any flat plains even along the rivers.

### **2.2. POPULATION**

The region is - as the middle hills of Nepal generally are - characterized by dense population, and thus progressive deforestation and erosion (KHADKA 1981). Besides a few variably compact villages, the houses are scattered over the flanks and ridges of the area. The population density is recorded with 84 per km<sup>2</sup>, with a per capita land holding of 0.04 ha (mean of Sindhupalchok and Dolakha District, Nepal District Profiles 1981).

A variety of people lives here. The dominating group are the Nepali-speaking Brahmin and Chhetri (over 50%) with their Hindu religion and caste system. They are followed by the Tamangs (over 20%), a Buddhist Tibeto-Burman-speaking community. Of the remaining groups the Newari (about 5%) should be mentioned (BISTA 1980, DOBREMEZ et al. 1974).

### **2.3. ECONOMY**

The economy depends almost exclusively on agronomy, which is, because of the variation in climate, diverse. An elaborate terrace-system with mainly rice, wheat, maize and mustard in the lower parts and potatoes,




Fig. 2.1. Nepal




Fig. 2.2. Situation of study area  
Abb. 2.2. Lage des Untersuchungsgebietes

maize, millet, barley and buckwheat in the higher areas can be seen, as can extensive pastures and shrubland or forests.

The whole region is easily accessible by the Arnico Highway, opened 1967 and leading from Kathmandu to Lhasa. It is one of the important trading routes to Tibet. The same highway is the gate to the whole Solu-Khumbu Region with its Sherpa community of Tibetan origin. This region, with Mount Everest as a magnet, is the destination of numerous mountaineering expeditions and trekkers.

A joint Nepali-Swiss Integrated Hill Development Project (IHDP) was started by the Swiss Technical Cooperation in 1974 with the aim of developing agronomy, education, forestry, health, small scale and cottage industry, and water supply and irrigation. The project entered its third phase, the handing-over, in 1985.

Also in 1974 and in connection with IHDP the Lamosangu-Jiri Road Project (LJRP) was started. From 1976-1985 a road of 110 km was built to connect the Arnico Highway with the Dolakha district (Fig. 2.2). In both projects much emphasis is laid on erosion control and protection of nature (NEVILLE 1985, SCHAFFNER 1985). The study in hand is part of this effort.

#### **2.4. GEOLOGY AND TECTONICS**

The Himalaya is a young tertiary mountain range still being uplifted. The most important structures of the present region are the WNW-ESE oriented folds and schuppen with the separating thrust faults. The rock can be divided into three major types:

- crystalline rocks (various gneisses)
- metamorphic sediments (phyllites and crystalline schists)
- quaternary deposits (river deposits).

The most important types of rock are sediments of a low metamorphic grade. These are usually exceptionally schistous to thin-foliated, while the gneisses are thickly bedded. Often, very deep weathering attributable to heavy cleavage and rock characteristics can be observed. This results in a general instability, in much broken and displaced rock units (LJRP 1977). This instability is aggravated by earthquakes which

occasionally occur (HAGEN 1950-1958 and 1969, DONNER 1972, MARUO et al. 1973).

In most places, a variably thick covering of loose rock conceals the bed. This is composed of detritus and scree. The particle size distribution of this loose rock varies significantly. However, the coarse components (gravel, stones, blocks) are typically embedded in a silty, fine sandy matrix. There is practically no clay (LJRP 1977).

## 2.5. CLIMATE

Nepal has a monsoon climate, with a pronounced rainfall maximum during the months of June to September.




Owing to the irregular nature of the midland hills, the uneven topography and the extreme differences in altitude there exist different climatic belts (DONNER 1972, DOBREMEZ 1974b):

---

Fig. 2.3 (p. 15). Climate diagrams of the study area  
(drawn by U. Schaffner, Emmenbrücke; IHDP-data)

Abb. 2.3 (S. 15). Klimadiagramme des Untersuchungsgebietes  
(gezeichnet von U. Schaffner, Emmenbrücke; IHDP-data)

- a - station
- b - height above sea level
- c - number of years of observation
- d - mean annual temperature (in degrees centigrade)
- e - mean annual precipitation (in millimeters)
- f - mean daily temperature minimum of the coldest month
- g - absolute minimum temperature (lowest recorded)
- h - mean daily temperature maximum of warmest month
- i - absolute maximum temperature (highest recorded)
- j - mean daily temperature fluctuation
- k - curve of mean monthly temperature (1 division =  $10^{\circ}\text{C}$ )
- l - curve of mean monthly precipitation (1 division = 20 mm, i.e.,  $10^{\circ}\text{C} = 20 \text{ mm}$ )
- m - period of relative drought (dotted) for the climate region concerned
- n - corresponding relatively humid season (vertical shading)
- o - mean monthly precipitation  $>100 \text{ mm}$  (scale reduced to 1/10) (black areas, perhumid season)
- r - months with absolute minimum below  $0^{\circ}\text{C}$  (diagonally shaded) i.e., with either late or early frosts



- hot monsoon belt (up to 1200 m)
- warm temperate monsoon belt (1200-2200 m)
- cool temperate monsoon belt (2200-3000 m)
- subarctic belt (3000-3800 m)
- arctic belt (from 3800 m).

IHDP has installed three meteorological stations. The rainfall figures recorded in the hills vary considerably according to the local topography. Bonch (2130 m a.s.l.; mean 1979-1984) has a yearly average of 3053 mm, Dandapakhar (1715 m a.s.l.; mean 1976-1984) 2830 mm and at Kirantichap (1240 m a.s.l.; mean 1981-1984) 1600 mm. 80-90% of the yearly rainfall occurs during the monsoon. There is a certain amount of rainfall during the winter months with possible snowfall above 2000 m from December to February (IHDP 1976-1985, Fig. 2.3).

The temperature varies mainly according to the altitude: Bonch shows a maximum mean of 18.5°C, Dandapakhar 21.8°C and Kirantichap 24.0°C. The minimum mean is 9.6°C at Bonch, 12.6°C at Dandapakhar and 13.8°C at Kirantichap. There can be frost in the coldest months from December to January above 1900 m (IHDP 1976-1985, Fig. 2.3).

## 2.6. VEGETATION AND SOIL


The vegetation is strongly connected with soil and climate, therefore we find a great variation in the Himalaya (SCHWEINFURTH 1957, STANTON 1972). A comparison of the vegetation zonation schemes of the Nepal Himalaya by different authors is given Fig. 2.4 (after OHSAWA 1974).

Owing to the high pressure of population the natural situation of the region has been considerably transformed: forests are rare. There is

---

Fig. 2.4 (p. 17). Comparisons of vegetation zonation schemes in the Nepal Himalaya (after OHSAWA 1977)  
Abb. 2.4 (S. 17). Vergleich verschiedener Darstellungen von Vegetationszonen im Nepal-Himalaya (nach OHSAWA 1977)





hardly any forest below 1900 m except in inaccessible areas or for cultural reasons or property conditions.

According to ESPINOSA (1974), the soils of the Lamosangu-Kharidhunga area have developed from metamorphic and sedimentary rocks, chiefly augengneiss, phyllite and metamorphosed sandstone. The major soils on bench terraces are well to poorly drained, strongly to moderately acid, slightly gravelly loams, silt loams and clay loams. Where forested, the soils range from excessively drained, stony sandy loams to well drained silty clay loams. Soil reaction is strongly to very strongly acid. The soils are mainly Cambisols, Acrisols and Regosols; a few Gleysols (classification by FAO-UNESCO 1974) occur along the rivers (ESPINOSA 1974, DOBREMEZ 1976).