

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 94 (1989)

Artikel: The impact of draining, burning and fertilizer treatments on the nutrient status of floating "Typha" mats in a freshwater marsh = Der Einfluss der Drainage, kontrolliertem Abbrennen und Düngung auf die Nährstoffverhältnisse in schwimmenden "Typha"-Beständen

Autor: Krüsi, Bertil O.

Kapitel: 1: Introduction

DOI: <https://doi.org/10.5169/seals-308910>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

1. INTRODUCTION

In cool, northern wetlands nutrients tend to be withdrawn from circulation through incorporation into organic matter and surface litter which accumulates because of slow decomposition in this type of climate (e.g. WEIN 1983). ROWE and SCOTTER (1973) called this process nutrient lock-up. WARD (1968) reported for wetland areas where water levels had been artificially stabilized that organic matter can accumulate to substantial depths in comparatively short periods of time that is 10 to 20 years. And WELLER (1975) found in a newly established cattail stand the formation of a 35 to 43 cm thick mat in as little as 3 years. Unless there is a continuous supply of nutrients, the lock-up process must lead to a decrease in available minerals. WHITMAN (1974, 1976) observed this process in water level stabilized marshes similar to the one studied in the present paper, where in a relatively short period of time after impounding and flooding the amount of available nutrients declined. As was shown by BEAUCHAMP and KEREKES (1980), newly flooded impoundments tend initially to be limited by nitrogen and later on by phosphorus.

In areas where in more or less regular intervals drought periods occur, freshwater marshes tend to undergo cyclic changes (e.g. WELLER and SPATCHER 1965, WELLER 1978, 1982, VAN DER VALK and DAVIS 1978a,b, 1979, VAN DER VALK 1981). During wet periods nutrients are removed from circulation, accumulated and locked-up in surface litter and soil organic matter, during dry periods they are released through decomposition. Due to recurrent drought, prairie marshes in North America are periodically rejuvenated, a complete cycle requiring from 5 to 35 or more years (VAN DER VALK and DAVIS 1978a). WELLER (1982) argues that natural fires must have occurred regularly in the prairie eliminating the bulk of accumulated plant material, returning locked-up nutrients into circulation and rejuvenating the marsh in a way very similar to that of drought. Whereas drought periods in cool northern ecosystems are too limited in duration and too infrequent to reverse the lock-up process, recent studies (e.g. TOLONEN 1983) indicate that fire has always been an essential ecological factor in northern latitudes. The marsh studied proved to burn anytime of the year provided the weather was sunny and windy since there was a sufficient quantity of contiguous fuel (surface litter) to permit fire

spread even among green Typha stems (KRÜSI and WEIN 1988).

The hypothesis that ageing ecosystems can be revitalized through improved nutrient cycling conditions brought about by fire has been termed the "paludification-fire-nutrient release hypothesis" (WEIN 1983), "paludification" or "swamping" being the term proposed by HEILMANN (1966) for the species change associated with the nutrient lock-up process. Up to date, quantitative information and experimental studies concerning the paludification-fire-nutrient release hypothesis are rare. Moreover, there is conflicting opinion as to the degree of revitalization and as to whether paludification is enhanced in moist ecosystems (e.g. HEINSELMANN 1975).

The objective of the present study was to test the paludification-fire-nutrient release hypothesis for a Typha glauca floating mat in a water-level stabilized marsh and to compare the effectiveness of burning with that of draining. In order to get a more precise idea of the extent to which draining and burning affect the nutrient status of the system studied, their impact was compared with that of the application of known amounts of fertilizers viz. (i) nitrogen, (ii) phosphorus, (iii) lime and (iv) nitrogen, phosphorus and lime combined. Burning has been reported to have similar effects on plant growth as liming (FOWELLS and STEPHENSON 1934 cited in RAISON 1979) or as fertilizing with nitrogen and phosphorus (e.g. BURTON 1944, RAISON 1979). Treatment effects were evaluated in terms of phenological and growth characteristics of Typha glauca.

Acknowledgments

It is a pleasure to acknowledge the help of a large number of people. I am grateful to Prof. Dr. E. Landolt, Swiss Federal Institute of Technology, Zurich, who supported my application for a postdoctoral fellowship in Canada. Special thanks are due to Dr. Ross W. Wein, Director of the Fire Science Center of the University of New Brunswick in Fredericton, Canada, and his staff who received me cordially and encouraged me to undertake this study. Janice M. Moore and Katherine J. Turner assisted in the field and laboratory. Scott Whitman provided considerable support as summer student. Timothy G. Dilworth, Mohamed El-Bayoumi and Peter Thomas helped when burning was conducted. The National Wildlife Area Committee permitted the study to be carried out on the Tintamarre National Wildlife Area. Mr. and Mrs. Walter Phinney and Mr. Larry Phinney were our neighbours at the research site and helped us kindly in many ways. Sincere thanks of the author are addressed to all these persons as well as

to all the other colleagues of the University of New Brunswick, Fredericton N.B., and the Université de Moncton, Moncton N.B., who helped occasionally in the marsh. The advice of Dr. George P. Malanson, Iowa City, who read an early draft of the paper is greatly appreciated.

The Postdoctoral Fellowship awarded by the University of New Brunswick as well as the operating funds provided by the World Wildlife Fund (Canada), the Canadian Sportsmen's Fund, Ducks Unlimited (Canada), the Canadian Wildlife Service, the Natural Sciences and Engineering Research Council of Canada and the University of New Brunswick Research Fund are gratefully acknowledged. Last but not least, statistical analysis of the data would not have been possible without the permission to use the computing facilities at the Centre National Universitaire Sud de Calcul, Montpellier, France, generously granted by the Module "Dynamique et Succession" of the Centre Louis Emberger, Centre National de la Recherche Scientifique, during a stage in 1984 at Montpellier, France.