

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 105 (1991)

Artikel: Pflanzenökologische und limnologische Untersuchungen des Reussdelta-Gebietes (Kanton Uri) : Aufnahme des Ist-Zustandes von 1987/88 = Phytoecological and limnological investigations in the region on the Reuss delta (canton Uri)

Autor: Elber, Fredy / Marti, Karin / Niederberger, Klemens

Register: List of tables :limnology

DOI: <https://doi.org/10.5169/seals-308914>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Sector 1: on the right of the River Reuss, Allmeini and Schützenrütli
Sector 2: on the left of the River Reuss, Flüelerschachen and Schanz
Sector 3: on the left of the River Reuss, Schwäb and Seedorferbucht. 188

7.2.-

7.7. Comparison of the vegetation area (size and flora composition) in the littoral 1982/83 and 1988.

Sector 1 = Allmeini and Schützenrütli

Sector 2 = Flüelerschachen and Schanz

Sector 3 = Schwäb and Seedorferbucht

----- = 5 m isobath (boundary of the actual shallow water zone with regard to the mean water level of many years, altitude above sea level: 433.6 m.

Investigation from 1982/83: x = occurrence of the species without declaration of the relative frequency .

Investigation from 1988: x = occurrence of one or a small number of

191-197

8.1. Density of individuals for *Tabellaria fenestrata*, *Rhodomonas minura* and the group of the diverse flagellates in August (before the flood event) and in September (after the flood event) at the sampling stations in the littoral and in the pelagial. Pel li = pelagial on the left of River Reuss mouth, Pel re = pelagial on the right of River Reuss mouth. 205

9.1. Possible development of the phytoplankton community in the littoral of Lake Uri as a result of the new formation of the River Reuss mouth. 211

9.2. Possible development of the macroinvertebrate communities in the littoral of Lake Uri as a result of the new formation of the River Reuss mouth. 214

List of tables / Limnology

2.1. Latitude and longitude of the sampling stations in the pelagial and the littoral. Latitude and longitude refers to the contour map 1:5000, Reusshochwasser, Canton Uri. 79

2.2. Dates of the chemical, physical and biological investigations and exposition time of the glass slides. 81

3.1. Annual mean and standard deviation of the investigated parameters during the period April 1987 to April 1988. M = mean, S = standard deviation. 86

3.2. Results of the statistical comparisons between the six sampling stations in the littoral and between the two in the pelagial as well as between the littoral and the pelagial. Significance level $p \leq 0.05$, * = $p > 0.1$. 98

3.3. Results of the bacteriological investigation from August 5, 1987 (carried out by the Labor der Urkantone) and February 1, 1988 (our investigation). The investigation were carried out in the Giessen (a channel which acts as a receiving body of water for the water treatment plant of Altdorf), within the bay behind the island for the birds and to the left of the island.
* = at the mouth of Giessen, ** = about 50 m upstream of the Giessen mouth, *** = on the right to the Giessen mouth, near the shore. 102

3.4. Comparison of the nutrient concentrations of some Swiss lakes. Values of Lake Uri from January to March 1988. * = Mean of the littoral stations

(average concentration from the 0-4 m depth interval), ** = Mean of the pelagial stations (Average concentration from the 0-10 m depth interval). Data from Lake Zuerich: Seenuntersuchungen der Wasserversorgung (Water Supply Service) Zürich (1988), from the other lakes: Bulletin des Bundesamtes für Umweltschutz (1984-1988). V = Vierwaldstättersee, Lake Lucerne.

103

- 4.1. Phytoplankters and their volumina. Investigation period April 1987 to April 1988.
X = net plankton. Data of phytoplankter's volumina from BÜRGI (1977), BÜRGI and LEHN (1979), FRIEDLI (1978), Water Supply Service of Zuerich and our measurements. 109-111
- 4.2. Minimum, mean and maximum of individual density, wet weight, chlorophyll a concentration, dry weight and ash-free dry weight at the six sampling stations in the littoral (mixed water, water column from the 0-4 m depth interval) and at the two sampling sites in the pelagial (0-10 m depth interval, mean of samples taken at 0, 1, 2.5, 5 and 10 m depth) during the investigation period from April 1987 to April 1988. Pel-l = pelagial on the left, Pel-r = pelagial on the right, Min. = minimum, Mittel. = mean, Max. = maximum. 113
- 4.3. List of zooplankters found in Lake Uri during the investigation period April 1987 to April 1988. 123
- 5.1. Mean chlorophyll a concentration (values in $\mu\text{g}/\text{cm}^2$) of the periphyton on the glass slides at 2.5 m and 5 m depths and ratio of the values of the two depths during the investigation period 1987/88. 129
- 5.2. Mean ash-free dry weight (values in mg/cm^2) of the periphyton on the glass slides at 2.5 m and 5 m depths and ratio of the values of the two depths during the investigation period 1987/88. 131
- 5.3. Annual sum of the biomass, mean daily growth rate (measured as ash-free dry weight, values in g/m^2) of the periphyton and ratio of the values of the two depths during the investigation period 1987/88. 131
- 5.4. Mean portion of chlorophyll a (values in %) at the ash-free dry weight of the periphyton on the glass slides at 2.5 m and 5 m depths and ratio of the values of the two depths during the investigation period 1987/88. 136
- 5.5. Mean degree of cover of the periphyton on the glass slides (values in %) at 2.5 m and 5 m depths during the investigation period 1987/88. 137
- 5.6. List of algae (excluding the diatoms) found on the glass slides at 2.5 m and 5 m depths during the investigation period 1987/88. 138
- 5.7. List of diatoms found on the glass slides at 2.5 m and 5 m depths during the investigation period 1987/88 and the index of sensitivity. x = species found at the corresponding sampling station, bold = major species. Index of sensitivity: 1 = very sensitive, 2 = sensitive, 3 = less tolerant, 4 = tolerant species (see Chapter 5.4). 141-146
- 5.8. Total, mean, maximum and minimum species numbers of the diatom community at the six sampling stations in the littoral during the investigation period May 1987 to April 1988. Constant species = species which occur with at least 1% relative frequency during the entire investigation period.
* = smallest number of species of all samplings in September 1987, after the flood event. 146
- 5.9. Mean relative frequency of the major diatom species (values in %) at the six sampling stations in the littoral during the investigation period May 1987 to April 1988. Decreasing order with regard to the average relative frequency. 150

5.10.	Maximum, minimum and mean diversity (values in bit/ind.) of the diatom communities at 2.5 m and 5 m depths during the investigation period May 1987 to April 1988. * = lowest diversity of all samplings in September 1987, after the flood event.	152
5.11.	Statistical comparison of the six sampling stations with regard to periphyton biomass, number of species, diversity of the diatom community and the relative frequencies of the 18 major species. Friedmann Test, significance level 5%. * = $p > 0.05$. H = stations with high mean range, T = stations with low mean range. Species listed in order of the frequency of significant differences.	154
5.12.	Statistical comparison of the two depths, 2.5 m and 5 m, with regard to periphyton biomass, number of species, diversity of the diatom community and the relative frequencies of the 18 major species. Wilcoxon matched pairs test, significance level 5%. * = $p > 0.05$. Species listed in order of the frequency of significant differences.	155
5.13.	Water quality with percent of the diatom differential groups. üs = very sensitive, s = sensitive, wt = less tolerant, t = tolerant species. * = after HOFMANN (1987), other classifications after KRAMMER and LANGE-BERTALOT (1986).	157
6.1.	Mean sedimentation rate (measured as dry weight, TG, and ash-free dry weight, AFTG) at the six sampling stations in the littoral during the investigation period May 1987 to April 1988. Number of samplings (N) = 12, station 6: N = 11. Without the sampling from September: N = 11, station 6 N = 10.	164
6.2.	List of macroinvertebrates found in the littoral of Lake Uri during the investigation period 1987/88.	167-168
6.3.	Mean number of individuals per labyrinth, total and mean number of species of macroinvertebrates at the six sampling stations during the investigation period 1987/88.	170
6.4.	Mean number of individuals per labyrinth of the main macroinvertebrates species or species groups at the six sampling stations during the investigation period 1987/88. N = 12.	171
7.1.	List of aquatic plants found in the littoral of Lake Uri. Data from the investigation 1982/82 by LACHAVANNE et al. (1985). a = species which are protected or should be protected in Canton Uri, b = endangered and nearly extinct species, c = species rare in Lake Lucerne, x = occurrence cartographically recorded, + = occurrence mentioned, cartographically not recorded. I = groups of indicators after LANG (1981): 1 = Indicator of oligotrophic to little mesotrophic places, 2 = Indicator of mesotrophic to little eutrophic places, 3 = Indicator of eutrophic to very eutrophic places.	186