

**Zeitschrift:** Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich  
**Herausgeber:** Geobotanisches Institut, Stiftung Rübel (Zürich)  
**Band:** 107 (1992)  
  
**Artikel:** The "Wolosate" peat bog reserve in the Bieszczady Mts. (S.E. Poland)  
**Autor:** Ralska-Jasiewiczowa, Magdalena  
**DOI:** <https://doi.org/10.5169/seals-308958>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 18.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## The "Wołosate" peat bog reserve in the Bieszczady Mts. (S.E. Poland)

Magdalena RALSKA-JASIEWICZOWA

### 1. SITUATION, VEGETATION AND STRATIGRAPHY

The peat bog is situated in the upper part of the Wołosatka river valley (Fig. 1), about 3 km SSE of the village, Ustrzyki Górne. The peat bog lies at 700 m a.s.l., and is fairly regularly elliptic in shape, measuring about 120 by 260 m. Its dome is raised about 3.20 m above the surrounding area (LIPKA and GODZIEMBA-CZYŻ 1970). The surface of the peat bog was once cut with drainage ditches and was badly eroded. It is now protected as a nature reserve and there are signs that active growth is beginning to restore a natural surface. It is now a typical raised bog with a *Sphagnetum medii*-dominated surface and with *Ledum palustre*, *Oxycoccus quadripetalus*, *Oxycoccus microcarpus*, *Andromeda polifolia*, *Eriophorum vaginatum*, *Carex pauciflora*, *Drosera rotundifolia*, *Vaccinium uliginosum*, *Vaccinium myrtillus* and *Vaccinium vitis-idaea*, and the mosses, *Sphagnum medium*, *Sphagnum acutifolium*, *Sphagnum papillosum*, *Campylopus flexuosus*, *Dicranodontium denudatum*, growing in it. Sparse shrubby birches (*Betula pendula*), single non-flowering specimens of *Picea abies* and sporadic *Pinus sylvestris* seedlings occur on its surface. *Carex fusca*, *Carex brizoides*, *Potentilla erecta*, *Juncus effusus*, *Menyanthes trifoliata*, *Sphagnum recurvum* and *Polytrichum commune* grow in the lagg of the bog.

The lower terrace of the Wołosatka river supports *Alnus incana* scrubs. The valley floor is covered by degraded *Deschampsietum* and *Nardetum* mead-




Fig. 1. The location of the "Wolosate" peat bog.

ows with *Juniperus communis*, and fallows overgrown by *Alnus incana* showing a mixture of succession stages typical of ground abandoned by man. Now fields are being cultivated again. The slopes of the valley are occupied by beech forests without fir and spruce. LIPKA and GODZIEMBA-CZYŻ (1970) reconstructed the stratigraphy of the bog. Details are given in Fig. 2.




Fig. 2. Stratigraphy of the "Wolosate" peat bog shown by the NWW-SEE transect. The bore sites at 25 m intervals are indicated. (LIPKA and GODZIEMBA-CZYŻ 1970, modified).  
1 - *Sphagnum* peat, 2 - *Eriophorum vaginatum* peat, 3 - wood peat, 4 - clay.

## 2. THE SHORT HISTORY OF VEGETATION AND HUMAN IMPACT

### 2.1. DESCRIPTION OF THE DIAGRAM

Wolosate is the youngest of the peat bogs in the Bieszczady Mts. examined by RALSKA-JASIEWICZOWA (1980). Peat accumulation started there during the declining phase of mixed deciduous forests (climatic optimum). The pollen diagram (Fig. 3) covers only the latest period of forest development and can only with difficulty be divided into the following pollen assemblage zones (PAZ):

- 380-365 cm: *Corylus-Tilia-Ulmus-Picea* PAZ characterized by the falls of *Corylus*, *Ulmus* and *Tilia*, and a rapid rise of *Carpinus* and *Fagus* pollen curves. The *Picea* pollen curve is high. The indicators of man presence appear sporadically.
- 365-315 cm: *Fagus-Carpinus* PAZ. *Carpinus* and *Fagus* become the dominant pollen taxa, *Picea* still shows rather high frequencies and the indicators of human activities form the continuous pollen curves.
- 315-190 cm: *Fagus-Abies-Betula* PAZ. *Carpinus* pollen values decline and those of *Abies* become substantial. *Fagus* pollen curve shows some depression, and *Betula* and the frequencies of culture indicators increase.
- 190-0 cm: *Fagus-Abies-NAP* PAZ. Oscillations in pollen curves of dominant trees are evidently correlated with those of culture indicators, their sum in places, rising up to 25% of the total pollen sum. Therefore, it seems reasonable to describe the anthropogenic changes recorded in the pollen diagram independently from the regional forest history.

The pollen assemblage zones (PAZ) reveal the following development:

1. Since 370 cm: the beginning of continuous *Artemisia*, *Chenopodiaceae* and *Plantago lanceolata* pollen curves.
2. Since 345 cm: the slight rise of *Gramineae*, *Plantago lanceolata*, the beginning of *Rumex* pollen curve and sporadic appearances of *Plantago media* pollen are correlated with the decline of *Carpinus* and *Corylus*, and first rise of *Betula* since around 315 cm. They are followed by the declining tendencies of *Fagus*, *Picea*, a maximum of *Artemisia* and the first appearance of the *Humulus/Cannabis* pollen type. The maximum of *Betula* and of *Gramineae* pollen is correlated with the minimum of the *Fagus* curve at 270 cm. Since 280 cm the first pollen grains of cereals (*Triticum* type) appear. The decreased pollen frequencies of culture indicators between 220 and 195 cm correspond with the subsequent maximum of *Fagus* pollen.
3. At about 190 cm, the above mentioned culture indicators increase again along with the increase in cereals (including *Secale*!) and the beginning of the regular appearances of *Urtica*, *Polygonum aviculare* and *Plantago major* pollen. The *Picea* pollen values fall to a minimum followed by a transitory depression in the *Fagus* curve, and a rise in frequencies of *Abies*, *Betula*, *Corylus* and *Alnus* pollen. Around 135 cm the pollen values of some culture plants decrease again. The curves of *Cerealia* undiff. and *Triticum* type become discontinuous; *Secale* pollen occur only as single grains. This decline is short-lived in the cases of *Artemisia*, *Plantago lanceolata* and *Plantago major*.

4. The youngest settlement phase, covering the upper 80 cm of the profile, consists of two subphases divided by a more or less distinct depression in most pollen curves of culture indicators between 45 and 25 cm. It is characterized by double peaks of nearly all indicators mentioned so far and by the appearance of *Centaurea cyanus* and *Trifolium pratense* pollen and in the younger subphase, pollen of *Fagopyrum*. In this subphase the final decline of *Carpinus*, *Abies* and *Fagus* and a rise of *Alnus* and *Corylus* pollen curves takes place.




Fig. 3. Pollen diagram from the "Wolosate" peat bog.

## 2.2. INTERPRETATION

The profile from the Wolosate peat bog has not been radiocarbon dated and its interpretation is based on comparisons with other dated profiles. The decline of *Ulmus* and *Corylus* coinciding with the rise of *Carpinus* and *Fagus* pollen curves - the phenomenon which defines the basal *Corylus-Tilia-Ulmus-Picea* pollen zone at Wolosate - have been dated in pollen diagrams from the Smerek and Tarnawa Wyzna peat bogs at c. 4'400-4'300 B.P. (RALSKA-JASIEWICZOWA 1980). These dates set the beginning of *Carpinus* and *Fagus* expansion in the whole region of the Polish East Carpathians, opening the development processes of modern forest zones in this area (*Carpinus-Fagus* PAZ). Their formation was completed by the expansion of *Abies* (*Fagus-Abies-Betula* PAZ) dated at Smerek and Tarnawa Wyzna at c. 2'800-2'700 B.P.

An interesting, individual feature of the Wolosate pollen diagram is the record of prehistoric human activities in the surroundings of the peat bog. It may be interpreted as follows:

1. The oldest traces of man, coinciding with a fall of *Ulmus* and an increase of *Fagus* pollen values, date probably from the late Neolithic. They might have been produced by the tribes of the Corded Ware Culture, which depended mostly on animal husbandry, practicing clearing by fire and axe to increase the area of pastures, but also grazing their cattle within forests. Their settlements might have been seasonal.
2. The next phase may correspond to the Late Bronze/Early Iron Age Lusatian Phase at Tarnawa Wyzna and Smerek. The subsequent changes reflect probably;
  - a. the clearances in deciduous forests of the lower elevations and in spruce forests, the development of pastures (increase in grass pollen) and of secondary birch copses;
  - b. the extension of forest clearances up to the beech forests, the encroachment of fir and the progressive development of post-farming communities (hazel and alder);
  - c. the further increase of pasture areas at the cost of beech-fir and spruce-fir forests and the first evidence of agriculture. The settlement was presumably situated at a moderate distance from the peat bog.
3. The third cycle of changes may correspond to phase 3 at Tarnawa Wyzna, referred to the period of Roman influences. It is very clearly expressed in the Wolosate diagram by peaks of nearly all culture indicators. They point to the presence of a settlement surrounded by pastures and arable fields in the Wolosatka river area. It may have been connected with the presence of a trade route, running along the broad and easily passable valley of the Wolosatka and through the Beskid Pass towards the valley of the river Uh, although the existence of such a route has not, as yet, been confirmed archaeologically. It may have been another branch of the route detected by REYMAN (1960) running on the southern side of the Carpathians along the valley of the Latorica or Uh towards the Przelecz Uzocka pass.

The following section of diagram represents a period of a partial reduction in human activities. The disappearance of pollen of autogamous cereals indicates the end of farming near the peat bog (there are still traces of anemophilous rye). Presumably the settlement was abandoned, but the route continued to be used and the areas adjacent to it were grazed.

4. The youngest, historical period of colonization also showed two phases. The depression of human indicators, separating the two subphases, may have been due to the economic standstill at the time preceding the Wallachian migrations and the foundation of villages in the Late Medieval Period.

## REFERENCES

- LIPKA K. and GODZIEMBA-CZYŻ W., 1970: Peat bogs and fens of the Wolosatka stream watershed in the Western Bieszczady Mts. (In Polish). *Zes.Nauk.WSR Krakow, Melioracje* 59(4), 3-28.
- RALSKA-JASIEWICZOWA M., 1980: Late-Glacial and Holocene vegetation of the Bieszczady Mts. (Polish Eastern Carpathians). PWN, Warszawa/ Krakow. 202 p.
- REYMAN J., 1960: Réflexions sur le rôle du Col d'Uzok mises en lumière par les trouvailles des monnaies romaines. (In Polish with French summary). *Acta Archeol.Carp.* 1(1), 55-59.

Address of the author: Prof. Dr. hab. Magdalena RALSKA-JASIEWICZOWA  
W. Szafer Institute of Botany  
Polish Academy of Sciences  
Lubicz 46  
31-512 Krakow, Poland