Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech.

Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 130 (2002)

Artikel: Alpine vegetation of the Teberda Reserve, the northwestern Caucasus

= Die Alpine Vegetation des Teberda Reservates, Nordwest-Kaukasus

Autor: Onipchenko, V.G.

Kapitel: 1: Introduction

DOI: https://doi.org/10.5169/seals-308994

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 04.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

1. Introduction

High elevations are home to unique assemblages of plants and animals. Although the overall economic importance of alpine and subalpine ecosystems has probably never been particularly great, high mountains remain an integral and important part of the biosphere. Admittedly, this is one of the most fragile and poorly studied ecosystems in the world (BLISS 1985, WALTER & BRECKLE 1986, ARCHIBOLD 1995). Alpine tundras and subalpine meadows, grasslands and scrub communities are used as important grazing areas worldwide. The great aesthetic quality and solitude of alpine ridges attracts multitudes of tourists to the high-mountain wilderness areas (Lukschanderl 1983).

The Teberda State Biosphere Nature Reserve (Teberdinskii Zapovednik) is one of the largest in the Caucasus, covering 85'000 ha. It has played an extremely important role in keeping biological diversity of thousands of species. The flora of the reserve is well-studied (Vorob'eva & Kononov 1991). The vegetation, on the other hand, has not yet been well documented (Korotkov 1994). This hampers conservation work in the region, since the preserve remains essentially the only "baseline" pristine area undamaged by grazing and other human activities in Karachaevo-Cherkessian Republic.

The main aim of our study was to describe the vegetation of the reserve in terms of its syntaxonomy. Our interests were focused mainly on the alpine (non-forest) vegetation, but some transitional to forest communities have also been included.

The present volume describes, for the first time in English, all principal types of alpine and subalpine plant communities of the Teberda Reserve. The monograph is intended as a handy reference for plant ecologists, park rangers, and amateur naturalists interested in studying plant ecology in the Northern Caucasus – a land of unsurpassed beauty and remarkable biological diversity.

Among different approaches to studying terrestrial ecosystem diversity, floristic classification (syntaxonomy) is the most widespread in Europe. We used this method of community classification to characterize communities since it has long ago become a common language for plant ecologists in many countries of the world. Position of a community within the syntaxonomic system may serve as its "address" when more detailed investigations take place. Syntaxonomic "reviews" are very important in national parks and nature

reserves, and in poorly studied vegetation areas as a first step in detailed vegetation surveys.

The syntaxonomic (floristic) classification was at first developed by Braun-Blanquet and colleagues early in this century (BRAUN-BLANQUET 1921). It very much resembles the conventional biological classification of living organisms. The highest level of classification is represented by classes, subdivided into orders. The latter are further subdivided into alliances (unions) and then into particular associations. Once classifications are published, scientists and land use managers can both use them as a common reference point.

The plant releves for this volume were collected over 16 years (1980-1995). Due to similar ecological conditions and vegetation structure of the Alps and the Caucasus, we chose to adjust the well-developed European system of higher syntaxa to our conditions. Regional specificity was reflected mostly at the level of alliance or association. Such an approach seems to allow comparison of vegetation structure between large regions.