

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich
Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)
Band: 130 (2002)

Artikel: Alpine vegetation of the Teberda Reserve, the northwestern Caucasus
= Die Alpine Vegetation des Teberda Reservates, Nordwest-Kaukasus
Autor: Onipchenko, V.G.
Kapitel: 8: Alpine heaths : "Juncetea trifidi"
DOI: <https://doi.org/10.5169/seals-308994>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 14.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

8. Alpine heaths – *Juncetea trifidi*

Prodromus

Juncetea trifidi HADAC 1946

Caricetalia curvulae BRAUN-BLANQUET in BRAUN-BLANQUET & JENNY 1926

Anemonion speciosae MINAEVA ex ONIPCHENKO all.nov.

Campanulo ciliatae-Chamaesciadietum acaulis MINAEVA &
ONIPCHENKO ass. nov.

Pediculari comosae-Eritrichietum caucasicum MINAEVA & ONIPCHENKO
ass. nov.

P.c.-E.c. typicum subass.nov.

P.c.-E.c. oxytropidetosum kubanensis MINAEVA & ONIPCHENKO
subass. nov.

P.c.-E.c. bromopsietosum variegatae subass.nov.

Anemonion speciosae

The alliance comprises alpine lichen heaths (alpine tundra) and fellfields on windward crests and slopes on acid soils (ONIPCHENKO *et al.* 1987). Diagnostic species of *Juncetea trifidi*, *Caricetalia curvulae* are represented by *Helictotrichon versicolor*, *Luzula spicata*, *Minuartia recurva*, *Trifolium polyphyllum* (geographical vicarious species of *Trifolium alpinum*), *Euphrasia ossica* (geographical vicarious species of *Euphrasia minima*) (OBERDORFER 1978; GRABHERR 1993b). Due to the severe ecological regime (deep winter freezing, frequent summer drought, thin soil profile, great day-to-night temperature fluctuations), the communities have low production rates, high species richness and significant cover of lichens. Floristically and ecologically they are remarkably similar to the communities of *Carici rupestris-Kobresietea bellardii*. The low frequency of *Kobresia spp.*, *Polygonum viviparum*, *Lloydia serotina* and *Draba scabra*, as well as a preference for warmer slopes (south and east aspect), distinguish *Anemonion* communities from the communities of the previous class (Table 8.1.). Their nomenclature type (typus) is *Pediculari comosae-Eritrichietum caucasicum*.

Table 8.1.
Diagnostic table of *Juncetea trifidi*

	1	2	3	4
<i>D.sp. Campanulo ciliatae-Chamaesciadetum acaulis</i>				
<i>Anthyllis vulneraria</i>	V	-	I	III
<i>Dactylina madreporiformis</i>	V	-	I	-
<i>Chamaesciadum acaule</i>	V	I	II	IV
<i>Primula algida</i>	V	I	II	IV
<i>Pulsatilla albana</i>	V	I	-	III
<i>Aetheopappus caucasicus</i>	V	III	-	III
<i>Campanula ciliata</i>	IV	I	I	-
<i>Gentiana aquatica</i>	IV	-	I	II
<i>Thymus nummularius</i>	III	-	-	-
<i>Jurinella moschus</i>	III	-	-	II
<i>Campanula saxifraga</i>	II	-	-	-
<i>D.sp. Pediculari comosae-Eritrichietum caucasici</i>				
<i>Gentiana pyrenaica</i>	I	V	IV	III
<i>Carum caucasicum</i>	II	V	IV	II
<i>Vaccinium vitis-idaea</i>	II	V	IV	IV
<i>Rhytidium rugosum</i>	-	V	II	IV
<i>D.sp. P.c.-E.c. oxytropidetosum kubanensis</i>				
<i>Oxytropis kubanensis</i>	II	V	-	IV
<i>Fritillaria lutea</i>	-	IV	-	III
<i>Plantago atrata</i>	II	IV	-	IV
<i>D.sp. P.c.-E.c. typicum</i>				
<i>Pedicularis caucasica</i>	II	-	IV	-
<i>Polytrichum juniperinum</i>	II	-	IV	II
<i>D.sp. P.c.-E.c. bromopsietosum varigatae</i>				
<i>Scabiosa caucasica</i>	I	III	-	V
<i>Bromopsis variegata (dom.)</i>	V	V	-	V
<i>Muscari racemosum</i>	-	-	-	IV
<i>Myosotis alpestris</i>	III	I	I	IV
<i>Silene saxatilis</i>	II	I	-	III
<i>Carex humilis</i>	-	-	-	III
<i>Polygala alpicola</i>	I	-	-	III
<i>Seseli alpinum</i>	I	-	-	III
<i>D.sp. Anemonion speciosae</i>				
<i>Arenaria lychnidea</i>	V	V	V	IV
<i>Carex sempervirens</i>	V	IV	V	I
<i>Carex umbrosa</i>	IV	V	V	III
<i>Eritrichium caucasicum</i>	V	V	V	II
<i>Minuartia circassica</i>	IV	V	V	V
<i>Anemone speciosa</i>	V	V	V	IV
<i>Pedicularis comosa</i>	V	V	V	V
<i>Thamnolia vermicularis</i>	V	V	V	I
<i>Cetraria nivalis</i>	V	V	II	I
<i>Cetraria cucullata</i>	V	III	III	-

Table 8.1. (continued)

	1	2	3	4
D.sp. <i>Caricetalia curvulae, Juncetea trifidi</i>				
<i>Trifolium polyphyllum</i>	III	V	II	III
<i>Euphrasia ossica</i>	IV	V	IV	III
<i>Luzula spicata</i>	III	V	V	II
<i>Helictotrichon versicolor</i>	V	V	V	V
<i>Minuartia recurva</i>	III	III	II	V
Other frequent species				
<i>Alchemilla caucasica</i>	IV	V	III	V
<i>Antennaria dioica</i>	IV	V	III	II
<i>Anthemis marshalliana</i>	V	V	II	V
<i>Aster alpinus</i>	V	IV	I	IV
<i>Campanula collina</i>	V	IV	I	IV
<i>Campanula tridentata</i>	IV	V	V	IV
<i>Cetraria islandica</i>	V	V	V	IV
<i>Cetraria laevigata</i>	IV	IV	I	-
<i>Cladonia gracilis</i>	III	V	III	-
<i>Cladonia mitis</i>	III	V	IV	I
<i>Cladonia pyxidata</i>	V	III	V	IV
<i>Erigeron alpinus</i>	V	V	III	II
<i>Festuca ovina</i>	V	V	V	V
<i>Gentiana biebersteinii</i>	V	II	II	V
<i>Gentiana verna</i>	III	II	II	IV
<i>Polygonum bistorta</i>	V	V	I	IV
<i>Ranunculus oreophilus</i>	IV	I	I	IV
<i>Veronica gentianoides</i>	V	V	IV	V

Syntaxa:

1 - *Campanulo ciliatae-Chamaesciadietum acaulis*, 2 - *Pediculari comosae - Eritrichietum caucasici oxytropidetosum kubanensis*, 3 - *P.c.-E.c. typicum*, 4 - *P.c.-E.c. bromopsietosum variegatae*

8.1. *Campanulo ciliatae-Chamaesciadietum acaulis*

Synonym: *Pediculari chroorrhynchae* - *Eritrichietum caucasici chamaesciadetosum acaule* MINAEVA 1987

Floristic features

The association includes communities on stabilized dry alpine talus (scree, fellfields). *Anthyllis vulneraria*, *Dactylina madreporeiformis*, *Chamaesciadum acaule*, *Pulsatilla albana*, *Aethopappus caucasicus*, *Campanula ciliata* (Table 8.2.), represent the diagnostic set. Due to substrate peculiarities, some common species of *Gypsophilion tenuifoliae* are rather frequent.

The floristic richness of this community type is high. We registered 80 vascular plant species, 12 bryophytes and 14 macrolichens in 9 relevels of the

association. Mean values per releve were 38, 1 and 8 species respectively. A more precise estimate of the species richness shows values of 4.5, 22.1 and 46.0 vascular plant species for 0.01, 1 and 100 square m respectively (ONIPCHENKO & SEMENOVA 1995). In comparison with the next association, the role of lichens is not great. Their cover ranges between 2% and 20% (with the mean of 10%).

Vascular plant cover is larger (15-50%, mean 32%). Typus, or nomenclature type, is releve No. 10/84.

Ecological features

These open communities on dry stabilized screes are typical of snowfree areas of the alpine zone (2700-2950 m a.s.l., mean 2820 m). They prefer steep (15-30°, mean 27°) slopes of "warm" aspect (south, east, southeast). There are many stones covering 3-70% (mean 40%). The high floristic richness is due to sparse plant cover and low competition between the plants. The soil seed bank of the communities is substantial (about 2800 germinable seed per square m) (SEMENOVA & ONIPCHENKO 1994). *Luzula spicata*, *Primula algida*, *Euphrasia ossica* are the main components of the bank, each containing more than 200 seeds /square m. According to their floristic composition and structure, we may consider the communities as seral, being gradually succeeded by the communities belonging to the next association.

8.2. *Pediculari comosae-Eritrichietum caucasici*

Synonym: *Pediculari chroorrhynchae-Eritrichietum caucasici* MINAEVA 1987
(*Pedicularis chroorrhyncha* Vved.=*P.comosa* L.)

Floristic features

This association combines closed winter-snowfree communities of the alpine and upper subalpine zones. The diagnostic species set is represented by several common alpine species (*Gentiana pyrenaica*, *Carum caucasicum*, *Vaccinium vitis-idaea*, *Rhytidium rugosum*). The main floristic features of the association consist of a low frequency (or absence) of *Campanulo ciliatae-Chamaesciadietum acaulis* - species and a good representation of *Anemonion speciosae* and *Juncetea trifidi* - species (Table 8.3.). The other characteristic property of the communities is an abundance of fruticose

lichens. Lichens cover from less than 1% to 55% (average 28%), while the same values for vascular plants and bryophytes are 15-80% (47%) and less than 1 - 25% (4%) respectively.

We registered 102 vascular plant species, 27 bryophytes and 20 lichens in 28 relevés of the association. Mean numbers per relevé were 31, 3, and 6 species respectively. Vascular plant species richness was estimated as 6.2, 25.0, and 39.8 species per 0.01, 1 and 100 square-m plots respectively (ONIPCHENKO & SEMENOVA 1995).

Three subassociations can be further distinguished according to their floristic composition and ecological properties:

P.c.-E.c. typicum (typus, or nomenclature type, No. 75/93) *Pedicularis caucasica* and *Polytrichum juniperinum* are the diagnostic species of the association. The diagnostic species of the other subassociations are practically absent. Floristic richness is relatively low (25 vascular plant species per relevé). The communities are more typical of the ridges of various aspects.

P.c.-E.c. oxytropidetosum kubanensis (Typus, or nomenclature type, No. 18/81) comprises floristically rich alpine lichen heaths with high frequency of *Oxytropis kubanensis*, *Fritillaria lutea*, *Plantago atrata*. They occupy mainly southern and eastern slopes.

P.c.-E.c. bromopsietosum variegatae (Typus, or nomenclature type, No. 103/95) pools low alpine and high subalpine variants of the association. The lichen cover here is the lowest (mean 7%), while the vascular plant cover is the highest (mean 65%) among the communities of the association. Some species of subalpine grasslands (*Scabiosa caucasica*, *Myosotis alpestris*, *Polygala alpicola*, *Seseli alpinum*) as well as steppe species (*Carex humilis*, *Muscari racemosa*) form the diagnostic set of the subassociation. The communities are well known in Russian publications as species-rich *Bromus variegatus* - grasslands (SHIFFERS 1953, VOROB'EVA 1977b, TANFIL'EV *et al.*, 1979).

Ecological features

The communities occupy windward crests and slopes within the altitude range of 2350-3000 m (mean 2710 m). They prefer moderate (2°-30°, mean 15°) slopes of southern and eastern aspects. As a rule, stone cover is not great (0-30%, mean 5%). The communities of the association have been the subjects of our long-term ecological investigation (RABOTNOV 1987, ONIPCHENKO 1994).

Table 8.2.

Campanulo ciliatae-Chamaesciadietum acaulis

Releve No.	5	4	28	32	10	6	42	13	55
Year	89	83	83	83	84	84	89	84	83
Altitude (* 10)	285	280	280	285	280	295	290	270	280
Steepness	30	30	15	30	30	15	30	25	35
Exposition	se	s	e	se	sse	se	se	s	e
Vascular plant cover	35	50	25	20	25	15	40	50	35
Bryophyte cover	0.5	1	0	0.5	0.5	0.5	10	3	0.5
Lichen cover	5	10	15	5	2	20	10	15	15
Stone cover	60	20	40	70	60	60	20	5	3
Lichen species number	7	8	10	7	6	8	7	12	7
Bryophytes species number	2	0	0	0	0	1	2	2	6
Vascular plants species number	54	38	29	38	37	34	37	34	40
D.sp. <i>Campanulo ciliatae-Chamaesciadietum acaulis</i>									
<i>Anthyllis vulneraria</i>	1	2	1	1	1	1	1	1	1
<i>Dactylina madreporiformis</i>	+	+	1	+	+	+	+	+	+
<i>Chamaesciadum acaule</i>	+	+	1	+	1	+	1	1	+
<i>Primula algida</i>	+		+	+	+	+	+	+	+
<i>Pulsatilla albana</i>	+	1	+		+	+	+	1	+
<i>Aetheopappus caucasicus</i>	1	1	+	+	1	+	1	1	
<i>Campanula ciliata</i>	+	+	+	1	1	1			+
<i>Gentiana aquatica</i>	+	+	+		+	+	+	+	
<i>Thymus nummularius</i>	+	+		+	1				+
<i>Jurinella moschus</i>		+		+	1		1	+	
<i>Campanula saxifraga</i>	+			+	1				
D.sp. <i>Anemonion speciosae</i>									
<i>Arenaria lychnidea</i>	+	1	1	1	1	1	1	1	+
<i>Carex sempervirens</i>	1		1	1	1	1	2	1	1
<i>Carex umbrosa</i>	+	+			1		+	1	1
<i>Eritrichium caucasicum</i>	+	+	+	+	+	+	+		+
<i>Minuartia circassica</i>		1	1	1	1	1	+		1
<i>Anemone speciosa</i>	1	1	+	+	+	1		1	+
<i>Pedicularis comosa</i>	1	1	+	+	+	+	+	+	+
<i>Thamnolia vermicularis</i>	+	1	1	+		1	1	1	1
<i>Cetaria nivalis</i>	+	+	1	+	+	1	+	1	
<i>Cetaria cucullata</i>	+	+	1	+	+	1		1	+
D.sp. <i>Caricetalia curvulae, Juncetea trifidi</i>									
<i>Trifolium polyphyllum</i>	+		1			1		1	1
<i>Euphrasia ossica</i>	+	+		+	+		1	+	+
<i>Luzula spicata</i>	+			1		+	+		+
<i>Helictotrichon versicolor</i>	1	+	1	+	+	+	+		+
<i>Minuartia recurva</i>	+				+		+		+
Other species									
<i>Alchemilla caucasica</i>		1	1		1		1	1	2
<i>Antennaria dioica</i>	1	+	1	+		+			2
<i>Anthemis marshalliana</i>	1	+	1	+	+	+	1	+	+
<i>Aster alpinus</i>	1	1	1	+	1	+	+	+	+
<i>Astragalus levieri</i>	+	+							+
<i>Bromopsis variegata (dom.)</i>	1	2	1	+	1		1	2	1
<i>Campanula collina</i>	1	+	+	+	+		1	1	1

Table 8.2. (continued)

Releve No.	5	4	28	32	10	6	42	13	55
Year	89	83	83	83	84	84	89	84	83
<i>Campanula tridentata</i>			1	+	+	1	1	+	1
<i>Carum caucasicum</i>	+			+		+			
<i>Cetraria islandica</i>	1	1	1	1	+	1	+	1	1
<i>Cetraria laevigata</i>	+	+	2	+	+	1			1
<i>Cladonia gracilis</i>		+	1				+	1	+
<i>Cladonia mitis</i>			+			1		1	1
<i>Cladonia pyxidata</i>	1	+	+	+	1	1	1	1	+
<i>Erigeron alpinus</i>	+	+	+	+	+	+	+	1	
<i>Festuca ovina</i>	1	1	1	1	+	1	1	1	1
<i>Gentiana biebersteinii</i>	+	+	+	+	+	+	+	+	+
<i>Gentiana septemfida</i>		+			+	+	+	1	
<i>Gentiana verna</i>	+					+	+	+	+
<i>Lloydia serotina</i>	+			+	+				
<i>Myosotis alpestris</i>	+	+			+				+
<i>Oxytropis kubanensis</i>	1		1			1		1	
<i>Plantago atrata</i>		1						1	+
<i>Polygonum bistorta</i>	1	1	+	+	+	+	+	+	
<i>Polytrichum juniperinum</i>	+					+			+
<i>Potentilla gelida</i>	1			+		+		+	
<i>Potentilla nivea</i>		+	+					+	
<i>Primula ruprechtii</i>	+	+							+
<i>Ranunculus oreophilus</i>	+	1		+	+		+	+	
<i>Vaccinium vitis-idaea</i>	2			1		1		1	
<i>Veronica gentianoides</i>	1	1	+	+	+	+	1	1	+

Sporadic species (number of releve in parenthesis, abundance are shownen after ":", unless it is not "+", Braun-Blanquet scale)

Alopecurus glacialis (42/89), *Anthemis cretica* (5/89), *Asperula alpina* (5/89, 10/84), *Bryoerythrophyllum recurvirostrum* (42/89), *Bryum* sp. (42/89), *Carum meifolium* (4/83:1), *Centaurea cheiranthifolia* (4/83, 42/89), *Cladonia foliacea* (13/84:1), *Cladonia furcata* (28/83), *Cornicularia muricata* (13/84:1), *Deschampsia flexuosa* (5/89), *Draba hispida* (42/89), *Draba siliquosa* (5/89), *Empetrum nigrum* (5/89), *Eurhynchium pulchellum* (55/83), *Festuca varia* (5/89), *Galium verum* (5/89), *Gentiana pyrenaica* (5/89), *Gnaphalium supinum* (5/89, 6/84), *Hieracium umbellatum* (55/83), *Hymenostomum microstomum* (13/84), *Juniperus communis* (5/89, 55/83), *Kobresia schoenoides* (32/83:1), *Minuartia aizoides* (5/89:1), *Mnium ambiguum* (55/83), *Parmelia vagans* (42/89, 13/84:1), *Pedicularis caucasica* (32/83, 6/84), *Physconia muscigena* (13/84), *Poa badensis* (4/83), *Pohlia cruda* (55/83), *Polygala alpicola* (42/89:1, 55/83), *Polytrichum piliferum* (5/89, 55/83), *Potentilla crantzii* (55/83), *Sanionia uncinata* (55/83), *Saxifraga flagellaris* (32/83, 6/84), *Saxifraga juniperifolia* (32/83), *Saxifraga moschata* (55/83), *Scabiosa caucasica* (55/83), *Scorzonera cana* (5/89), *Sedum tenellum* (5/89), *Seseli alpinum* (55/83), *Silene saxatilis* (10/84, 42/89:1), *Taraxacum confusum* (42/89), *Tortella tortuosa* (13/84:1), *Valeriana alpestris* (5/89, 32/83), *Viola altaica* (4/83, 55/83).

Date (day.month), size (sq.m) and location of the releves.

5/89 - 09.08, 100, M.Khatipara; 4/83 - 20.08, 15, M.Khatipara (G.Levitskaya); 28/83 - 31.08, 15, M.Khatipara (A.Baykalova); 32/83 - 31.08, 16, M.Khatipara (A.Baykalova); 10/84 - 01.09, 25, M.Khatipara; 6/84 - 25.08, 25, M.Khatipara; 42/89 - 30.08, 25, Bol.Khatipara; 13/84 - 01.09, 25, M.Khatipara; 55/83 - 10.09, 25, Mukhu

A short account on the research results is presented below.

The average composition of the aboveground **biomass** in this community is the following: vascular plants - 113, lichen - 440, mosses - 3, litter and dead plant parts - 230 g/square m d.w. Belowground biomass and dead matter were estimated as 480 and 400 g/sq. m d.w. respectively. Total annual net-production is low (about 150 g/sq.m d.w) (ONIPCHENKO 1985).

The communities have homogeneous fine **spatial structure** (ONIPCHENKO & POKARZHEVSKAYA 1994). This structure is very peculiar: small lichen patches alternate with graminoid bunches or isolated shoots of vascular plants. The following hypothesis was put forward to explain such structure (ONIPCHENKO 1985, 1994, GRABHERR 1989): in poor shallow soils, the roots of vascular plants occupy a larger area than their aboveground shoots. Thus, vacant space above ground becomes available for the fruticose lichens, since they compete very little for nutrients with the vascular plants. Results of a special series of long-term experiments confirmed this hypothesis.

The high **floristic richness** may probably be attributed to a long period of stable existence of the communities, which was demonstrated both by spore and pollen, and phytolith soil analyses (PAVLOVA & ONIPCHENKO 1992, BLINNIKOV 1994). It is interesting to note that floristic richness of soil and epiphytic algae was also very high in this community (SHTINA *et al.* 1995).

Long term experiments of reciprocal sod transplantations between pairs of different alpine communities showed that most species of the alpine heath could successfully grow under more favourable conditions, but that they are excluded from the grasslands due to competition. On the other hand, dominants of *Nardetalia* - grasslands cannot survive the abiotic stress of the harsh alpine heaths' environment (SENNOV & ONIPCHENKO 1994).

Soil **seed bank** is relatively small (about 350 seed/ sq.m). Species producing many seeds (*Euphrasia ossica*, *Primula algida*, *Gentiana pyrenaica*, *Veronica gentianoides*) are well represented in the bank (SEMENOVA & ONIPCHENKO 1994). Due to the generally very severe environment, the recovery process after disturbance of alpine heaths is very slow. For example, revegetation after mild digging by wild boar on a single occasion (*Sus scrofa L.*) took at least 15 years (ONIPCHENKO & GOLIKOV 1996). The communities serve as winter pastures for wild ungulates. Lichen cover degradation takes place in overgrazed areas. Some special protection (restricted grazing of livestock) is absolutely necessary for the preservation of these fragile and species-rich communities, which have such exceptional aesthetic value.

Table 8.3. *Pediculari comosae-Eritrichietum caucasici*

Table 8.3. (continued)

Table 8.3. (continued)
Releve No.

Table 8.3. (continued)

Scandinavian species number of subspecies in monotypic genera are shown after "in natus" unless it is not $n = 1$. Because Diomus and

Sporadic species (number of relevés in parentheses, abundance are shown after "...", unless it is not ...+, Braun-Blanquet scale)

Aconitum nasutum (51/93:r), *Androsace albana* (12/84), *Ajuga orientalis* (12/84), *Bartsia alpina* (1/89), *Barbilophozia lycopodioides* (159/94), *Botrychium lunaria* (70/94), *Brachythecium sp.* (70/93), *Brachythecium velutinum* (77/93), *Bryum capillare* (77/93, 16/94), *Bryum imbricatum* (77/93, 18/91), *Bupleurum faictatum* (51/93:r, 86/93), *Calamagrostis arundinacea* (8/84:2, 47/93), *Campanula ciliata* (51/83, 54/83), *Campylotropus fragilis* (75/93), *Carex atrata* (70/93), *Carex caryophyllea* (70/94:1, 103/95), *Cladonia coniocraea* (70/94), *Cladonia crispata* (70/93), *Cladonia ecmocyna* (103/95), *Cladonia foliacea* (51/93:1), *Ceratium purpurascens* (19/81, 70/94), *Cladonia coniocraea* (70/94), *Dactylina madereponiformis* (75/93), *Daphne glomerata* (8/84), *Deschampsia flexuosa* (11/81, 87/95), *Desmodium latifolium* (77/93, 70/94:1), *Dicranum congestum* (54/83:1), *Distichlis capillacea* (103/95), *Draba hispida* (51/93, 86/93), *Draba scabra* (54/83), *Encalypta ciliata* (87/95, 103/95), *Gaulium verum* (103/95), *Gnaphalium supinum* (159/94, 45/91:r), *Heterocladium dimorphum* (86/93), *Hieracium macrolepis* (70/93, 70/94), *Hieracium prenanthoides* (103/95), *Hypnum cupressiforme* (103/95), *Isopterygiopsis pulchella* (77/93), *Jurinea moschata* (47/93, 86/93:r), *Kobresia schoenoides* (54/83:1), *Leskeella nervosa* (159/94), *Lloydia serotina* (54/83, 18/91:r), *Minuartia imbricata* (18/91), *Parmelia vagans* (51/93), *Pedicularis nordmanniana* (16/94), *Peltigera canina* (70/93, 70/94), *Physconia muscigena* (86/93), *Plagiochila porelloides* (103/95), *Polygonum viviparum* (12/84), *Potentilla nivea* (16/81, 18/81), *Pterigynandrum filiforme* (103/95), *Ranunculus caucasicus* (51/93), *Rhinanthus minor* (70/93, 86/93), *Saxifraga flagellaris* (18/91), *Sedum tenellum* (18/91, 8/84), *Sempervivum caucasicum* (51/93), *Senecio aurantiacus* (70/94), *Senecio karelinii* (16/81), *Stereocaulon apinum* (159/94), *Tortula ruralis* (70/93, 86/93), *Tritolum ambiguum* (70/93:1, 51/93:2), *Vicia cracca* (8/84, 86/93), *Weissia wimmeriana* (87/95)

Date (day month) size (sq m) and location of the relevées

Date (y.e./mon.), size (sq.m.) and location of the trees.

8/81 - 23.07, 100, M.Khatipara;	11/81 - 30.07, 25, M.Khatipara;	17/81 - 01.08, 16, M.Khatipara;	17/81 - 12.08, 16, M.Khatipara;	18/81 12.08, 25, M.Khatipara; 19/81 - 12.08,
25, M.Khatipara; 39/81 - 08.09, 100, M.Khatipara;	34/81 - 06.08, 25, M.Khatipara;	51/83 - 10.09, 25, Mukhu;	36/83 - 31.08, 16, M.Khatipara;	77/93 - 22.08, 9, M.Khatipara;
34/83 - 31.08, 25, M.Khatipara;	54/83 - 10.09, 25, Mukhu;	12/84 - 01.09, 16, M.Khatipara;	75/93 - 22.08, 9, M.Khatipara;	18/91 - 04.08, 6,
16/94 - 08.07, 16, Azgek;	15/94 - 05.09, 16, Oriuchat;	45/91 - 19.08, 16, Baduk;	70/93 - 17.08, 8, Baduk;	70/94 - 16.07, 15, Bol.Khatipara; 87/95 -
24.07, 16, Bol.Khatipara;	81/84 - 26.08, 25, M.Khatipara;	51/93 - 14.08, 16, M.Khatipara;	47/93 - 10.08, 16, M.Khatipara;	86/93 - 26.08, 25, M.Khatipara; 103/95 - 05.08, 25,
M.Khatipara.				