

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübél, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübél (Zürich)

Band: 130 (2002)

Artikel: Alpine vegetation of the Teberda Reserve, the northwestern Caucasus = Die Alpine Vegetation des Teberda Reservates, Nordwest-Kaukasus

Autor: Onipchenko, V.G.

Kapitel: 15: Multivariate analysis of the syntaxa

DOI: <https://doi.org/10.5169/seals-308994>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

15. Multivariate analysis of the syntaxa

We used detrended correspondence analysis (DCA) and weighted pair group cluster analysis (MVSP Software, KOVACH 1995) to estimate the relationships between the syntaxa of the vegetation. All (47) subassociations or associations (except *Ranunculo oreophili-Polygonetum aviculais*) were included in the analysis. Each syntaxon was represented by a complete list of species with frequency classes (1-5) as the abundance value of each species. Rare species found infrequently (class 1) in one syntaxon only were excluded from the general matrix before the processing. The matrix analyzed included 651 species.

The DCA -ordination patterns are represented in Fig. 15.1. on three main axes. The results of the cluster analysis are represented in the dendrogramm (Fig. 15.2.). Overall, the results showed a considerable correspondence with the position of the syntaxa in the floristic system, but some discrepancies have also been noted.

Communities of *Juncetea trifidi*, *Carici rupestris-Kobresietea bellardii* and *Polygono viviparum-Salicetum kazbekensis* form a separate group in all projections. They build a distinct cluster as well. All of these low productive alpine communities are typical of windward slopes with thin or absent snow cover in winter, so the extreme environment is responsible for their distinct floristic composition. On the other hand, as we have noted above, the floristic differences between *Juncetea trifidi* and *Carici rupestris-Kobresietea bellardii* are very subtle in our region. Perhaps, it would be better to consider them within the same class. The position of *Polygono viviparum-Salicetum kazbekensis* in *Loiseleurio-Vaccinietea* is more or less formal and might be revised in the future.

It is necessary to point out that the position of *Violo altaicae-Festucetum variae* is quite different from *Juncetea trifidi* (*Anemonion speciosae*) syntaxa. It confirms the position of the Caucasian *Festuca varia* dominated grasslands within *Nardetalia*.

All syntaxa of *Mulgedio-Aconitetea*, except one, form a separate cluster (Fig.15.2.). Their position on the DCA -axis surface is rather distinct as well. *Senecioni propinqui-Betuletum litwinowii* is positioned closely to the subalpine tall herbaceous communities, as we have discussed above (ch.13).





Fig. 15.1. (see next page)




Fig. 15.1.

Results of DCA-analysis (projection of syntaxa on three main ordination axes) Syntaxa:

Thlaspietea rotundifolii:

A - *Veronico minutae-Chaerophylletum humilis typicum*, **B** - *V.t.-Ch.h. lamietosum tomentosum*, **C** - *V.t.-Ch.h. saxifragetosum flagellaris*, **D** - *Hyalopoo ponticae-Oxyrietum digynae ranunculetosum oreophili*, **E** - *H.p.-O.d. typicum*, **F** - *Scrophulario olympicae-Epilobietum dodonaei*, **G** - *Dicranoweisio crispulae-Rubetum idaei*, **H** - *Silene compactae-Salicetum elbursensis*.

Asplenietea trichomanis:

I - *Potentilletum divinae*, **J** - *Astragaletum levieri*, **K** - *Galio valantioides-Polypodietum vulgare*, **L** - *Thymo-Seseliolum petraei*.

Scheuchzerio-Caricetea fuscae:

M - *Caro caucasici-Caricetum nigrae salicetosum kazbekensis*, **N** - *C.c.-C.n. typicum*, **O** - *Swertio ibericae-Caricetum nigrae*, **P** - *Caricetum rostratae*.

Montio-Cardaminetea: **Q** - *Cerastio cerastioidis-Cardaminetum uliginosi*.

Carici rupestris-Kobresietea bellardii:

R - *Drabo scabri-Kobresietum schoenoidis*, **S** - *Alchemillo-Kobresietum capilliformis*.

Juncetea trifidi:

T - *Campanulo ciliatae-Chamaesciadietum acaulis*, **U** - *Pediculari comosae-Eritrichietum caucasici oxytropidetosum kubanensis*, **V** - *P.c.-E.c. typicum*, **W** - *P.c.-E.c. bromopsietosum variegatae*.

Calluno-Ulicetea:

X - *Violo altaicae-Festucetum varia geranietosum renardii*, **Y** - *V.a.-F.v. typicum*, **Z** - *V.a.-F.v. nardetosum*, **a** - *Hedysaro caucasicae-Geranietum gymnocauli senecionetosum kolenatiani*, **b** - *H.c.-G.g. typicum*.

Salicetea herbaceae:

c - *Ranunculetum brachylobi*, **d** - *Hyalopoo ponticae-Pedicularietum nordmanniana*, **e** - *Saxifragetum sibiricae typicum*, **f** - *S.s. primuletosum amoena*, **g** - *S.s. saxifragetosum moschatae*.

Mulgedio-Aconitetea:

h - *Betonici macranthae-Calamagrostietum arundinaceae typicum*, **i** - *B.m.-C.a. veronicetosum peduncularis*, **j** - *Anthriscio sylvestris-Rumicetum alpini typicum*, **k** - *A.s.-R.a. senecionetosum platyphyllidis*, **l** - *Cephalario giganteae-Ligusticetum alani*, **m** - *Poetum longifoliae*.

Loiseleurio-Vaccinietea:

n - *Lerchenfeldio-Rhododendretum caucasici L.-R.c. oxalidetosum*, **o** - *L.-R.c. typicum*, **p** - *L.-R.c. pleurozietosum*, **q** - *Aconito nasuti-Juniperetum typicum*, **r** - *A.n.-J. chaerophylletosum rosei*, **s** - *Polygono viviparum-Salicetum kazbekensis*.

t - *D.c. Senecioni nemorensis-Betuletum litwinowii*.

Vaccinio-Piceetea: **u** - *Rhododendro caucasici-Betuletum litwinowii*.




Fig.15.2. Dendrogram of cluster analysis results. For the list of syntaxa see fig.15.1.

The alpine grassland (*Calluno-Ulicetea*) forms another compact group in the dendrogram. It occupies the central part of the ordination surfaces. The position can be attributed to rather favourable ecological conditions for this community where many alpine, as well as some subalpine species can grow. We consider *Hedysaro caucasicae-Geranium gymnocauli* as the "core" (according to KEDDY 1989, 1990) alpine community.

Rhododendron- and *Juniperus-*shrubs (*Loiseleurio-Vaccinietae*) are positioned close to the *Nardetalia* (Fig. 15.1.). One of the birch woodlands (*Rhododendro caucasici-Betuletum litwinowii*) also fits into this group.

Despite their significant floristic variability, rock outcrop communities (*Asplenieta trichomanis*) form a separate cluster in the dendrogram (Fig. 15.2.). They assume a relatively isolated position in the ordination surfaces as well. Syntaxa of *Thlaspietea rotundifolii* can be divided into 3 groups. *Silene compactae-Salicetum elbursensis* has the most distinct position from all other communities due to its great floristic richness and varied composition. Subalpine stable screes (*Dicranoweisio crispulae-Rubetum idaei*) are very closely related to *Rhododendron* - scrub (*Lerchenfeldio-Rhododendretum caucasici*). On the other hand alpine and subalpine unstable screes are the closest to the snowbeds and fens.

Alpine fens (*Scheuchzerio-Caricetea fuscae*) demonstrated a very distinct position on the first ordination axis (Fig. 15.1.a). The dry rock communities of the lower altitudes (*Thymo-Seselietum petraei*) occupy the opposite position. Therefore, we can consider the first axis as a complex moisture-altitude gradient.

The interpretation of the ecological sense of the second axis is more difficult. Species-poor high productive *Anthriscu sylvestris-Rumicetum alpini* communities occupied the extreme positions on the one side, on the other side there were species of rich unproductive dry rock communities (*Thymo-Seselietum petraei*).

Acknowledgements

I would like to thank all members of Moscow University expeditions, especially A. Egorov, O. Gorbachevskaya, S. Zenyakin, O. Cherednichenko, A. Aksenova, A. Sennov, G. Semenova, N. Lubeznova, for their indispensable help in the fieldwork. Professor B.M. Mirkin was the first who actively stimulate our syntaxonomic studies in Teberda. A.R. Ishbirdin, A.I. Solomeshch and V.V. Akatov made many helpful comments on the tables.

I am particularly thankful to K. Thompson for the English text correction and M.S. Blinnikov for helpful comments. D.E. Aksenov made the computer version of the map of Teberda Reserve. E.A. Ignatova determined all bryophyte samples. I would like to thank R. Stupf and T. Wilhalm for their help with the manuscript. Thanks are also due to the administration and staff of Teberda State Reserve for their continual assistance in the organization of field research.

John D. and Catherine T. MacArthur Foundation supported our investigations, as well as Russian Foundation for Fundamental Research (N 93-04-6498, 99-04-48012). Prof. Peter Edwards and Rübél Foundation made possible this publication in "Veröffentlichungen des Geobotanischen Institutes der ETH" series.