

Zeitschrift: Helvetia : magazine of the Swiss Society of New Zealand
Herausgeber: Swiss Society of New Zealand
Band: 75 (2009)
Heft: [8]

Artikel: The European mole
Autor: [s.n.]
DOI: <https://doi.org/10.5169/seals-944524>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Sender:

The Secretary of the Swiss Society of New Zealand Inc
Trudi Brühlmann
401 Marine Drive
Mahina Bay
Eastbourne
Lower Hutt 5013

New Zealand
Permit No. 93966



HELVE⁺IA

Magazine for Members of the Swiss Society of New Zealand Inc

FAUNA IN SWITZERLAND

The European Mole

The European mole has a cylindrical body and is around 12 cm long. Females are typically smaller than males. The eyes are small and hidden behind fur. Its ears are just a small ridge in the skin. The fur is usually dark grey, but the actual range of colors is larger, as due to the subterranean habits there is no disadvantage in off-colored fur.

Moles are found throughout Middle Asia and Europe, but not in Ireland. They are present in most habitats where the soil is deep enough to allow tunnelling but are uncommon in coniferous forests, on moorlands and in sand dunes, probably because their prey is scarce.

Moles spend almost all their lives underground in a system of permanent and semi-permanent tunnels which form a complex network hundreds of metres long at varying depths in the soil. The deepest tunnels are used most in times of drought and low temperatures. Permanent tunnels are used repeatedly for feeding over long periods of time, sometimes by several generations of moles. The fore limbs are used to dig, shearing soil from the sides of the tunnel with alternate strokes. Hind limbs are used to brace the mole's body against the tunnel walls. The mole turns round, scoops up accumulated soil with its fore limbs and pushes it along a previously dug side tunnel leading to the surface. The soil is pushed out above ground to form a molehill.

Within the tunnel system moles construct one or more spherical nest chambers, each lined with a ball of dry plant material. Nests are used for sleeping and for raising young.

Moles sometimes construct very large mounds containing more than 750kg of soil. The mounds have an internal structure with one or more nests

and a network of tunnels and can have food stores. They are built by males and females, and they occur most often in areas with shallow soil on hard substrate and in areas prone to flooding. The fortress acts as a refuge in times of flooding and also helps insulate the nest against low temperatures.



European mole, *Talpa europaea*

Earthworms are the most important component of the mole's diet; an 80g mole needs 50g of earthworms per day. Moles also eat many insect larvae particularly in the summer. Earthworms dominate the winter diet. Moles sometimes collect and store them alive in special chambers. The stored worms are immobilised by a bite to the head segment; 470 worms have been recorded in one chamber. Food is either actively dug out of the soil by the mole or more often collected from the floor of the tunnel. Many soil animals fall through into the tunnels. Moles rarely forage on the surface. *internet*