

Zeitschrift: Helvetia : magazine of the Swiss Society of New Zealand
Herausgeber: Swiss Society of New Zealand
Band: 76 (2010)
Heft: [8]

Artikel: Kreuzotter = Vipera berus
Autor: [s.n.]
DOI: <https://doi.org/10.5169/seals-944230>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Sender:

The Secretary of the Swiss Society of New Zealand Inc
Trudi Brühlmann
401 Marine Drive
Mahina Bay
Eastbourne
Lower Hutt 5013

New Zealand
Permit No. 93966



HELVE⁺IA

Magazine for Members of the Swiss Society of New Zealand Inc

FAUNA IN SWITZERLAND

Kreuzotter - *Vipera berus*



Vipera berus, the common European viper, is a venomous viper species that is extremely widespread and can be found throughout most of Western Europe and all the way to Far East Asia. It is not regarded as highly dangerous. The snake is not aggressive and usually only bites when alarmed or disturbed; people are generally only bitten after stepping on it or attempting to pick it up. It will usually disappear into the undergrowth at a hint of any danger, but will return once all is quiet, often to the same spot. Bites can be very painful, but are seldom fatal. The last snake bite fatality in Switzerland was recorded in 1961; in the 50 years since, over 20,000 people would have died in car accidents in Switzerland. Even so, professional medical help should always be sought as soon as possible after a bite.

The Kreuzotter feeds on small mammals, birds, lizards, amphibians and in some cases on spiders, worms and insects. Females breed once every two or three years with litters usually born in late summer to early autumn in the Northern hemisphere. The Kreuzotter, like most other vipers, is ovoviviparous, which means that the young develop inside eggs that are retained within the mother's body until they are ready to hatch. The young are usually born encased in a transparent sac from which they must free themselves. The neonates, measuring 14 to 23 cm, are born with a fully functional venom apparatus and a reserve supply of yolk within their bodies. They shed their skins for the first time within a day or two. Females do not appear to take much interest in their offspring, but the young have been observed to remain near their mothers for several days after birth. Litters range from 3 to 20.

Adults grow to a length of 60 to 90 centimetres. Most have some kind of zigzag dorsal pattern down

the entire length of the body and tail. Females are usually brownish in hue with dark-brown markings, while the males are pure grey with black markings.



Kreuzotter – *Vipera berus*

Males find females by following their scent trails, sometimes tracking them for hundreds of meters a day. If a female is found and flees, the male follows. Courtship involves side-by-side parallel "flowing" behavior, tongue flicking along the back and excited lashing of the tail. Pairs stay together for one or two days after mating. Males chase away their rivals and engage in combat, culminating in the dramatic "adder dance": In this act, the males confront each other, raise up the front part of the body vertically, make swaying movements and attempt to push each other to the ground. This is repeated until one of the two becomes exhausted and crawls off to find another mate.

from Wikipedia