

Zeitschrift: Helvetia : magazine of the Swiss Society of New Zealand
Herausgeber: Swiss Society of New Zealand
Band: 77 (2011)
Heft: [6]

Artikel: Fragaria vesca - wild or alpine strawberry
Autor: Brühlmann, Trudi
DOI: <https://doi.org/10.5169/seals-943396>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use


The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 24.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Sender:
The Secretary of the Swiss Society of New Zealand Inc
Trudi Brühlmann
401 Marine Drive
Mahina Bay
Eastbourne
Lower Hutt 5013

New Zealand
Permit No. 93966



HELVE^{CH}IA

Magazine for Members of the Swiss Society of New Zealand Inc

FLORA IN SWITZERLAND

Fragaria vesca – Wild or Alpine Strawberry

Do you remember all the secret places where you used to find Walderdbeeren in Switzerland? I still remember some – but I'm not going to tell you where they are.

Fragaria vesca, commonly known as the Woodland Strawberry, occurs naturally throughout the Northern Hemisphere. Other names for this species include Wild Strawberry, European Strawberry and Alpine Strawberry. Wild strawberries are usually everbearing, and produce few runners.

Typical habitat is along trails and roadsides, embankments, hillsides, stone- and gravel-laid paths and roads, meadows, young woodlands, sparse forest, woodland edges, and clearings. Often plants can be found where they do not get sufficient light to form fruit. In the southern part of its range, it can only grow in shady areas; further north it tolerates more sun. It is tolerant of a variety of moisture levels. It can survive mild fires and/or establish itself after fires.

Although *Fragaria vesca* primarily propagates via runners, viable seeds are also found in soil seed banks and seem to germinate when the soil is disturbed.

Walderdbeeren leaves serve as significant food source for a variety of ungulates, such as mule deer and elk, and the fruit are eaten by a variety of mammals and birds that also help to distribute the seeds in their droppings.

Evidence from archaeological excavations suggests that *Fragaria vesca* has been consumed by humans since the

Stone Age. The woodland strawberry was first cultivated in ancient Persia. Its seeds were later taken along the Silk Road towards the far East and to Europe where it was widely cultivated until the 18th century, when it began to be replaced by the garden strawberry, which has much larger fruit and showed



Fragaria vesca –Walderdbeeren

greater variation, making them better suited for further breeding.

Woodland strawberry fruit is strongly flavored, and is still collected and grown for domestic use and on a small scale commercially for the use of gourmets and as an ingredient for commercial jam, sauces, liqueurs, cosmetics and alternative medicine. In Turkey hundreds of tons of wild fruit are harvested annually, mainly for export. *tb/internet*