

**Zeitschrift:** Helvetica Physica Acta

**Band:** 21 (1948)

**Heft:** VI

**Artikel:** On the Magnetic Moments of the Neutron and Proton

**Autor:** Luttinger, J.M.

**DOI:** <https://doi.org/10.5169/seals-111924>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 23.02.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# On the Magnetic Moments of the Neutron and Proton

by J. M. Luttinger<sup>1)</sup>, ETH., Zürich.

(14. X. 1948.)

*Zusammenfassung:* In dieser Arbeit wird das magnetische Moment eines Nukleons berechnet für die neutrale, geladene und symmetrische pseudoskalare Mesonentheorie. Die Nukleonen werden durch die Löchertheorie beschrieben, so dass die Rechnung vollständig relativistisch ist. Die üblichen Divergenzen werden durch eine vom Verfasser angegebene Methode<sup>2)</sup> vermieden und — wie beim Electron — werden alle Resultate endlich. Die magnetischen Momente der Nukleonen lassen sich, in der  $g^2$ -Näherung, durch geschlossene Formeln darstellen.

## I. Introduction.

It is the purpose of this paper to show that a method previously developed<sup>2)</sup> for calculating the radiative correction to the  $g$ -factor of the electron may be extended to calculation of the  $g$ -factors of the neutron and proton in meson theory. For concreteness we have chosen the case of pseudoscalar mesons. Within the framework of this theory there are still two types of coupling possible<sup>3)</sup>: that in which a pseudovector is constructed out of the meson field, and that in which a pseudoscalar is constructed. It may be shown, however, by simple generalization and correction of a proof due to Dyson<sup>4)</sup> that the two theories will yield identical results for all the magnetic moments calculated in this paper. We shall therefore only give the calculation for the pseudoscalar type of

<sup>1)</sup> National Research Fellow.

<sup>2)</sup> J. M. LUTTINGER, Phys. Rev., **74**, 893 (1948). Henceforth cited as I.

<sup>3)</sup> N. KEMMER, Proc. Roy. Soc., A **166**, 127 (1938).

<sup>4)</sup> F. J. DYSON, Phys. Rev., **73**, 929 (1948). Dyson drops the terms quadratic in the coupling constant on the grounds that they are without physical significance. Actually these terms play an essential role in making the two types of coupling equivalent. The same proof shows that the nuclear forces are identical in the  $g^2$  approximation. Direct calculation verifies this and shows that the interaction terms quadratic in the coupling constant remove the  $1/r^3$  singularity in the forces, and thus possibly allow a ground state for the deuteron. These forces, however, are not at all static in nature and cannot be applied to calculate the ground state of the deuteron. Similar conclusions have been reached independently by Dr. L. VAN HOVE (unpublished). I am grateful to Dr. VAN HOVE for a valuable correspondence on this point.

coupling, as it is considerably simpler in structure. As a check on the algebraic work we have also carried out the calculations for the pseudovector coupling, and of course obtained the same results. It may be mentioned that it is the pseudovector type of coupling which has usually been discussed in connection with the meson theory of nuclear forces.

In handling the heavy particles we have used hole theory throughout, as it represents the only consistent relativistic theory for particles of spin one half. That is, we have taken the neutron and the proton to be quantized fields which obey Fermi-Dirac commutation relations and the Dirac equation without the presence of "Pauli terms"—it is just the hope of meson theory that it will account for the existence of such terms.

Three types of pseudoscalar meson field have been considered: neutral, charged and symmetric. From the point of view of comparison with the experimental facts perhaps only the latter is of interest, the nuclear forces generally being assumed to be charge independent<sup>1)</sup>. However, for completeness the other two cases are included.

## II. Neutral Theory.

For this case the meson field will only yield an additional magnetic moment for the proton—the magnetic moment of the neutron remaining zero. The Lagrangian density is given by<sup>2)</sup>.

$$\begin{aligned}
 L &= L_p + L_M + L' + L'' & (1) \\
 L_p &= -\psi_p^* \left( \frac{1}{i} \frac{\partial}{\partial t} + \frac{1}{i} \alpha \cdot (\nabla - ie A_0) + M\beta \right) \psi_p \\
 L_M &= \frac{1}{2} \left( \left( \frac{\partial \Phi}{\partial t} \right)^2 - (\nabla \Phi)^2 - \mu^2 \Phi^2 \right) \\
 L' &= -\sqrt{4\pi} ig (\psi_p^* \beta \gamma_5 \Phi \psi_p) \\
 L'' &= -2\pi g^2 \lambda [\psi_p^* \sigma \psi_p] \cdot [\psi_p^* \sigma \psi_p] - (\psi_p^* \gamma_5 \psi_p) (\psi_p^* \gamma_5 \psi_p)
 \end{aligned}$$

In these equations  $A_0$  is the vector potential of an external homogeneous magnetic field,  $\psi_p$  the (quantized) proton wave function,  $\alpha, \beta$  the well known Dirac matrices<sup>3)</sup>,  $\sigma$  the (four rowed) spin ma-

<sup>1)</sup> This charge independence has recently been questioned. See J. M. BLATT, Phys. Rev., **74**, 92 (1948). Blatt's results may, however, be dependent on his special assumptions concerning the potential.

<sup>2)</sup> We have used natural units,  $\hbar = c = 1$ .  $M$  is the nucleon mass,  $\mu$  the meson mass.

<sup>3)</sup> The specific representation which we use, however, has the usual representations of  $\alpha_x$  and  $\alpha_y$  interchanged. Cf. I, appendix.

trices,  $\gamma_5 = i\alpha_x\alpha_y\alpha_z$  and  $g$  the coupling constant. The number  $\lambda$  which appears in  $L''$  is an arbitrary constant. The presence of this term corresponds to the fact that one may get different versions of the pseudoscalar theory by adding an invariant term quadratic in the coupling constant (see Kemmer, op. cit.). The value of  $\lambda$  cannot be fixed a priori, but we shall see that  $L''$  gives no contribution to the magnetic moment and therefore this arbitrariness is in no way disturbing for our calculations.

Using this Lagrangian density it is an easy matter to find the Hamiltonian density.

$$H = H_p + H_M + H' + H'' \quad (2)$$

Where

$$H_p = \psi_p^* \left( \frac{1}{i} \alpha \cdot (\nabla - ieA_0) + M\beta \right) \psi_p$$

$$H_M = \frac{1}{2} (\pi^2 + (\nabla \Phi)^2 + \mu^2 \Phi^2)$$

$$H' = \sqrt{4\pi} ig (\psi_p \beta \gamma_5 \Phi \psi_p)$$

$$H'' = 2\pi g^2 \lambda [(\psi_p^* \sigma \psi_p) \cdot (\psi_p^* \sigma \psi_p) - (\psi_p^* \gamma_5 \psi_p) (\psi_p^* \gamma_5 \psi_p)].$$

To carry out the quantization<sup>1)</sup> we write<sup>2)</sup>:

$$\begin{aligned} \psi_p &= \sum_n a_n \psi_n \\ \Phi &= \sum_p \frac{\alpha_p + \alpha_{-p}^*}{\sqrt{2\omega_p}} e^{ip \cdot x} \\ \pi &= i \sum_p \sqrt{\frac{\omega_p}{2}} (\alpha_{-p}^* - \alpha_p) e^{ip \cdot x}. \end{aligned}$$

The  $\psi_n$  are the solutions corresponding to the energy  $E_n$  of the Dirac equation in the presence of an homogeneous external field (cf. I, appendix). The quantities  $a_n$  and  $\alpha_p$  satisfy the commutation rules of FERMI-DIRAC and EINSTEIN-BOSE particles respectively, i. e. all

$$\begin{aligned} a_n^* a_{n'} + a_{n'} a_n^* &= \delta_{nn'} \\ \alpha_p \alpha_{p'}^* - \alpha_{p'}^* \alpha_p &= \delta_{pp'} \end{aligned}$$

other commutators being zero.  $\omega_p$  represents the energy of a free meson,  $\omega_p = \sqrt{\mu^2 + p^2}$ .

<sup>1)</sup> As a general reference for the methods of quantization used here, see G. WENTZEL, Einf. in die Quant. Theorie der Wellenfelder, Franz Deuticke, Wien (1943).

<sup>2)</sup> Our solutions are normalized in a large box of volume  $V$ . However, since in all end results the quantity  $V$  drops out, we simply put it equal to 1.

Substituting in the Hamiltonian we obtain:

$$\begin{aligned}
 H_p &= \sum_n a_n^* a_n E_n \\
 H_M &= \sum_p \alpha_p^* \alpha_p \omega_p \\
 H' &= \sqrt{4\pi} ig \sum_{n, n'} \sum_p a_n^* a_{n'} \alpha_p^* \frac{Q_{nn'}^{(p)}}{\sqrt{2\omega_p}} + C. C. \\
 H'' &= 2\pi g^2 \lambda \sum_{\substack{n, n' \\ m, m'}} a_n^* a_{n'} a_m^* a_{m'} R_{nn' mm'}
 \end{aligned}$$

where

$$Q_{nn'}^{(p)} \equiv \int d^3 x (\psi_n^* \beta \gamma_5 \psi_{n'}) e^{-ip \cdot x}$$

$$R_{nn' mm'} \equiv \int d^3 x [(\psi_n \sigma \psi_{n'}) \cdot (\psi_m^* \sigma \psi_{m'}) - (\psi_n^* \gamma_5 \psi_{n'}) (\psi_m^* \gamma_5 \psi_{m'})].$$

Proceeding exactly as in the case of the electron, we calculate the energy of one proton in a state “ $m$ ” (and no mesons present), to terms proportional to the square of the coupling constant. This means calculating the mean value of  $H''$  for this state, and performing second order perturbation theory with respect to  $H'$ . The method is straightforward and proceeds exactly as in I. The results are:

$$E = E' + E'' \quad (3)$$

$$\begin{aligned}
 E' &= -2\pi g^2 \left[ \sum_p \frac{1}{\omega_p} \left\{ \sum_{E_n > 0} \frac{|Q_{nn}^{(p)}|^2}{\omega_p + E_n - E_m} \right. \right. \\
 &\quad \left. \left. - \sum_{E_n < 0} \frac{|Q_{nn}^{(p)}|}{\omega_p + |E_n| + E_m} \right\} + \sum_p \frac{Q_{nn}^{(p)*} Q_{mm}^{(p)} + Q_{mm}^{(p)*} Q_{nn}^{(p)}}{\omega_p^2} \right] \\
 E'' &= 2\pi g^2 \lambda \left( \sum_{E_n > 0} - \sum_{E_n < 0} \right) R_{mnmm} + 2\pi g^2 \lambda \sum_{E_n < 0} R_{mmnn}.
 \end{aligned}$$

The characteristic minus signs which arise in the summations over negative energies come from the use of hole theory, and the definition of the true energy as the energy of the particle plus vacuum, minus the energy of the vacuum.

To make use of the method of I, it is now necessary to specialize the state “ $m$ ”. As was there shown, if we take “ $m$ ” to be the solution of the DIRAC equation in which the energy is simply  $M$ , then any infinities in the mass or charge of the electron will not effect the magnetic moment. This means that simply by developing the quantity  $E$  in a power series in  $H_0$  (external field) the linear

term in  $H_0$  will converge and will give us directly the change in the magnetic moment of the nucleon. This expectation is borne out by the detailed calculation, from which one may conclude that in the pseudoscalar theory<sup>1)</sup> all the infinities are simply mass and charge infinities, just as in electrodynamics.

With this choice of state the evaluation of the matrix elements is simple, and we shall only quote the results<sup>2)</sup>:

$$E' = -\pi g^2 \sum_p \frac{1}{\omega_p} \left\{ \sum_{n=0}^{\infty} \left[ \frac{\left( \frac{2eH_0}{E_n(E_n+M)} \right)}{E_n + \omega_p - M} + \frac{\left( \frac{p_3^2}{E_n(E_n+M)} \right)}{E_n + \omega_p - M} \right] \frac{\xi^n}{n!} e^{-\xi} \right\} - (M \rightarrow -M) \quad (4)$$

where

$$E_n \equiv \sqrt{M^2 + p_3^2 + 2eH_0n}$$

$$\xi = \frac{p_1^2 + p_2^2}{2eH_0}$$

and  $(M \rightarrow -M)$  means that the same function of  $-M$  is to be constructed.

$$E'' = \pi g^2 \lambda \sum_{p_2, p_3} \sum_{n=0}^{\infty} \left( \frac{2(E_{n+1}+M)}{E_{n+1}} + \frac{E_n+M}{E_n} - \frac{E_n-M}{E_n} \right) - M \rightarrow -M \quad (5)$$

Simplification gives

$$E' = -\pi g^2 \sum_p \frac{1}{\omega_p} \left\{ \sum_{n=0}^{\infty} \frac{E_n-M}{\omega_p+E_n-M} \frac{1}{E_n} \frac{\xi^n}{n!} e^{-\xi} \right\} - (M \rightarrow -M) \quad (6)$$

$$E'' = (2M)(eH_0) \lambda g^2 \sum_{p_3} \sum_{n=0}^{\infty} \left( \frac{1}{E_{n+1}} + \frac{1}{E_n} \right). \quad (7)$$

In (7) we have replaced the summation over  $p_2$  by  $\frac{eH_0}{2\pi}$ , which corresponds to keeping the particles present within the normali-

<sup>1)</sup> VILLARS (in unpublished calculations) has shown that even with this special state the magnetic moment of the nucleons diverges for the vector theory—with tensor coupling—which seems to indicate that in this version of the vector meson theory there are infinities in addition to the mass and charge terms. This result, incidentally, prevents the calculation of the magnetic moments on the basis of a MØLLER-ROSENFIELD mixture.

The other version of vector meson theory (vector coupling, similar to coupling of electron to the electromagnetic field) has been investigated and found to give convergent magnetic moments.

<sup>2)</sup> The calculation involves an integration over Hermite polynomials which has been carried through in I.

zation volume<sup>1)</sup>. Let us first calculate  $E''$ . To terms linear in  $H_0$  we have

$$E'' = 2 M e H_0 \lambda g^2 \sum_{p_3} \sum_{n=0}^{\infty} \left( \frac{2}{E_n} - \frac{e H_0}{E_n^3} \right).$$

The EULER sum formula

$$\sum_{n=0}^{\infty} f(n) = \int_0^{\infty} f(x) dx + \frac{f(0)}{2} + \dots$$

provides an expansion in powers of  $H_0$ . Keeping only the linear terms one obtains

$$E'' = 2 M (e H_0) \lambda g^2 \sum_{p_3} \left\{ \int_0^{\infty} d n \left( \frac{2}{E_n} - \frac{e H_0}{E_n^3} \right) + \left[ \frac{1}{E_n} \right]_{n=0} \right\}.$$

The first term is independent of  $H_0$ , and represents an infinite self energy. Dropping this, we get for the terms linear in  $H_0$ :

$$E'' = 2 M (e H_0) \lambda g^2 \sum_{p_3} \left( \left[ \frac{1}{E_n} \right]_{n=0} - e H_0 \int_0^{\infty} \frac{1}{E_n^3} d n \right).$$

However,

$$\int_0^{\infty} d n \frac{e H_0}{E_n^3} = \left[ \frac{1}{E_n} \right]_{n=0}$$

and therefore

$$E'' = 0. \quad (8)$$

Returning now to the discussion of  $E'$ , we make use of the identity

$$\frac{1}{\omega_p} \frac{E_n - M}{\omega_p + E_n - M} = \left( \frac{1}{\omega_p} - \frac{1}{\omega_p + E_n - M} \right)$$

which transforms (6) into

$$E' = \pi g^2 \sum_p \sum_{n=0}^{\infty} \frac{1}{E_n} \frac{1}{(\omega_p + E_n - M)} \frac{\xi^n}{n!} e^{-\xi} - (M \rightarrow -M). \quad (9)$$

Defining a quantity (Cf. I)

$$\begin{aligned} f(n) &= \frac{1}{E_n} \frac{1}{\omega_p + E_n - M} \\ &= f(\xi) + (n - \xi) f'(\xi) + \frac{(n - \xi)^2}{2} f''(\xi) + \dots \end{aligned}$$

<sup>1)</sup> BETHE-SOMMERFELD, Hand. der Phys., XXIV, 2, page 478, Julius Springer, Berlin (1933).

we get

$$\begin{aligned} E' &= \pi g^2 \sum_p \sum_{n=0}^{\infty} \left( f(\xi) + (n-\xi) f'(\xi) + \frac{(n-\xi)^2}{2} f''(\xi_1) + \dots \right) \frac{\xi^n}{n!} e^{-\xi} \\ &\quad - (M \rightarrow -M) \\ &= \pi g^2 \sum_p \left( f(\xi) + \frac{\xi}{2} f''(\xi) + \dots \right) - (M \rightarrow M). \end{aligned}$$

The first term of this expression is independent of  $H_0$  and represents a self energy of the proton. We shall therefore drop it along with the remainder terms in the expansion, the latter representing higher powers of  $H_0$  than the first. Therefore, for the purpose of calculating the magnetic moment we have

$$E' = \pi g^2 \sum_p \frac{\xi}{2} f''(\xi) - (M \rightarrow -M). \quad (10)$$

Carrying through the indicated differentiations one obtains

$$E' = \frac{\pi g^2 (e H_0)}{4} \sum_p (p_1^2 + p_2^2) \left[ \frac{3}{\Omega^2 (\Omega + \omega - M)} + \frac{3}{\Omega (\Omega + \omega - M)^2} + \frac{2}{(\Omega + \omega - M^3)} \right] \frac{1}{\Omega^3} - (M \rightarrow -M). \quad (11)$$

Here

$$\Omega \equiv \sqrt{M^2 + p^2}, \quad \omega \equiv \sqrt{\mu^2 + p^2}.$$

Noting the fact that the factor multiplying  $p_1^2 + p_2^2$  in (11) is a function of  $p^2$  only, so that we may replace  $p_1^2 + p_2^2$  by  $\left(\frac{2}{3}\right) p^2$ , we get

$$\begin{aligned} E' &= \frac{\pi g^2 (e H_0)}{3} (M) \sum_p p^2 \frac{1}{\Omega^3} \\ &\quad \left[ \frac{3}{\Omega^2 ((\omega + \Omega)^2 - M^2)} + \frac{6 (\omega + \Omega)}{\Omega ((\omega + \Omega)^2 - M^2)^2} + \frac{2 (3 (\omega + \Omega)^2 + M^2)}{((\omega + \Omega)^2 - M^2)^3} \right]. \end{aligned}$$

Finally, we replace the sum by an integral

$$\left( \sum_p \rightarrow \left( \frac{1}{2\pi} \right)^3 \int d^3 p \right),$$

and carry out the angle integration:

$$\begin{aligned} E' &= \frac{g^2 (e H_0) M}{6\pi} \int_0^\infty dp \frac{p^2}{\Omega^3} \\ &\quad \left[ \frac{3}{\Omega^2 ((\omega + \Omega)^2 - M^2)} + \frac{6 (\omega + \Omega)}{\Omega ((\omega + \Omega)^2 - M^2)^2} + \frac{2 (3 (\omega + \Omega)^3 + M^2)}{((\omega + \Omega)^2 - M^2)^3} \right]. \quad (12) \end{aligned}$$

The integral in (12) is elementary, as may be seen by introducing the new variable  $z$ , defined by  $\Omega + \omega = \sqrt{z}$ . The result is

$$E'_p = \frac{g^2 (e H_0)}{8 \pi M} \left( 1 + 2 \delta^2 - \frac{2 \delta^3 (3 - \delta^2)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} - 2 (1 - \delta^2) \delta^2 \log \frac{1}{\delta} \right)$$

where  $\delta$  is the ratio of meson to proton (or neutron) mass.

$$\mu_p = - \frac{\partial E'_p}{\partial H_0} = - \frac{g^2}{\pi} \left( \frac{1}{4} + \frac{\delta^2}{2} - \frac{\delta^3 (3 - \delta^2) \cos^{-1} \delta/2}{2 \sqrt{4 - \delta^2}} - \frac{(1 - \delta^2) \delta^2}{2} \log \frac{1}{\delta} \right) \quad (13)$$

in units of the nuclear magneton.

### III. Charged Theory.

The Lagrangian density is given by

$$L_c = L_p + L_N + L_{MC} + L'_c + L''_c \quad (14)$$

Here  $L_p$  is defined as above, while

$$\begin{aligned} L_N &= -\psi_N^* \left( -\frac{1}{i} \frac{\partial}{\partial t} + \frac{1}{i} \alpha \cdot \nabla + M \beta \right) \psi_N \\ L_{MC} &= -\left( \frac{\partial \Phi^*}{\partial t} \right) \left( \frac{\partial \Phi}{\partial t} \right) - (\nabla \Phi^* + i e A_0 \Phi^*) \cdot (\nabla \Phi - i e A_0 \Phi) - \mu^2 \Phi^* \Phi \\ L'_c &= -\sqrt{8\pi} i g [(\psi_p^* \beta \gamma_5 \Phi \psi_N) + (\psi_N^* \beta \gamma_5 \Phi^* \psi_p)] \\ L''_c &= 4\pi g^2 \lambda [(\psi_p^* \sigma \psi_N) \cdot (\psi_N^* \sigma \psi_p) - (\psi_p^* \gamma_5 \psi_N) (\psi_N^* \gamma_5 \psi_p) \\ &\quad + (\psi_N^* \sigma \psi_p) \cdot (\psi_p^* \sigma \psi_N) - (\psi_N^* \gamma_5 \psi_p) (\psi_p^* \gamma_5 \psi_N)] \end{aligned}$$

$\psi_N$  is the (quantized) wave function of the neutron,  $\Phi$  the charged meson field. The term  $L''_c$  gives a contribution  $(-L'_p)$  in the Hamiltonian density. Just as in the case of neutral mesons this term will give no contribution to the magnetic moment, and in what follows we shall drop it entirely. We then obtain for the Hamiltonian density

$$\begin{aligned} H_c &= H_p + H_N + H_{MC} + H'_c \\ H_N &= \psi_N^* (1/i \alpha \cdot \nabla + M \beta) \psi_N \\ H_{MC} &= \pi^* \pi + (\nabla \Phi^* + i e A_0 \Phi^*) \cdot (\nabla \Phi - i e A_0 \Phi) + \mu^2 \Phi^* \Phi \\ H'_c &= -L'_p. \end{aligned} \quad (15)$$

We now carry through the quantization by means of the relationships

$$\begin{aligned}\psi_N &= \sum_n b_n \tilde{\psi}_n \\ \psi_p &= \sum_n a_n \psi_n \\ \Phi_l &= \sum_l \frac{\alpha_l + \beta_l^*}{\sqrt{2 \varepsilon_l}} \Phi_l \\ \pi^* &= -i \sum_l \sqrt{\frac{\varepsilon_l}{2}} (\alpha_l - \beta_l^*) \Phi_l.\end{aligned}\tag{16}$$

$\psi_n$  are solutions of the DIRAC equation in the presence of an homogeneous magnetic field,  $\tilde{\psi}_n$  are solutions of the field free DIRAC equation (corresponding to the energy  $\tilde{E}_n$ ), and the  $\Phi_l$  are solutions of the KLEIN-GORDON equation in the magnetic field. The quantities  $\varepsilon_l$  are the corresponding (positive) frequencies, *i.e.*

$$((\nabla - ie A_0)^2 - \mu^2) \Phi_l = \varepsilon_l^2 \Phi_l\tag{17}$$

The solutions of (17) are well known<sup>1)</sup>. One finds that  $l$  is given by an integer  $N \geq 0$  and two momenta  $k_2, k_3$ :

$$\begin{aligned}\Phi_l &= e^{i(k_2 y + k_3 z)} e^{-\eta^2/2} H_N(\eta) \frac{(e H_0)^{1/4}}{\pi^{1/4} 2^{N/2} \sqrt{N!}}, \quad \eta \equiv \sqrt{e H_0} \left( x - \frac{k_2}{e H_0} \right) \\ \varepsilon_l &= \sqrt{\mu^2 + k_3^2 + 2e H_0 (N + \frac{1}{2})}.\end{aligned}$$

The quantities  $H_N$  are the ordinary Hermite polynomials.

The quantities  $a_n, b_n$  satisfy FERMI-DIRAC commutation rules, the quantities  $\alpha_l, \beta_l$  EINSTEIN-BOSE ones. Using (16) we obtain

$$\begin{aligned}H_p &= \sum_n a_n^* a_n E_n \\ H_N &= \sum_n b_n^* b_n \tilde{E}_n \\ H_{MC} &= \sum_l (\alpha_l^* \alpha_l + \beta_l^* \beta_l) \varepsilon_l \\ H_c' &= \sqrt{4\pi} ig \sum_{n,m} \frac{1}{\sqrt{\varepsilon_l}} a_n^* b_m T_{nm}^{(l)} (\alpha_l + \beta_l^*) + c.c. \\ T_{nm}^{(l)} &\equiv \int d^3 x (\psi_n^* \beta \gamma_5 \Phi_l \tilde{\psi}_m).\end{aligned}\tag{18}$$

It is now necessary to calculate the energy of a neutron or proton in the state "m", where "m" is again the special state which avoids the divergencies in the mass and charge. This choice is necessary

<sup>1)</sup> BETHE-SOMMERFELD, Hand. der Phys., XXIV, 2, page 478, Julius Springer, Berlin (1933).

here only in the case of the proton: for the neutron (since its energy is always independent of  $H_0$ ) any state would do. We calculate with the state “ $m$ ”, however, because it is mathematically the simplest state to handle. Straightforward second order perturbation theory then leads to the result ( $E_N^c, E_P^c$  being the corrections to the energy of the neutron and proton respectively):

$$E_N^c = -4\pi g^2 \sum_l \frac{1}{\varepsilon_l} \left\{ \sum_{e_m > 0} \frac{|T_{nm}^{(l)}|^2}{E_n + \varepsilon_l - M} - \sum_{\tilde{E}_n < 0} \frac{|T_{nm}^{(l)}|^2}{|\tilde{E}_n| + \varepsilon_l + M} \right\}. \quad (19)$$

$$E_P^c = -4\pi g^2 \sum_l \frac{1}{\varepsilon_l} \left\{ \sum_{\tilde{E}_n > 0} \frac{|T_{mn}^{(l)}|^2}{\tilde{E}_n + \varepsilon_l - M} - \sum_{E_n < 0} \frac{|T_{nm}^{(l)}|^2}{|E_n| + \varepsilon_l + M} \right\}. \quad (20)$$

Calculation of the matrix elements gives

$$E_N^c = -2\pi g^2 \sum_{p_2, p_3} \sum_{n=0}^{\infty} \left\{ \frac{2eH_0(n+1)}{\varepsilon_{n+3/2} E_{n+1}(E_{n+1}+M)(E_{n+1}+\varepsilon_{n+3/2}-M)} + \frac{p_3^2}{\varepsilon_{n+1/2} E_n(E_n+M)(E_n+\varepsilon_{n+1/2}-M)} \right\} - (M \rightarrow -M). \quad (21)$$

$$E_P^c = -2\pi g^2 \sum_p \sum_{N=0}^{\infty} \left\{ \frac{p_1^2 + p_2^2}{(\Omega + \varepsilon_{N+1/2} - M)} + \frac{p_3^2}{(\Omega + \varepsilon_{N+1/2} - M)} \right\} \frac{e^{-\xi}}{\Omega(\Omega+M)} \frac{\xi^N}{N!} - (M \rightarrow -M). \quad (22)$$

$$E_n \equiv \sqrt{M^2 + p_3^2 + 2eH_0 n}$$

$$\varepsilon_n \equiv \sqrt{\mu^2 + p_3^2 + 2eH_0 n}$$

$$\Omega \equiv \sqrt{M^2 + p^2}$$

$$\xi = \frac{p_1^2 + p_2^2}{2eH_0}.$$

Let us first calculate  $E_N^c$ .

$$E_N^c = -2\pi g^2 \frac{eH_0}{2\pi} \sum_{p_3} \sum_{n=0}^{\infty} \left\{ \frac{p_3^2 + 2eH_0 n}{\varepsilon_{n+1/2} E_n(E_n+M)(E_n+\varepsilon_{n+1/2}-M)} \right\} - (M \rightarrow -M).$$

$$= -g^2 (eH_0) \sum_{p_3} \sum_{n=0}^{\infty} \left\{ \frac{E_n - M}{\varepsilon_{n+1/2} E_n(E_n+\varepsilon_{n+1/2}-M)} \right\} - (M \rightarrow M).$$

$$= g^2 (eH_0) \sum_{p_3} \sum_{n=0}^{\infty} \frac{1}{E_n(E_n+\varepsilon_{n+1/2}-M)} - (M \rightarrow -M).$$

$$= g^2 (eH_0) \sum_{p_3} \sum_{n=0}^{\infty} \left\{ \frac{1}{E_n(E_n+\varepsilon_n-M)} - \frac{1}{E_n(E_n+\varepsilon_n-M)^2} \frac{eH_0}{2\varepsilon_n} \right\}$$

$$+ O(H_0^2) - (M \rightarrow -M_0^2).$$

Using once more the EULER sum formula used in the evaluation of (7), and dropping self energy terms and those which are proportional to the square of the field strength, we get

$$\begin{aligned}
 E_N^c &= g^2 (e H_0) \sum_{p_3} \left\{ \frac{1}{2} \left[ \frac{1}{E_n (E_n + \varepsilon_n - M)} \right]_{n=0} \right. \\
 &\quad \left. - \frac{e H_0}{2} \int_0^\infty d n \frac{1}{E_n \varepsilon_n (E_n + \varepsilon_n - M)^2} \right\} - (M \rightarrow -M). \\
 &= g^2 \frac{e H_0}{2 \pi} \int_{-\infty}^{+\infty} d p_3 \left\{ \frac{1}{2} \left[ \frac{1}{E_n (E_n + \varepsilon_n - M)} \right]_{n=0} \right. \\
 &\quad \left. - \frac{e H_0}{2} \int_0^\infty d n \frac{1}{E_n \varepsilon_n (E_n + \varepsilon_n - M)^2} \right\} - (M \rightarrow -M).
 \end{aligned}$$

The second integral may be considerably simplified. Let  $r^2 = 2 e H \cdot n$ ,  $\omega = \sqrt{\mu^2 + p^2}$ ,  $\Omega = \sqrt{\mu^2 + p^2}$ ,  $p^2 = p_3^2 + r^2$ , then

$$\begin{aligned}
 \int_{-\infty}^{\infty} d p_3 \int_0^{\infty} d n \frac{e H_0}{E_n \varepsilon_n (E_n + \varepsilon_n - M)^2} &= \int_{-\infty}^{\infty} d p_3 \int_0^{\infty} r dr \frac{1}{\Omega \omega (\omega + \Omega - M)^2} \\
 &= \frac{1}{2 \pi} \int d^3 p \frac{1}{\omega \Omega (\omega + \Omega - M)^2} = 2 \int_0^{\infty} d p \frac{p^2}{\omega \Omega (\omega + \Omega - M)^2} \\
 &= \int_{-\infty}^{\infty} d p \frac{p^2}{\omega \Omega (\omega + \Omega - M)^2}.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 E_N^c &= \frac{(e H_0) g^2}{4 \pi} \int_{-\infty}^{\infty} d p \left\{ \frac{1}{\Omega (\omega + \Omega - M)} - \frac{p^2}{\Omega \omega (\omega + \Omega - M)^2} \right\} - (M \rightarrow -M) \\
 &= \frac{(e H_0) g^2}{\pi} \int_0^{\infty} d p \left\{ \frac{1}{\Omega ((\omega + \Omega)^2 - M^2)} - \frac{p^2}{\omega \Omega} \frac{\omega + \Omega}{((\omega + \Omega)^2 - M^2)^2} \right\}. \quad (23)
 \end{aligned}$$

The integral in (23) is elementary, and may be reduced to known form by the same substitution that was used in (12). The result is

$$\begin{aligned}
 E_N^c &= \frac{(e H_0) g^2}{2 \pi} \frac{1}{M} \left( 1 - \frac{\delta (2 - \delta^2)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} + \delta^2 \log \frac{1}{\delta} \right) \\
 \mu_N^c &= - \frac{g^2}{\pi} \left( 1 - \frac{\delta (2 - \delta^2)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} + \delta^2 \log \frac{1}{\delta} \right) \quad (24)
 \end{aligned}$$

\*

Proceeding to  $E_P^c$ , we may write (22) as

$$\begin{aligned} E_P^c &= +2\pi g^2 \sum_p \sum_{N=0}^{\infty} \left\{ \frac{1}{\Omega(\Omega + \varepsilon_N + \frac{1}{2} - M)} \right\} \frac{\xi^N}{N!} e^{-\xi} - (M \rightarrow -M) \\ &= 2\pi g^2 \sum_p \sum_{N=0}^{\infty} \left\{ \frac{1}{\Omega(\Omega + \varepsilon_N - M)} - \frac{eH_0}{2\varepsilon_N} \frac{1}{(\Omega + \varepsilon_N - M)^2} \right\} \frac{\xi^N}{N!} e^{-\xi} \\ &\quad - (M \rightarrow -M) + O(H_0^2). \end{aligned}$$

Define a function

$$g(N) = \frac{1}{\varepsilon_N + \Omega - M} = g(\xi) + (N - \xi) g'(\xi) + \frac{(N - \xi)^2}{2} g''(\xi) + \dots$$

Then, after dropping the self energy terms and terms in  $H_0^2$ , we get

$$E_P^c = 2\pi g^2 \sum_p \frac{1}{\Omega} \left\{ \frac{\xi}{2} g''(\xi) - \frac{eH_0}{2\omega} \frac{1}{(\omega + \Omega - M)^2} \right\} - (M \rightarrow -M).$$

Carrying out the  $x$  differentiation yields:

$$\begin{aligned} E_P^c &= \pi g^2 (eH_0) \sum_p \frac{1}{\Omega} \left\{ \frac{p_1^2 + p_2^2}{2(\omega + \Omega - M)^2 \omega^2} \left( \frac{1}{\omega} + \frac{2}{\omega + \Omega - M} \right) - \frac{1}{\omega} \frac{1}{(\omega + \Omega - M)^2} \right\} - (M \rightarrow -M) \\ &= \pi g^2 (eH_0) \sum_p \frac{1}{\Omega} \left\{ \frac{p^2}{3(\omega + \Omega - M)^2 \omega^2} \left( \frac{1}{\omega} + \frac{2}{\omega + \Omega - M} \right) - \frac{1}{\omega} \frac{1}{(\omega + \Omega - M)^2} \right\} - (M \rightarrow -M) \\ &= \frac{g^2 (eH_0)}{2\pi} \int_0^\infty dp \frac{p^2}{\Omega \omega} \left[ \frac{2}{3} \frac{p^2}{\omega (\omega + \Omega - M)^3} + \frac{1}{(\omega + \Omega - M)^2} \left( \frac{1}{3} \frac{p^2}{\omega^2} - 1 \right) \right] - (M \rightarrow -M) \\ &= \frac{2g^2 (eH_0)}{\pi} \int_0^\infty dp \frac{p^2}{\omega \Omega} \left[ \frac{1}{3} \frac{p^2 [3(\omega + \Omega)^2 + M^2]}{\omega ((\omega + \Omega)^2 - M^2)^3} - \frac{(\Omega + \omega)}{((\omega + \Omega)^2 - M^2)^2} \left( 1 - \frac{1}{3} \frac{\omega^2}{p^2} \right) \right] \quad (25) \end{aligned}$$

Just as in (12) and (23) the integral of (25) is elementary. One obtains

$$\begin{aligned} E_P^c &= -\frac{g^2 (eH_0)}{2M\pi} \left( \frac{1}{2} - \delta^2 - \frac{\delta(2 - 4\delta^2 + \delta^4)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} + \delta^2 (2 - \delta^2) \log \frac{1}{\delta} \right) \\ \mu_P^c &= \frac{g^2}{\pi} \left( \frac{1}{2} - \delta^2 - \frac{\delta(2 - 4\delta^2 + \delta^4)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} + \delta^2 (2 - \delta^2) \log \frac{1}{\delta} \right). \quad (26) \end{aligned}$$

#### IV. The Pseudovector Coupling.

With this coupling the interactions  $L'$  and  $L'_c$  are replaced by

$$L' \text{ (pseudovector coupling)} = -\sqrt{4\pi} f \left( \psi_p^* \left( \sigma \cdot \nabla \Phi - \gamma_5 \frac{\partial \Phi}{\partial t} \right) \psi_p \right)$$

$$L'_c \text{ (pseudovector coupling)} = -\sqrt{8\pi} f \left[ \left( \psi_p^* \left[ \sigma \cdot (\nabla - ie A_0) \Phi - \gamma_5 \frac{\partial \Phi}{\partial t} \right] \psi_N \right) \right. \\ \left. + \left( \psi_N^* \left[ \sigma \cdot (\nabla + (ie A_0) \Phi^* - \gamma_5 \frac{\partial \Phi^*}{\partial t} \right] \psi_p \right) \right]$$

When the coupling constants are related by  $g^2 = (2Mf)^2$ , then the two types of coupling give identical results. The values usually given in the literature<sup>1)</sup> are  $(fM)^2 \sim 9$ , for the meson mass equal to either 200 or 325 electron masses. These values are extremely tentative, however, and possibly should be subject to suspicion even in so far as order of magnitude goes.

#### V. Summary of Results<sup>2)</sup>.

##### Neutral Mesons.

$$\mu_N = 0$$

$$\mu_p = -\frac{g^2}{\pi} \left( \frac{1}{4} + \frac{\delta^2}{2} - \frac{\delta^3 (3 - \delta^2)}{2\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} - \frac{\delta^2}{2} (1 - \delta^2) \log \frac{1}{\delta} \right).$$

##### Charged Mesons.

$$\mu_N^c = -\frac{g^2}{\pi} \left( 1 - \frac{\delta (2 - \delta^2)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} + \delta^2 \log \frac{1}{\delta} \right).$$

$$\mu_P^c = \frac{g^2}{\pi} \left( \frac{1}{2} - \delta^2 - \frac{\delta (2 - 4\delta^2 + \delta^4)}{\sqrt{4 - \delta^2}} \cos^{-1} \frac{\delta}{2} + \delta^2 (2 - \delta^2) \log \frac{1}{\delta} \right).$$

##### Symmetrical Theory.

$$\mu_N^s = \mu_N^c$$

$$\mu_P^s = \mu_P^c + \mu_P.$$

The units are those of the nuclear magneton, and  $\delta$  is the ratio of meson to proton mass. When the above numerical values for the coupling constant and meson mass are used, one obtains:

A)  $\mu \sim 325 m$  (electron).

$$\mu_N = 0 \quad \mu_N^c \sim -8 \quad \mu_N^s \sim -8$$

$$\mu_P \sim -2.8 \quad \mu_P^c \sim 2.5 \quad \mu_P^s \sim -.3.$$

<sup>1)</sup> F. VILLARS, Helv. Phys. Acta, XX, 476 (1947).

<sup>2)</sup> These results satisfy the relationship  $\mu_N^c + \mu_P^c = 2 \mu_N^N$ , which has a general validity in the  $g^2$ -approximation. (The factor 2 appears because we have chosen the coupling constant differently for charged and neutral theory.)

B)  $\mu \sim 200 m$  (electron).

$$\begin{array}{lll} \mu_N = 0 & \mu_N^c \sim -9.3 & \mu_N^s \sim -9.3 \\ \mu_P \sim -2.8 & \mu_P^c \sim 3.6 & \mu_P^s \sim .8 \end{array}$$

$$\begin{array}{ll} \mu_N \text{ (experimental)} = -1.9103 \\ \mu_P \text{ (experimental)} = 1.7896 \end{array}$$

It is clear that none of the theories give agreement with experiment. More disturbing than the incorrectness of the absolute values is that of the ratio of the two moments. This is independent of the value of the coupling constant, and depends only on the ratio  $\delta$ . It seems impossible to fit the data with any value of  $\delta$ . Whether the error lies in the model (pseudoscalar mesons) or in the use of perturbation theory remains an open question.

In conclusion I should like to thank Dr. F. VILLARS for suggesting this problem, and for much help in its solution. I am also indebted to Prof. W. PAULI and Dr. RES JOST for many stimulating and helpful discussions.