Zeitschrift:	Helvetica Physica Acta
Band:	22 (1949)
Heft:	IV
Artikel:	Remarks on the problem of the vacuum polarization and the photon- self-energy
Autor:	Jost, Res / Rayski, Jerzy
DOI:	https://doi.org/10.5169/seals-112011

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Remarks on the problem of the vacuum polarization and the photon-self-energy

by Res Jost*) and Jerzy Rayski**).

(17. V. 1949.)

As is well known the formulae for the polarization of the vacuum and the expression for the photon self energy are not well defined mathematical symbols and lead to conflict with the postulate of gauge invariance unless a certain regularizing procedure¹) is applied. This procedure of regularization means however a change in the fundamentals of field theory, so that the question as to other possibilities of obtaining the desired result (i.e. the gauge invariance of the vacuum polarization current and the vanishing of the ph. s. e.) seems still to be of some interest. Indeed, it has been shown by Umesawa²), and independently, by one of us³) that by coupling the electromagnetic field to several charged fields of spinor and scalar (or pseudoscalar) types the gauge invariance of the vacuum polarization and the vanishing of the ph. s. e. may be achieved (at least in the second order of approximation e^2) due to some compensations. It would be premature to decide whether such compensations posess any deeper meaning or whether they should be considered merely as accidental. In any case it should be noted that one of the rules for regularization (i.e. the regularization of the whole expressions without factorization) appears as if borrowed from the theory of several mutually compensating fields. On the other hand it should be stressed that there still remains another difficulty i. e. the infinite self charge produced by the external field. This is, of course, immaterial for the photon self energy but plays a role in the problem of the vacuum polarization. In order to obtain a reasonable value for the vacuum polarization effects, the charges of the spinor and scalar fields have to be renormalized.

An external electromagnetic field $A_{\mu}(x)$ coupled with *n* fields of a spinor type (with the mass constants m_i) and with N fields of

^{*)} Eidg. Techn. Hochschule, Zurich.

^{**)} N. COPERNICUS, Univ. Torun, Poland. At present: Eidg. Techn. Hochschule, Zurich.

a scalar (or pseudoscalar) type (with masses M_i) is given. The charge and current of the vacuum polarization then assumes the form:

$$< j_{\mu}(x) >_{\rm vac} = -e^2 \int d^4 x' K_{\mu\nu}(x-x') A_{\nu}(x')$$
 (1)

where $K_{\mu\nu}$ (in the e² approximation) consists additively of separate terms for different charged fields:

$$K_{\mu\nu} = \sum_{i=1}^{n} K_{\mu\nu}^{(1/2)}(m_i) + \sum_{t=1}^{N} K_{\mu\nu}^{(0)}(M_i)$$
(2)

with

$$K^{(1/2)}_{\mu\nu} = 4 \left\{ \frac{\partial \bar{\Delta}}{\partial x_{\mu}} \frac{\partial \Delta^{(1)}}{\partial x_{\nu}} + \frac{\partial \bar{\Delta}}{\partial x_{\nu}} \frac{\partial \Delta^{(1)}}{\partial x_{\mu}} - \delta_{\mu\nu} \left(\frac{\partial \bar{\Delta}}{\partial x_{\lambda}} \frac{\partial \Delta^{(1)}}{\partial x_{\lambda}} + m^2 \bar{\Delta} \Delta^{(1)} \right) \right\} \quad (2')$$

$$\begin{split} K^{(0)}_{\mu\nu} &= \left\{ -\frac{\partial\bar{\varDelta}}{\partial x_{\mu}} \frac{\partial\varDelta^{(1)}}{\partial x_{\nu}} - \frac{\partial\bar{\varDelta}}{\partial x_{\nu}} \frac{\partial\varDelta^{(1)}}{\partial x_{\mu}} + \frac{\partial^{2}\bar{\varDelta}}{\partial x_{\mu}\partial x_{\nu}} \varDelta^{(1)} + \bar{\varDelta} \frac{\partial^{2}\varDelta^{(1)}}{\partial x_{\mu}\partial x_{\nu}} \right. \\ &+ \left. \delta_{\mu\nu} \left(-\bar{\varDelta} \cdot \Box \varDelta^{(1)} - \varDelta^{(1)} \cdot \Box \bar{\varDelta} + 2 M^{2} \bar{\varDelta} \varDelta^{(1)} \right) \right\} \end{split}$$

where the $\overline{\Delta}$ and $\Delta^{(1)}$ functions are given by J. SCHWINGER⁴). As the expression for the vacuum polarization current of a spinor field has already been extensively discussed in the literature (compare e.g. footnote ¹), we limit ourselves here to giving only the derivation of $K_{\mu\nu}^{(0)}$:

The Hamiltonian of a scalar (or pseudoscalar) field coupled with an external electromagnetic field is⁵):

$$H_0 + H'$$

where
$$H_0 = \frac{1}{2} \int d^3 x \left(\pi^* \pi + \operatorname{grad} \varphi^* \operatorname{grad} \varphi + M^2 \varphi^* \varphi \right)$$
 (3')

and
$$H' = \int d^3 x \left(\varrho A_0 - \sum_{i=1}^3 s_i A_i + e^2 \varphi^* \varphi \sum_{i=1}^3 A_i A_i \right)$$
 (3")

with
$$\varrho = i e \left(\pi^* \varphi^* - \pi \varphi\right) \qquad s_i = i e \left(\frac{\partial \varphi^*}{\partial x_i} \varphi - \frac{\partial \varphi}{\partial x_i} \varphi^*\right) \tag{4}$$

The current four vector j_{μ} has the components:

$$j_4 = i \, \varrho \quad \text{and} \quad j_i = s_i - 2 \, e^2 \, \varphi^* \, \varphi \, A_i$$
 (5)

The Schrödinger equation is:

$$i \frac{\partial \Psi}{\partial t} = (H_0 + H') \Psi$$
(6)

458

Vacuum polarization and the photon-self-energy.

In the interaction representation⁶) this becomes:

$$i \frac{\partial \Psi'}{\partial t} = H'(t) \Psi'$$
(7)

with
$$\Psi' = e^{iH_0 t} \Psi$$
 and $H'(t) = e^{iH_0 t} H' e^{-iH_0 t}$ (8)

In general, the time dependence of an operator F is given by

$$F(t) = e^{iH_0 t} F e^{-iH_0 t} \tag{9}$$

In this representation the canonically conjugate momenta become

$$\pi(t) = \frac{\partial \varphi^{*}(t)}{\partial t} \qquad \pi^{*}(t) = \frac{\partial \varphi(t)}{\partial t} \qquad (10)$$

while the commutation relations remain the same as in the interaction free case

$$[\varphi^{*}(x), \varphi(x')] = i \Delta (x - x')$$

[\varphi(x), \varphi(x')] = [\varphi^{*}(x), \varphi^{*}(x')] = 0 (11)

where x is written instead of x, y, z, t. With the aid of the notation:

$$s_{\nu}(t) = i e \left(\frac{\partial \varphi^{*}}{\partial x_{\nu}} \varphi - \frac{\partial \varphi}{\partial x_{\nu}} \varphi^{*} \right)$$
(12)

we obtain for the interaction energy:

$$H'(t) = \int d^3 x \left(-s_{\nu} A_{\nu} + e^2 \varphi^* \varphi \sum_{i=1}^3 A_i A_i \right)$$
(13)

while the current j_{μ} becomes:

$$j_{\mu}(x) = s_{\mu}(x) + 2e^{2}\varphi^{*}(x)\varphi(x)(\delta_{\mu 4}\delta_{\nu 4} - \delta_{\mu \nu})A_{\nu}(x).$$
(14)

The vacuum expectation value of the perturbed (due to the existence of the external field) current j_{μ} is:

$$\leq j_{\mu}(x) \geq_{\mathrm{vac}} = \frac{i}{2} \int d^{4} x' \varepsilon (x - x') \leq [s_{\mu}(x), s_{\nu}(x')] \geq_{\mathrm{vac}} A_{\nu}(x')$$

+ $e^{2} \Delta^{(1)}(0) (\delta_{\mu 4} \delta_{\nu 4} - \delta_{\mu \nu}) A_{\nu}(x)$ (15)

The commutator in (15) yields

$$\frac{i}{2} \varepsilon (x - x') < [s_{\mu}, s'_{\nu}] >_{\text{vac}} = e^{2} \left\{ \frac{\partial \overline{\Delta}}{\partial x_{\mu}} \frac{\partial \Delta^{(1)}}{\partial x_{\nu}} + \frac{\partial \overline{\Delta}}{\partial x_{\nu}} \frac{\partial \Delta^{(1)}}{\partial x_{\mu}} - \frac{\partial^{2} \overline{\Delta}}{\partial x_{\mu} \partial x_{\nu}} \Delta^{(1)} - \overline{\Delta} \frac{\partial^{2} \Delta^{(1)}}{\partial x_{\mu} \partial x_{\nu}} \right\} - e^{2} \delta^{(4)} (x - x') \Delta^{(1)} (x - x') \delta_{\mu 4} \delta_{\nu 4}$$
(16)

Here use has been made of the identity:

$$\varepsilon \frac{\partial^2 \varDelta}{\partial x_{\mu} \partial x_{\nu}} = \frac{\partial^2}{\partial x_{\mu} \partial x_{\nu}} \varepsilon (x) \varDelta (x) - 2 \delta^{(4)} (x) \delta_{\mu 4} \delta_{\nu 4}$$

which, together with the definition of the Δ function, easily yields $(16)^*$).

Therefore

$$\langle j_{\mu}(x) \rangle_{\text{vac}} = e^{2} \int d^{4} x' \left(\frac{\partial \bar{\Delta}}{\partial x_{\mu}} \frac{\partial \Delta^{(1)}}{\partial x_{\nu}} + \frac{\partial \bar{\Delta}}{\partial x_{\nu}} \frac{\partial \Delta^{(1)}}{\partial x_{\mu}} - \frac{\partial^{2} \bar{\Delta}}{\partial x_{\mu} \partial x_{\nu}} \Delta^{(1)} \right)$$
$$- \bar{\Delta} \frac{\partial^{2} \Delta^{(1)}}{\partial x_{\mu} \partial x_{\nu}} A_{\nu}(x') - e^{2} \Delta^{(1)}(0) A_{\mu}(x)$$
(17)

or

$$< j_{\mu}(x) >_{\rm vac} = -e^2 \int d^4 x' K^{(0)}_{\mu\nu}(x-x') A_{\nu}(x')$$

with

$$K_{\mu\nu}^{(0)}(x) = -\frac{\partial \bar{\varDelta}(x)}{\partial x_{\mu}} \frac{\partial \varDelta^{(1)}(x)}{\partial x_{\nu}} - \frac{\partial \bar{\varDelta}(x)}{\partial x_{\nu}} \frac{\partial \varDelta^{(1)}(x)}{\partial x_{\mu}} + \frac{\partial^{2} \bar{\varDelta}(x)}{\partial x_{\mu} \partial x_{\nu}} \varDelta^{(1)}(x) + \bar{\varDelta}(x) \frac{\partial^{2} \varDelta^{(1)}(x)}{\partial x_{\mu} \partial x_{\nu}} + \varDelta^{(1)}(x) \delta^{(4)}(x) \delta_{\mu\nu}.$$
(18)

With the aid of the equations:

$$\Box \Delta^{(1)} - M^2 \Delta^{(1)} = 0 \quad \text{and} \quad \Box \overline{\Delta} - M^2 \overline{\Delta} = -\delta^4 \tag{19}$$

(18) may be put into the form (2'').

*) The arguments of $\overline{\varDelta}$ and $\varDelta^{(1)}$ are in (16): x - x'.

The condition for the gauge invariance of the total vacuum polarization current is:

$$\frac{\partial K_{\mu\nu}}{\partial x_{\nu}} = 0 \tag{20}$$

the separate terms, however, yield:

$$\frac{\partial K_{\mu\nu}^{(1/2)}(m_i)}{\partial x_{\nu}} = -4 \,\delta^{(4)}(x) \,\frac{\partial \Delta^{(1)}(m_i)}{\partial x_{\mu}}$$

$$\frac{\partial K_{\mu\nu}^{(0)}(M_i)}{\partial x_{\mu\nu}} = 2 \,\delta^{('')}(x) \,\partial \Delta^{(1)}(M_i)$$
(21)

and

$$\frac{\partial K^{(0)}_{\mu\nu}(M_i)}{\partial x_{\nu}} = 2 \,\delta^{(\prime\prime)}(x) \,\frac{\partial \Delta^{(1)}(M_i)}{\partial x_{\mu}}$$

whose right hand sides are not well defined at the origin¹). But the total $K_{\mu\nu}$ yields:

$$\frac{\partial K_{\mu\nu}}{\partial x_{\nu}} 2 \,\delta^{(4)}(x) \,\frac{\partial}{\partial x_{\mu}} \left(\sum_{i=1}^{N} \Delta^{(1)}(M_{i}) - 2 \sum_{i=1}^{n} \Delta^{(1)}(m_{i}) \right) \tag{22}$$

where the right hand side is regular and vanishes identically if two following conditions are satisfied:

$$N = 2 n$$
 I

$$\sum_{i=1}^{N} M_i^2 = 2 \sum_{i=1}^{n} m_i^2 \cdot$$
 II

These two conditions show a remarkable analogy with the conditions for regularization and are sufficient for the vanishing of the photon self energy. The mathematically simplest way to satisfy (I) and (II) is to take two scalar fields and to put their masses equal $M_1 = M_2$ = m. More general solutions exist too: we may introduce more than one spinor fields and assume the masses m_i and M_i very large except the mass of the electron m_1 so that the auxiliary charged particles would play a role in the extreme relativistic region only.

Next we proceed to the evaluation of the vacuum polarization current. We may FOURIER analyse the tensor $K_{\mu\nu}(x)$:

$$K_{\mu\nu}(x) = \frac{1}{(2\pi)^4} \int d^4 k \, K_{\mu\nu}(k) \, e^{ik_\lambda x_\lambda} \tag{23}$$

with

$$\begin{aligned} K^{(1/2)}_{\mu\nu}(k) &= \frac{4}{(2\pi)^3} \int d^4 \, p \left[+ p_{\mu} \left(p_{\nu} - k_{\nu} \right) + p_{\nu} \left(p_{\mu} - k_{\mu} \right) - \delta_{\mu\nu} \left(p_{\lambda} \left(p_{\lambda} - k_{\lambda} \right) \right) \\ &+ m^2 \right) \right] \frac{\delta \left(p_{\lambda} p_{\lambda} + m^2 \right)}{\left(p_{\lambda} - k_{\lambda} \right) \left(p_{\lambda} - k_{\lambda} \right) + m^2} \end{aligned} \tag{23'}$$

Res Jost and Jerzy Rayski.

$$\begin{split} K^{(0)}_{\mu\nu}(k) &= \frac{1}{(2\pi)^3} \int d^4 \, p \left[-p_{\mu} \left(p_{\nu} - k_{\nu} \right) - p_{\nu} \left(p_{\mu} - k_{\mu} \right) \right. \\ &- p_{\mu} \, p_{\nu} - \left(p_{\mu} - k_{\mu} \right) \left(p_{\nu} - k_{\nu} \right) + \delta_{\mu\nu} \left(p_{\lambda} p_{\lambda} + \left(p_{\lambda} - k_{\lambda} \right) \left(p_{\lambda} - k_{\lambda} \right) \right. \\ &+ 2 \, M^2 \right) \left] \frac{\delta \left(p_{\lambda} p_{\lambda} + M^2 \right)}{\left(p_{\lambda} - k_{\lambda} \right) \left(p_{\lambda} - k_{\lambda} \right) + M^2} \end{split}$$
(23")

Here use has been made of the FOURIER representations of the $\overline{\Delta}$ and $\Delta^{(1)}$ functions⁴). We introduce a new variable:

$$p' = p - \frac{1}{2}k$$

and denote for brevity:

$$p = p' + \frac{1}{2}k = p^{(1)}, \quad p - k = p' - \frac{1}{2}k = -p^{(2)}$$

Now (23') and (23'') assume the form:

$$\begin{split} K^{(1/_{2})}_{\mu\nu}(k) &= \frac{4}{(2\pi)^{3}} \int d^{4} p \left(-p^{(1)}_{\mu} p^{(2)}_{\nu} - p^{(1)}_{\nu} p^{(2)}_{\mu} \right. \\ &+ \left. \delta_{\mu\nu} \cdot \left(p^{(1)}_{\lambda} p^{(2)}_{\lambda} - m^{2} \right) \right) \frac{\delta \left(p^{(1)}_{\lambda} p^{(1)}_{\lambda} + m^{2} \right)}{p^{(2)}_{\lambda} p^{(2)}_{\lambda} + m^{2}} \quad (24') \end{split}$$

$$\begin{split} K^{(0)}_{\mu\nu}(k) &= \frac{1}{(2\pi)^3} \int d^4 \, p \left(p^{(1)}_{\mu} \, p^{(2)}_{\nu} + p^{(1)}_{\nu} \, p^{(2)}_{\mu} - p^{(1)}_{\mu} \, p^{(1)}_{\nu} - p^{(2)}_{\mu} \, p^{(2)}_{\nu} \right. \\ &+ \delta_{\mu\nu} \cdot \left(p^{(1)}_{\lambda} \, p^{(1)}_{\lambda} + p^{(2)}_{\lambda} \, p^{(2)}_{\lambda} + 2 \, M^2 \right) \right) \frac{\delta \left(p^{(1)}_{\lambda} \, p^{(1)}_{\lambda} + M^2 \right)}{p^{(2)}_{\lambda} \, p^{(2)}_{\lambda} + M^2} \quad (24'') \end{split}$$

where p has been written as integration variable instead of p' for convenience sake. The above integrals may be symmetrized by writing under the sign of integration:

$$U = \frac{1}{2} \left(\frac{\delta \left(p_{\lambda}^{(1)} p_{\lambda}^{(1)} + m^2 \right)}{p_{\lambda}^{(2)} p_{\lambda}^{(2)} + m^2} + \frac{\delta \left(p_{\lambda}^{(2)} p_{\lambda}^{(2)} + m^2 \right)}{p_{\lambda}^{(1)} p_{\lambda}^{(1)} + m^2} \right) \text{ instead of } \frac{\delta \left(p_{\lambda}^{(1)} p_{\lambda}^{(1)} + m^2 \right)}{p_{\lambda}^{(2)} p_{\lambda}^{(2)} + m^2}$$

which obviously does not change the values of the integrals in question but has the advantage of permitting the replacement of

$$U = \frac{1}{2} \left(\frac{\delta \left(p_{\lambda} p_{\lambda} + k_{\lambda} p_{\lambda} + m^2 + \frac{1}{4} k_{\lambda} k_{\lambda} \right)}{-2 k_{\lambda} p_{\lambda}} + \frac{\delta \left(p_{\lambda} p_{\lambda} - k_{\lambda} p_{\lambda} + m^2 + \frac{1}{4} k_{\lambda} k_{\lambda} \right)}{2 k_{\lambda} p_{\lambda}} \right)$$

by 7)

$$-\frac{1}{4}\int_{-1}^{1} du \,\delta'\left(p_{\lambda}p_{\lambda}+m^{2}+\frac{1}{4}\,k_{\lambda}k_{\lambda}+k_{\lambda}p_{\lambda}u\right). \tag{25}$$

We introduce again new variables

$$p' = p - \frac{1}{2} k u \tag{26}$$

whence

$$p^{(1)} = \frac{1}{2} k (1 + u) + p', \quad p^{(2)} = \frac{1}{2} k (1 - u) - p'$$
 (26')

which converts the argument of the δ' function in (25) into:

$$\delta' \left(p_{\lambda} p_{\lambda} + m^2 + \frac{1}{4} k_{\lambda} k_{\lambda} (1 - u^2) \right). \tag{25'}$$

Now we introduce (25) and (26) into (24) and split $K_{\mu\nu}$ into two parts:

$$K_{\mu\nu}(k) = \overline{K}_{\mu\nu}(k) + J_{\mu\nu}(k) \tag{27}$$

with:

$$\overline{K}_{\mu\nu}^{(1/2)}(k) = \frac{1}{2} \frac{1}{(2\pi)^3} \left(k_{\mu} k_{\nu} - k^2 \,\delta_{\mu\nu} \right) \int d^4 p \int_{-1}^{1} d u \left(1 - u^2 \right) \,\delta' \left(p^2 + m^2 + \frac{1}{4} \,k^2 \left(1 - u^2 \right) \right)$$
(28')

$$J_{\mu\nu}^{(1/2)}(k) = -\frac{2}{(2\pi)^3} \int d^4 p \int_{-1}^{1} du \left\{ p_{\mu} p_{\nu} - \frac{1}{2} \left(p^2 + m^2 + \frac{1}{4} k^2 \left(1 - u^2 \right) \right) \delta_{\mu\nu} \right\} \\ \delta' \left(p^2 + m^2 + \frac{1}{4} k^2 \left(1 - u^2 \right) \right)^*$$
(28")

and

$$\overline{K}^{(0)}_{\mu\nu}(k) = \frac{1}{4 (2 \pi)^3} \left(k_{\mu} k_{\nu} - k^2 \,\delta_{\mu\nu} \right) \int d^4 p \int_{-1}^{1} d \, u \, u^2 \, \delta' \left(p^2 + M^2 + \frac{1}{4} \, k^2 \left(1 - u^2 \right) \right)$$
(29')

$$J_{\mu\nu}^{(0)}(k) = \frac{1}{(2\pi)^3} \int d^4 p \int_{-1}^{1} du \left\{ p_{\mu} p_{\nu} - \frac{1}{2} \left(p^2 + M^2 + \frac{1}{4} k^2 (1 - u^2) \right) \delta_{\mu\nu} \right\} \\ \delta' \left(p^2 + M^2 + \frac{1}{4} k^2 (1 - u^2) \right)$$
(29")

*) As one sees from what follows, it is not permissible to put $[p^2+m^2+\frac{1}{4}k^2(1-u^2)] \delta'(p^2+m^2+\frac{1}{4}k^2(1-u^2)) = -\delta(p^2+m^2+\frac{1}{4}k^2(1-u^2))$, which, however, makes the non-gauge-invariant term vanish.

Here k^2 and p^2 have been written instead of $k_{\lambda}k_{\lambda}$ and $p_{\lambda}p_{\lambda}$ for brevity. We discuss first the part $J_{\mu\nu}$. Denoting

$$A = M^2 + rac{1}{4} \, k^2 \, (1 - u^2)$$

we may write:

$$\begin{split} \int\!d^{4}\,p\,\{p_{\mu}\,p_{\nu} - \frac{1}{2}\,(p^{2} + A)\,\delta_{\mu\,\nu}\}\delta'\,(p^{2} + A) = \frac{i}{2\,\pi}\!\!\!\!\int\limits_{-\infty}^{+\infty}\!\!\!\!dz\,z\!\int\!d^{4}\,p\,\{p_{\mu}\,p_{\nu} - \frac{1}{2}\,(p^{2} + A)\,\delta_{\mu\,\nu}\}e^{i\,(p^{2} + A)\,z}. \end{split}$$

With the aid of the formulae:

$$\int d^4 p \, e^{ip_{\lambda} p_{\lambda} z} = \frac{i \, \pi^2}{z^2} \, \varepsilon \, (z) \qquad \int d^4 p \, p_{\mu} p_{\nu} \, e^{ip_{\lambda} p_{\lambda} z} = - \, \delta_{\mu \nu} \frac{\pi^2}{2 \, z^3} \, \varepsilon \, (z) \quad (30)$$

the above integral yields

$$\begin{array}{c} \displaystyle -\frac{i\,\delta_{\mu\nu}\pi}{4} \int\limits_{-\infty}^{\infty} dz\,\varepsilon\left(z\right)\left(\frac{1}{z^2} - \frac{i\,A}{z}\right)e^{i\,A\,z} = -\frac{i\,\delta_{\mu\nu}}{4}\pi \int\limits_{-\infty}^{\infty} dz\,\varepsilon\left(z\right)\,\frac{d}{dz}\frac{e^{i\,A\,z}}{z} \\ \\ \displaystyle = \frac{i\pi}{2}\,\delta_{\mu\nu}\frac{e^{i\,A\,z}}{z}_{[z\,=\,0]} \end{array}$$

whence

$$J_{\mu\nu}(k) = \frac{i\pi}{2(2\pi)^3} \int_{-1}^{1} du \left\{ \frac{e^{ik^2/4}(1-u^2)z}{z} \left(\sum_{i=1}^{N} e^{iM_i^2 z} - 2\sum_{i=1}^{n} e^{im_i^2 z} \right) \right\}_{[z=0]}$$

It is immediately seen from (I) and (II) that this integral vanishes identically*). Only the gauge invariant part of $\overline{K}_{\mu\nu}(k)$ from (28) and (29) remains. We integrate $\overline{K}^{(0)}_{\mu\nu}(k)$ by parts:

$$\int_{-1}^{1} du \, u^{2} \, \delta' \left(p^{2} + M^{2} + \frac{1}{4} k^{2} \left(1 - u^{2} \right) \right) = \frac{u^{3}}{3} \, \delta' \left(p^{2} + M^{2} + \frac{1}{4} k^{2} \left(1 - u^{2} \right) \right) \Big|_{-1}^{+1} + \frac{k^{2}}{6} \int_{-1}^{1} du \, u^{4} \, \delta'' \left(p^{2} + M^{2} + \frac{1}{4} k^{2} \left(1 - u^{2} \right) \right)$$

*) Elementary integration of:

$$I_{\mu\nu} = \int d^4 p \left\{ p_{\mu} p_{\nu} - \frac{1}{2} \left(p^2 + A \right) \delta_{\mu\nu} \right\} \delta'(p^2 + A)$$

by performing the integration over p_0 first and with a large radius R gives (for A > 0):

$$I_{00} = 0, I_{11} = I_{22} = I_{33} = \frac{R^3}{\sqrt{R^2 + A}} = R^2 - \frac{1}{2}A + \dots$$

which leads to the same conditions for vanishing of $J_{\mu\nu}$.

whence

$$\begin{split} \overline{K}^{(0)}_{\mu\nu}(k) &= \frac{1}{4\,(2\,\pi)^3}\,(k_\mu\,k_\nu - k^2\,\delta_{\mu\nu})\int\!d^4\,p \left\{\!\frac{2}{3}\,\delta'\,(p^2 + M^2) \right. \\ &\left. + \frac{1}{6}k^2\int\limits_{-1}^1\!d\,u\,u^4\,\delta''\left(p^2 + M^2 + \frac{1}{4}k^2\,(1-u^2)\right)\right\}\,. \end{split}$$

The first term yields from (30):

$$-\frac{1}{(2\pi)^{3}}\frac{\pi}{12}(k_{\mu}k_{\nu}-k^{2}\delta_{\mu\nu})\int_{-\infty}^{+\infty}dz\frac{\varepsilon(z)}{z}e^{iM^{2}z}$$
(31')

while the second term may be integrated with the aid of the formula¹):

$$\int d^4 p \, \delta''(p_{\lambda}p_{\lambda}+A) = \pi/A$$

giving:

$$\frac{1}{(2\pi)^3} \frac{\pi}{24} k^2 (k_{\mu} k_{\nu} - k^2 \delta_{\mu\nu}) P \int_{-1}^{1} du \frac{u^4}{M^2 + \frac{1}{4} k^2 (1 - u^2)} = K^{(0)}_{\mu\nu}(k)_{\text{(regular)}}$$
(31")

In the same way $\overline{K}_{\mu\nu}^{(1/2)}$ yields:

$$-\frac{1}{(2\pi)^3}\frac{\pi}{3}(k_{\mu}k_{\nu}-k^2\delta_{\mu\nu})\int\limits_{-\infty}^{+\infty}dz\frac{\varepsilon(z)}{z}e^{im^2z}$$
(32')

and

$$\frac{1}{(2\pi)^3} \frac{\pi}{4} k^2 (k_{\mu} k_{\nu} - k^2 \delta_{\mu\nu}) P \int_{-1}^{+1} du \frac{u^2 \left(1 - \frac{u^2}{3}\right)}{m^2 + \frac{1}{4} k^2 (1 - u^2)} = K^{(1/2)}_{\mu\nu}(k)_{\text{(regular)}}$$
(32")

The parts (31') and (32') are logarithmically divergent and cannot compensate each other as the signs in both cases are equal. These terms are unphysical and have to be subtracted by means of a charge renormalization. The remaining terms (31'') and 32'') are finite and just responsible for the vacuum polarization effects.

In conclusions we state that the introduction of additional charged fields is sufficient to secure the gauge invariance of the vacuum polarization current and to yield a value zero for the photon self energy. On the other hand there remains the difficulty with the charge renormalization in an external field. This result contributes to the notion that the problem of the self charge is quite different from the problem of the self energy, and the analogy between the

30

procedures of mass and charge renormalizations may be more or less superficial.

Acknowledgement. The present authors wish to express their gratitude to Professor W. PAULI for his kind interest in this work as well as for much valuable aid.

References:

¹) W. PAULI & F. VILLARS, "On Invariant Regularization", Rev. M. Phys. in press.

²) H. UMEZAWA, J. YUKAWA, E. YAMADA, Progr. Theor. Phys. 3, 317 (1948).

³) J. RAYSKI, Acta Phys. Pol. IX, 129 (1948).

4) J. SCHWINGER, Phys. Rev. 74, 1439 (1948) and 75, 651 (1949).

⁵) G. WENTZEL, Quantentheorie d. Wellenfelder (1943).

⁶) S. TOMONAGA, Progr. Theor. Phys. 1, 27 (1946).

⁷) J. SCHWINGER, Quantumelectrodynamics, part III, in press.