

Zeitschrift: Helvetica Physica Acta
Band: 22 (1949)
Heft: VI

Artikel: Higher Approximation in the external field for the Problem of Vacuum Polarization
Autor: Källén, G.
DOI: <https://doi.org/10.5169/seals-112021>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Higher Approximations in the external field for the Problem of Vacuum Polarization

by G. Källén.

Swiss Federal Institute of Technology, Zurich, Switzerland*).

(13. VIII. 1949.)

Summary. The convergence and gauge invariance of the integrals appearing in the higher approximations of vacuum polarization are here discussed for the cases of spinor and scalar fields. Only the comparatively simple case of an external electromagnetic field is treated. The more complicated problem of a quantized field is not discussed.

As a direct result of the charge symmetry the vacuum expectation values of the current commutators vanish if an odd number of current operators are commuted. Hence the terms proportional to e^{2n+1} are identically zero, and in this form the statement is true for both the electrons and the bosons. Of the remaining terms it is shown that only the e^2 and e^4 approximations diverge, but that the still higher terms are both convergent and gauge invariant. It further appears that, apart from a numerical factor, which is the same in both approximations (and equal to $-\frac{1}{2}$), the strongest divergences are the same for the spinor and the scalar fields. The e^2 approximation has previously been treated by JOST and RAYSKI, who have shown that the non-gauge invariant (and divergent) terms compensate each other if one uses a suitable mixture of spinor and scalar fields. In this approximation, however, the logarithmically divergent charge renormalisation remains. The conditions of JOST and RAYSKI are

$$N = 2n \quad (I)$$

$$\sum_{i=1}^N M_i^2 = 2 \sum_{i=1}^n m_i^2 \quad (II)$$

where N and n are the number of scalar and spinor fields respectively and M_i and m_i are the corresponding masses. In the e^4 approximation the condition (I) alone is sufficient to secure a convergent and gauge invariance result.

*) At leave from: Department of Mechanics and Mathematical Physics, University of Lund, Sweden.

Introduction.

The e^2 approximation of the vacuum polarization has already been treated by several authors¹⁾ with the aid of the explicitly relativistically invariant quantum dynamics developed by SCHWINGER²⁾ and others. It has been shown by WENTZEL¹⁾ that it is possible to obtain different results for the photon self energy (which can be considered as a special case of the more general problem of the vacuum polarization) if one uses different methods of computing the integrals which appear in the formulae for the vacuum polarization. This question of uniqueness has been further discussed by PAULI and VILLARS¹⁾. In their paper the latter authors have given a method of invariant regularization of the different Δ -functions with the help of auxiliary masses. These masses, however, are only regarded as a mathematical aid for the computation and in the final result they are allowed to tend to infinity. It has been observed independently by RAYSKI and UMEZAWA³⁾ that it is also possible to regard these masses as observable if one assumes that the corresponding particles obey Bose-statistics. Detailed calculations by JOST and RAYSKI¹⁾ in the e^2 approximation have given as a result that the necessary assumptions in these realistic theories show a remarkable analogy with the more formalistical conditions of PAULI and VILLARS. We wish to extend the work of JOST and RAYSKI to the higher approximations in the fine-structure-constant and have as general equations

$$i \frac{\delta \psi[\sigma]}{\delta \sigma(x)} = H(x) \psi[\sigma] \quad (1)$$

$$H(x) = H_F(x) + H_B(x) \quad (1a)$$

$$H_F(x) = - \sum j_{\mu_F}(x) A_{\mu}(x) \quad (2)$$

$$j_{\mu_F}(x) = \frac{ie}{2} [\bar{\psi}(x) \gamma_{\mu} \psi(x) - \psi(x) \gamma_{\mu}^T \bar{\psi}(x)] \quad (2a)$$

$$H_B(x) = - \sum t_{\mu}(x) A_{\mu}(x) \quad (3)$$

¹⁾ E. g. J. SCHWINGER, Phys. Rev. **75**, 651 (1949); G. WENTZEL, Phys. Rev. **74**, 1070 (1948); W. PAULI, F. VILLARS, Rev. Mod. Phys. **21**, 434 (1949); R. JOST, J. RAYSKI, Helv. Phys. Acta **22**, 457 (1949). For earlier work without use of invariant formalism a summary has been given by V. WEISSKOPF, Det. Kgl. Danske Vid. Selskab, XIV 6 (1936).

²⁾ TOMONAGA, Progr. Theor. Phys. **1**, 27 (1946); J. SCHWINGER, Phys. Rev. **74**, 1439 (1948); *ibid.* **75**, 651 (1949); *ibid.* **76**, 790 (1949); F. J. DYSON: Phys. Rev. **75**, 486 (1949).

³⁾ J. RAYSKI, Acta Physica Polonica **9**, 129 (1948); H. UMEZAWA, J. YUKAWA, E. YAMADA, Progr. Theor. Phys. **3**, 317 (1948).

$$t_\mu(x) = ie s_\mu(x) + \xi_{\mu\nu} A_\nu(x) \cdot e^2 \cdot \varphi^*(x) \varphi(x) \quad (3a)$$

$$s_\mu(x) = \frac{\partial \varphi^*(x)}{\partial x_\mu} \varphi(x) - \frac{\partial \varphi(x)}{\partial x_\mu} \varphi^*(x) \quad (3b)$$

$$\xi_{\mu\nu} = \delta_{4\mu} \cdot \delta_{4\nu} - \delta_{\mu\nu} \quad (3c)$$

Here⁴⁾ $\psi(x)$ and $\bar{\psi}(x) = \psi^*(x) \gamma_4$ are the spinor fields and $\varphi(x)$ and $\varphi^*(x)$ the scalar fields⁵⁾. The symbols $t_\mu(x)$, $s_\mu(x)$ and $\xi_{\mu\nu}$ are defined by equations (3a) (3b) and (3c). The current operator for the scalar field is given by⁶⁾

$$j_{\mu B}(x) = ie s_\mu(x) + 2 \xi_{\mu\nu} A_\nu(x) \cdot e^2 \varphi^*(x) \varphi(x) \quad (4)$$

The total current is

$$j_\mu(x) = \sum j_{\mu F}(x) + \sum j_{\mu B}(x) \quad (4a)$$

It may be observed that none of the expressions (3a), (3b) or (3c) are tensors and that only the current $j_\mu(x)$ is a vector. The $A_\mu(x)$ are the four-dimensional vector potentials for the external electromagnetic field. They are here considered as given functions of space and time and not as operators. Hence we neglect the modification of the electromagnetic field due to polarization phenomena. The summations in equations (2) (3) and (4a) are to be extended over the spinor and scalar fields present.

The operators $\psi(x)$ and $\varphi(x)$ satisfy the following relations⁴⁾

$$\{\psi_\alpha(x); \bar{\psi}_\beta(x')\} = -i S_{\alpha\beta}(x-x') \quad (5)$$

$$\{\psi_\alpha(x); \psi_\beta(x')\} = \{\bar{\psi}_\alpha(x); \bar{\psi}_\beta(x')\} = 0 \quad (5a)$$

$$[\varphi^*(x); \varphi(x')] = i \Delta(x-x') \quad (6)$$

$$[\varphi^*(x); \varphi^*(x')] = [\varphi(x); \varphi(x')] = 0 \quad (6a)$$

Different kinds of fields always commute.

⁴⁾ The notations are essentially the same as those used by SCHWINGER but with natural units ($\hbar = c = 1$).

⁵⁾ Computations by FELDMANN have shown that it is not possible to compensate also the divergent charge renormalisation term by including fields with spin 1 (Unpublished letter to professor PAULI). The same result has also been gotten by H. UMEZAWA, R. KAWABE, Prog. Theor. Phys. (in press).

⁶⁾ G. WENTZEL: Quantentheorie der Wellenfelder.

We want to study the modified current operator, the expectation value of which may be written as²⁾

$$\begin{aligned} \langle j_\mu(x) \rangle &= \sum_{n=0}^{\infty} (i)^n \int_{-\infty}^{\sigma} dx' \int_{-\infty}^{\sigma'} dx'' \cdots \int_{-\infty}^{\sigma^{(n-1)}} dx^n \times \\ &\times \langle [H(x^n) [\cdots [H(x'); j_\mu(x)] \cdots]] \rangle_0 \end{aligned} \quad (7)$$

If the expressions (1a)–(4a) are substituted in equation (7) one gets a sum of commutators, some of which contains only one kind of fields (one of the $\psi(x)$ or one of the $\varphi(x)$ fields). The other terms contain at least one commutator between two different fields and are hence zero. This fact makes it possible to carry through the calculation for each field separately and in the end simply add the results together.

The Spinor fields.

We consider now only one of the spinor fields and write the corresponding parts of the current operator and of the hamiltonian as

$$j_\mu(x) = \frac{ie}{2} [\bar{\psi}(x); \gamma_\mu \psi(x)] \quad (8)$$

$$H(x) = -j_\mu(x) A_\mu(x) \quad (9)$$

In this case equation (7) gives

$$\begin{aligned} \langle j_\mu(x) \rangle &= \sum_{n=0}^{\infty} i^{-n} \int_{-\infty}^{\sigma} dx' \cdots \int_{-\infty}^{\sigma^{(n-1)}} dx^n A_{\nu_1}(x') \cdots A_{\nu_n}(x^n) \times \\ &\times \langle [j_{\nu_n}(x^n) [\cdots [j_{\nu_1}(x'); j_\mu(x)] \cdots]] \rangle_0 \end{aligned} \quad (10)$$

and our first task is to evaluate the iterated current commutators in this expression. If n is an even integer we have a commutator of an odd number of current operators, that is an expression of the following form

$$\begin{aligned} \left(\frac{ie}{2}\right)^{2n+1} \langle &[[\bar{\psi}(x^{2n}); \gamma_{\nu_{2n}} \psi(x^{2n})] [\cdots [[\bar{\psi}(x'); \gamma_{\nu_1} \psi(x')]; \\ &[\bar{\psi}(x); \gamma_\mu \psi(x)] \cdots]] \rangle_0 \end{aligned} \quad (11)$$

If we use the charge conjugate spinor $\psi'(x)$ equation (8) might as well be written as

$$j_\mu(x) = -\frac{ie}{2} [\bar{\psi}'(x); \gamma_\mu \psi'(x)] \quad (12)$$

and equation (11) as

$$-\left(\frac{ie}{2}\right)^{2n+1} \langle [\bar{\psi}'(x^{2n}); \gamma_{\nu_{2n}} \psi'(x^{2n})] [\dots [\bar{\psi}'(x'); \gamma_{\nu_1} \psi'(x')]; [\bar{\psi}'(x); \gamma_{\mu} \psi'(x)] \dots] \rangle_0 \quad (13)$$

As the two vacuum expectation values in (11) and (13) are equal we conclude that they are both zero and that equation (10) may be simplified to

$$\langle j_{\mu}(x) \rangle = \sum_{n=0}^{\infty} i^{2n-1} \int_{-\infty}^{\sigma} dx' \dots \int_{-\infty}^{\sigma^{2n}} dx^{2n+1} A_{\nu_1}(x') \dots A_{\nu_{2n+1}}(x^{2n+1}) \times \\ \times \langle [j_{\nu_{2n+1}}(x^{2n+1}) [\dots [j_{\nu_1}(x'); j_{\mu}(x)] \dots] \rangle_0 \quad (14)$$

If Ω is an arbitrary operator we have

$$[[\bar{\psi}(x); \Omega \psi(x')] \bar{\psi}_{\alpha}(x'')] = (\bar{\psi}(x) \Omega)_{\beta} \{ \psi_{\beta}(x'); \bar{\psi}_{\alpha}(x'') \} + \\ + \{ \bar{\psi}_{\alpha}(x''); \psi_{\beta}(x') \} (\Omega^T \bar{\psi}(x))_{\beta} = -2i (\bar{\psi}(x) \Omega S(x' - x''))_{\alpha} \quad (15)$$

In a similar way we get

$$[[\bar{\psi}(x); \Omega \psi(x')] \psi_{\alpha}(x'')] = 2i (S(x'' - x) \Omega \psi(x'))_{\alpha} \quad (16)$$

and hence

$$[[\bar{\psi}(x); \Omega \psi(x')] j_{\mu}(x'')] = \frac{ie}{2} ([\bar{\psi}(x); \Omega \psi(x')] \bar{\psi}_{\alpha}(x'')) (\gamma_{\mu} \psi(x''))_{\alpha} + \\ + (\bar{\psi}(x'') \gamma_{\mu})_{\alpha} [[\bar{\psi}(x); \Omega \psi(x')] \psi_{\alpha}(x'')] - [[\bar{\psi}(x); \Omega \psi(x')] \psi_{\alpha}(x'')] \times \\ \times (\gamma_{\mu}^T \bar{\psi}(x''))_{\alpha} - (\psi(x'') \gamma_{\mu}^T)_{\alpha} [[\bar{\psi}(x); \Omega \psi(x')] \bar{\psi}_{\alpha}(x'')] = \\ = e ([\bar{\psi}(x); \Omega S(x' - x'') \gamma_{\mu} \psi(x'')] - \\ - [\bar{\psi}(x''); \gamma_{\mu} S(x'' - x) \Omega \psi(x')]) \quad (17)$$

From equation (17) it is immediately seen that the commutator

$$[j_{\nu_n}(x^n) [\dots [j_{\nu_1}(x') j_{\mu}(x)] \dots]]$$

consists of 2^{n+1} terms of the form

$$\pm \frac{ie^{n+1}}{2} [\bar{\psi}(x^i); \gamma_{\nu_i} S(x^i - x^j) \dots \gamma_{\nu_r} S(x^r - x^s) \gamma_{\nu_s} \psi(x^s)] \quad (18)$$

where $i; \dots r; s$ is some permutation of the numbers $0; 1; \dots n$. With the aid of the formula

$$\langle [\bar{\psi}_{\alpha}(x); \psi_{\beta}(x')] \rangle_0 = S_{\beta\alpha}^{(1)}(x' - x)$$

the vacuum expectation value of (18) is given by

$$\pm \frac{i e^{n+1}}{2} Sp[S^{(1)}(x^s - x^i) \gamma_{\nu_i} S(x^i - x^j) \cdots \gamma_{\nu_s} S(x^r - x^s) \gamma_{\nu_s}] \quad (19)$$

Hence each term in the series (14) may be written as a sum of 2^{2n+2} terms of the form

$$(-1)^n \left(\pm \frac{e^{2n+2}}{2} \right) \int_{-\infty}^{\sigma} dx' \cdots \int_{-\infty}^{\sigma^{2n}} dx^{2n+1} Sp[S^{(1)}(x^s - x^i) \gamma A(x^i) S(x^i - x^j) \cdots \gamma A(x^r) S(x^r - x^s) \gamma A(x^s)] \quad (20)$$

If we suppose that the external field does not allow any real pairs to be created we can transform the expression (20) into

$$(-1)^n \left(\pm \left(\frac{e}{2} \right)^{2n+2} \right) \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} dx' \cdots dx^{2n+1} \varepsilon(01) \varepsilon(12) \cdots \cdots \varepsilon(2n; 2n+1) Sp[S^{(1)}(si) \gamma A(i) \cdots S(rs) \gamma A(s)]. \quad (21)$$

After changing the notations and rearranging the terms in the traces it is always possible to write each term in equation (14) as

$$(-1)^n \left(\frac{e}{2} \right)^{2n+2} \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} dx' \cdots dx^{2n+1} (Sp[\gamma_{\mu} S^{(1)}(01) \gamma A(1) S(12) \cdots \cdots S(2n+1, 0)] E_1^{(2n+1)} + Sp[\gamma_{\mu} S(01) \gamma A(1) S^{(1)}(12) \cdots \cdots S(2n+1, 0)] E_2^{(2n+1)} + \cdots + Sp[\gamma_{\mu} S(01) \gamma A(1) S(12) \cdots \cdots S^{(1)}(2n+1, 0)] E_{2n+2}^{(2n+1)}) \quad (22)$$

where $E_1^{(2n+1)}; E_2^{(2n+1)}; \cdots E_{2n+2}^{(2n+1)}$ are some functions of the $\varepsilon(i, j)$. It will be proven in the appendix that the expression (22) may be written as

$$(-1)^n \frac{e^{2n+2}}{2} \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} dx' \cdots dx^{2n+1} ((Sp[\gamma_{\mu} S^{(1)}(01) \gamma A(1) \bar{S}(12) \cdots \cdots \bar{S}(2n+1, 0)] + Sp[\gamma_{\mu} \bar{S}(01) \gamma A(1) S^{(1)}(12) \cdots \bar{S}(2n+1, 0)] \cdots + \cdots + Sp[\gamma_{\mu} \bar{S}(01) \gamma A(1) \bar{S}(12) \cdots S^{(1)}(2n+1, 0)]) \quad (23)$$

and hence

$$\langle j_{\mu}(x) \rangle = \frac{1}{2} \sum_{n=0}^{\infty} (-1)^n e^{2n+2} \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} dx' \cdots dx^{2n+1} (Sp[\gamma_{\mu} S^{(1)}(01) \times \cdots \times \gamma A(1) \cdots \bar{S}(2n+1, 0)] + \cdots + Sp[\gamma_{\mu} \bar{S}(01) \gamma A(1) \cdots S^{(1)}(2n+1, 0)]) \quad (24)$$

In momentum space this formula reads

$$\begin{aligned} \langle j_\mu(x) \rangle = & \frac{1}{2} \sum_{n=0}^{\infty} (-1)^n e^{2n+2} \left(\frac{1}{2\pi} \right)^{8n+7} \int \dots \int dp dp' \dots dp^{2n+1} \times \\ & \times e^{ix(p-p^{2n+1})} Sp[\gamma_\mu (i\gamma p - m) \gamma A(p' - p) (i\gamma p' - m) \dots \\ & \dots \gamma A(p^{2n+1} - p^{2n}) (i\gamma p^{2n+1} - m)] \left(\frac{\delta(p^2 + m^2)}{(p'^2 + m^2) \dots (p^{2n+1^2} + m^2)} + \right. \\ & \left. + \frac{\delta(p'^2 + m^2)}{(p^2 + m^2) (p'^2 + m^2) \dots (p^{2n+1^2} + m^2)} + \dots + \frac{\delta(p^{2n+1^2} + m^2)}{(p^2 + m^2) \dots (p^{2n^2} + m^2)} \right) \end{aligned} \quad (25)$$

where

$$Av(p) = \int Av(x) e^{ipx} dx \quad (26)$$

The equations (24) and (25) give a formal expression for the expectation value of the current operator, but the questions of convergence and gauge invariance are still open.

We first want to give a formal proof of the gauge invariance in momentum space. It is true that this proof is only valid if the integrals converge but we will leave this question unsettled for the moment.

For this calculation we need the following formula

$$\begin{aligned} Sp[\Omega(i\gamma a - m) \gamma q (i\gamma(a + q) - m)] = & i[(a + q)^2 + m^2] Sp[\Omega(i\gamma a - m)] - \\ & - i[a^2 + m^2] Sp[\Omega(i\gamma(a + q) - m)] \end{aligned} \quad (27)$$

In equation (27) a and q are two arbitrary fourdimensional vectors, m a number and Ω an arbitrary operator. The formula (27) may be proven by an explicate calculation e. g. in the following way

$$\begin{aligned} Sp[\Omega(i\gamma a - m) \gamma q (i\gamma(a + q) - m)] = & - Sp[\Omega \cdot \gamma a \cdot \gamma q \cdot \gamma(a + q)] - \\ & - im (Sp[\Omega \cdot \gamma q \cdot \gamma(a + q)] + Sp[\Omega \cdot \gamma a \cdot \gamma q]) + m^2 Sp[\Omega] \end{aligned} \quad (28)$$

Using the equation

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2 \delta_{\mu\nu}$$

we get from (28)

$$\begin{aligned} & -2qa Sp[\Omega \cdot \gamma a] + a^2 Sp[\Omega \cdot \gamma q] - q^2 Sp[\Omega \cdot \gamma a] - \\ & - mi(q^2 + 2aq) Sp[\Omega] + m^2 Sp[\Omega] = i[(a + q)^2 + m^2] \times \\ & \times Sp[\Omega(i\gamma a - m)] - i(a^2 + m^2) Sp[\Omega(i\gamma(a + q) - m)] \end{aligned}$$

which is the right hand side of (27).

One typical term in (25) may be written as

$$\int \dots \int dp dp' \dots dp^n e^{ix(p-p^n)} Sp[\gamma_\mu (i\gamma p - m) \gamma A(p' - p) \dots \dots (i\gamma p^n - m)] \frac{\delta(p^{i^2} + m^2)}{(p^2 + m^2) \dots (p^{i-1^2} + m^2) (p^{i+1^2} + m^2) \dots (p^{n^2} + m^2)} \quad (29)$$

(In (29) n is supposed to be odd but for our present purpose we do not need this fact.)

If we make an infinitesimal gauge transformation

$$\left. \begin{aligned} A_\nu(x) &\rightarrow A_\nu(x) + \frac{\partial A(x)}{\partial x_\nu} \\ \varepsilon(p) &= \int e^{ixp} A(x) dx \end{aligned} \right\} \quad (30)$$

the integral (29) will be changed by the following amount

$$\begin{aligned} &\frac{1}{i} \int \dots \int dp \dots dp^n e^{ix(p-p^n)} \frac{\delta(p^{i^2} + m^2)}{(p^2 + m^2) \dots (p^{i-1^2} + m^2) (p^{i+1^2} + m^2) \dots (p^{n^2} + m^2)} \times \\ &\times (Sp[\gamma_\mu (i\gamma p - m) \gamma (p' - p) (i\gamma p' - m) \gamma A(p'' - p') \dots] \times \\ &\times \varepsilon(p' - p) + Sp[\gamma_\mu (i\gamma p - m) \gamma A(p' - p) (i\gamma p' - m) \times \\ &\times \gamma (p'' - p') \dots] \varepsilon(p'' - p') + \dots) \end{aligned} \quad (31)$$

Inserting equation (27) in formula (31) we get

$$\begin{aligned} &\int \dots \int dp \dots dp^n e^{ix(p-p^n)} \frac{\delta(p^{i^2} + m^2)}{(p^2 + m^2) \dots (p^{i-1^2} + m^2) (p^{i+1^2} + m^2) \dots (p^{n^2} + m^2)} \times \\ &\times [(p'^2 + m^2) Sp[\gamma_\mu (i\gamma p - m) \gamma A(p'' - p') \dots] \varepsilon(p' - p) - \\ &- (p^2 + m^2) Sp[\gamma_\mu (i\gamma p' - m) \gamma A(p' - p) \dots] \varepsilon(p' - p) + \\ &+ (p''^2 + m^2) Sp[\gamma_\mu (i\gamma p - m) \gamma A(p' - p) \dots] \times \varepsilon(p'' - p') - \\ &- (p'^2 + m^2) Sp[\gamma_\mu (i\gamma p - m) \gamma A(p' - p) \dots] \varepsilon(p'' - p') + \dots] \end{aligned} \quad (32)$$

With the notation

$$\int dp' A_\nu(p' - p) \varepsilon(p' - p'') = \int dq \varepsilon(q) A_\nu(p'' - p - q) = E_\nu(p'' - p) \quad (33)$$

and the formula

$$\int dp \varepsilon(p' - p) e^{ixp} = \left(\frac{1}{2\pi}\right)^4 A(x) e^{ixp'} \quad (34)$$

the expression (32) may be written (the terms containing $p^{i^2} + m^2$ give zero due to the factor $\delta(p^{i^2} + m^2)$).

$$\begin{aligned} & \int \dots \int dp' \dots dp^n e^{ix(p'-p'')} \frac{\delta(p^{i^2} + m^2)}{(p'^2 + m^2) \dots (p^{i-1^2} + m^2) (p^{i+1^2} + m^2) \dots (p^{n^2} + m^2)} \times \\ & \times [Sp[\gamma_\mu (i\gamma p' - m) \gamma E(p'' - p') (i\gamma p'' - m) \dots] - \left(\frac{1}{2\pi}\right)^4 A(x) \times \\ & \times Sp[\gamma_\mu (i\gamma p' - m) \gamma A(p'' - p') (i\gamma p'' - m) \dots] + \\ & + Sp[\gamma_\mu (i\gamma p' - m) \gamma A(p'' - p') (i\gamma p'' - m) \gamma E(p''' - p'') \dots] - \\ & - Sp[\gamma_\mu (i\gamma p' - m) \gamma E(p'' - p') (i\gamma p'' - m) \dots] + \dots \\ & \dots + \left(\frac{1}{2\pi}\right)^4 A(x) Sp[\gamma_\mu (i\gamma p' - m) \gamma A(p'' - p') \dots]] = 0. \end{aligned} \quad (35)$$

From (35) we conclude that each term in (25) if convergent is also gauge invariant.

To discuss the convergence properties of (25) we write the bracket as

$$\begin{aligned} & \frac{\delta(p^2 + m^2)}{(p'^2 - p^2) \dots (p^{n^2} - p^2)} + \frac{\delta(p'^2 + m^2)}{(p^2 - p'^2) (p''^2 - p'^2) \dots (p^{n^2} - p'^2)} + \dots \\ & \dots + \frac{\delta(p^{n^2} + m^2)}{(p^2 - p^{n^2}) \dots (p^{n-1^2} - p^{n^2})} \end{aligned} \quad (36)$$

We now use the formula

$$\begin{aligned} & \frac{1}{(p'^2 - p^2) \dots (p^{n^2} - p^2)} + \frac{1}{(p^2 - p'^2) \dots (p^{n^2} - p'^2)} + \dots \\ & \dots + \frac{1}{(p^2 - p^{n^2}) \dots (p^{n-1^2} - p^{n^2})} = 0 \end{aligned} \quad (37)$$

to write (36) as

$$\frac{\delta(p'^2 + m^2) - \delta(p^2 + m^2)}{(p^2 - p'^2) (p''^2 - p'^2) \dots (p^{n^2} - p'^2)} + \dots + \frac{\delta(p^{n^2} + m^2) - \delta(p^2 + m^2)}{(p^2 - p^{n^2}) \dots (p^{n-1^2} - p^{n^2})} \quad (38)$$

Following SCHWINGER this can also be written

$$\begin{aligned} & -\frac{1}{2} \int_{-1}^{+1} du \left[\frac{\delta' \left(\frac{1}{2} (p^2 + p'^2) + m^2 + \frac{1}{2} u (p^2 - p'^2) \right)}{(p''^2 - p'^2) \dots (p^{n^2} - p'^2)} + \dots \right. \\ & \left. \dots + \frac{\delta' \left(\frac{1}{2} (p^2 + p^{n^2}) + m^2 + \frac{1}{2} u (p^2 - p^{n^2}) \right)}{(p'^2 - p^{n^2}) \dots (p^{n-1^2} - p^{n^2})} \right] \end{aligned} \quad (39)$$

By repeating this process it is obviously possible to write (36) in the following form

$$\begin{aligned} & \left(-\frac{1}{2}\right)^{\frac{n(n+1)}{2}} \int_{-1}^{+1} du_1 (1-u_1)^{n-1} \int_{-1}^{+1} du_2 (1-u_2)^{n-2} \dots \int_{-1}^{+1} du_n \delta^{(n)} \left(m^2 + \right. \\ & \quad + \frac{p^2}{2} + \frac{p'^2}{4} \dots + \frac{p^{n-1^2}}{2^n} + \frac{p^{n^2}}{2^n} - u_1 \left(-\frac{p^2}{2} + \frac{p'^2}{4} + \frac{p''^2}{8} + \dots + \right. \\ & \quad + \frac{p^{n-1^2}}{2^n} + \frac{p^{n^2}}{2^n} \Big) - u_2 (1-u_1) \times \left(-\frac{p'^2}{4} + \frac{p''^2}{8} + \dots + \frac{p^{n-1^2}}{2^n} + \frac{p^{n^2}}{2^n} \right) - \\ & \quad \dots - u_n (1-u_{n-1}) \dots (1-u_1) \left(-\frac{p^{n-1^2}}{2^n} + \frac{p^{n^2}}{2^n} \right) \Big) \end{aligned} \quad (40)$$

It is now convenient to make a translation of the origins and write (25) in the following way

$$\begin{aligned} \langle j_\mu(x) \rangle &= \frac{1}{2} \sum_{n=0}^{\infty} e^{2n+2} (-1)^n \left(\frac{1}{2\pi} \right)^{8n+7} \int \dots \int dq' \dots dq^{2n+1} \times \\ & \quad \times e^{-ix(q' + \dots + q^{2n+1})} K_{\mu \nu_1 \dots \nu_{2n+1}}^{(2n+1)}(q' \dots q^{2n+1}) A_{\nu_1}(q') \dots \\ & \quad \dots A_{\nu_{2n+1}}(q^{2n+1}) \end{aligned} \quad (41)$$

where the kernel $K_{\mu \nu_1 \dots \nu_n}^{(n)}$ according to (40) is given by

$$\begin{aligned} K_{\mu \nu_1 \dots \nu_n}^{(n)}(q' \dots q^n) &= \left(-\frac{1}{2}\right)^{\frac{n(n+1)}{2}} \int dp \int_{-1}^{+1} du_1 (1-u_1)^{n-1} \int_{-1}^{+1} du_2 \times \\ & \quad \times (1-u_2)^{n-2} \dots \int_{-1}^{+1} du_n \delta^{(n)}(m^2 + p^2 + 2pQ + Q^2 + \varphi) Sp[\gamma_\mu \times \\ & \quad \times (i\gamma p - m) \gamma_{\nu_1} (i\gamma(p+q') - m) \gamma_{\nu_2} \times (i\gamma(p+q'+q'') - m) \dots \\ & \quad \dots \gamma_{\nu_n} (i\gamma(p+q'+\dots+q^n) - m)] \end{aligned} \quad (42)$$

Here

$$Q = (1-u_1) \frac{q'}{2} + (1-u_1)(1-u_2) \frac{q''}{4} + \dots + (1-u_1) \dots (1-u_n) \frac{q^n}{2^n} \quad (43)$$

and φ is a bilinear expression in $q' \dots q^n$.

Another translation of the origin transforms (42) into

$$\begin{aligned} K_{\mu \nu_1 \dots \nu_n}^{(n)}(q' \dots q^n) &= \left(-\frac{1}{2}\right)^{\frac{n(n+1)}{2}} \int_{-1}^{+1} du_1 (1-u_1)^{n-1} \dots \int_{-1}^{+1} du_n \int dq \times \\ & \quad \times \delta^{(n)}(q^2 + m^2 + \varphi) Sp[\gamma_\mu (i\gamma(q+a_1) - m) \gamma_{\nu_1} (i\gamma(q+a_2) - m) \dots \\ & \quad \dots \gamma_{\nu_n} (i\gamma(q+a_{n+1}) - m)] \end{aligned} \quad (44)$$

where $a_1 \dots a_{n+1}$ are linear combinations of $q' \dots q^n$ with coefficients which depend on $u_1 \dots u_n$.

We now suppose that the potentials $A_\nu(q)$ vanish for large momenta so that the integrations over $q' \dots q^{2n+1}$ in (41) converge and are thus left with only the q integration in (44). The strongest divergence in this integral is of the form

$$\int dq \delta^{(n)}(q^2 + \alpha^2) q_\mu q_{\nu_1} \dots q_{\nu_n} \quad (45)$$

With the well-known representation of the δ -function (45) may be written

$$\frac{1}{2\pi} \cdot i^n \int_{-\infty}^{+\infty} x^n dx \int dq e^{ix(q^2 + \alpha^2)} q_\mu q_{\nu_1} \dots q_{\nu_n} \quad (46)$$

The q -space integration in (46) can be performed without difficulty and gives some constant factor λ multiplied by $|x| \cdot x^{-\frac{n+1}{2}-3}$. The integral (46) can therefore be written as

$$\frac{\lambda}{2\pi} \cdot i^n \int_{-\infty}^{+\infty} |x| \cdot x^{\frac{n-7}{2}} e^{i\alpha^2 x} dx \quad (47)$$

The expression (47) and the kernel (44) converge for $n \geq 4$ but diverge for $n \leq 3$. As we are only interested in odd integers n , the only divergent integrals are

$$(n = 1) \int_{-\infty}^{+\infty} \frac{e^{i\alpha^2 x}}{x \cdot |x|} dx \quad (48a)$$

and

$$(n = 3) \int_{-\infty}^{+\infty} \frac{e^{i\alpha^2 x}}{|x|} dx \quad (48b)$$

The first one appears in the e^2 approximation and as this case has already been treated by several authors we limit ourselves to the e^4 approximation, where the integral (48b) is the only divergent one. The kernel is here

$$\begin{aligned} K_{\mu\nu_1\nu_2\nu_3}^{(3)}(q'q''q''') &= \frac{1}{64} \int_{-1}^{+1} du_1 (1-u_1)^2 \int_{-1}^{+1} du_2 (1-u_2) \int_{-1}^{+1} du_3 \times \\ &\times \int_{-\infty}^{+\infty} x^3 dx \int dq e^{ix(q^2 + m^2 + \varphi)} Sp[\gamma_\mu (i\gamma(q + a_1) - m) \dots \\ &\dots \gamma_{\nu_3} (i\gamma(q + a_4) - m)] \end{aligned} \quad (49)$$

and the divergent term

$$\frac{1}{64} \int_{-1}^{+1} du_1 (1-u_1)^2 \int_{-1}^{+1} du_2 (1-u_2) \int_{-1}^{+1} du_3 \int_{-\infty}^{-\infty} x^3 dx \int dq e^{ix(q^2+m^2+\varphi)} \times \\ \times Sp[\gamma_\mu \gamma q \gamma_{v_1} \gamma q \gamma_{v_2} \gamma q \gamma_{v_3} \gamma q] \quad (50)$$

The trace in (50) is easily evaluated and is equal to

$$4\{8q_\mu q_{v_1} q_{v_2} q_{v_3} - 2q^2(q_\mu q_{v_1} \delta_{v_2 v_3} + q_\mu q_{v_3} \delta_{v_1 v_2} + q_{v_1} q_{v_3} \delta_{\mu v_3} + \\ + q_{v_2} q_{v_3} \delta_{\mu v_1}) + q^4(\delta_{\mu v_1} \delta_{v_2 v_3} - \delta_{\mu v_2} \delta_{v_1 v_3} + \delta_{\mu v_3} \delta_{v_1 v_2})\}. \quad (51)$$

The scalar fields.

In this case equation (7) reads

$$\langle j_\mu(x) \rangle = \sum_{n=0}^{\infty} i^{-n} \int_{-\infty}^{\sigma} dx' \int_{-\infty}^{\sigma'} dx'' \cdots \int_{-\infty}^{\sigma^{(n-1)}} dx^n A_{v_1}(x') \cdots A_{v_n}(x^n) \times \\ \times \langle [t_{v_n}(x^n) [\cdots [t_{v_1}(x') j_\mu(x)] \cdots]] \rangle_0. \quad (52)$$

As both $t_v(x)$ and $j_\mu(x)$ here contain two different powers of e , formula (52) is not an expansion in powers of the charge. It is, however, always possible to rearrange the terms so as to get such an expansion. For this purpose we write

$$\langle j_\mu(x) \rangle = \sum_{n=0}^{\infty} \sum_{k=0}^{n+1} e^{n+k+1} \langle j_\mu(x) \rangle^{(n)(k)} = \sum_{n=0}^{\infty} e^{n+1} \langle j_\mu(x) \rangle^{(n)} \quad (53)$$

where

$$\langle j_\mu(x) \rangle^{(n)} = \sum_{k=0}^{\frac{n+1}{2}} \langle j_\mu(x) \rangle^{(n-k)(k)} \quad (53a)$$

for odd n , and

$$\langle j_\mu(x) \rangle^{(n)} = \sum_{k=0}^{\frac{n}{2}} \langle j_\mu(x) \rangle^{(n-k)(k)} \quad (53b)$$

when n is even. From equations (52) and (53) we also get

$$\langle j_\mu(x) \rangle^{(n)(k)} = \left(\frac{1}{2}\right)^n i^{1-k} \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} dx' \cdots dx^n \varepsilon(01) \varepsilon(12) \cdots \varepsilon(n-1, n) \times \\ \times A_{v_1}(x') \cdots A_{v_n}(x^n) \{ \langle [\varphi^*(x^n) \varphi(x^n) [\cdots [\varphi^*(x^{n-k+1}) \varphi(x^{n-k+1}) \times \\ \times [s_{v_{n-k}}(x^{n-k}) [\cdots [s_{v_1}(x') s_\mu(x)] \cdots]] \cdots]] \rangle_0 \xi(n-k+1, n+1) \cdots \\ \cdots \xi(n, n+k) A_{v_{n+1}}(x^{n-k+1}) \cdots A_{v_{n+k}}(x^n) + \cdots \} \quad (54)$$

The bracket in (54) contains one term for each way in which it is possible to pick out $n - k + 1$ indices $\nu_{i_1} \dots \nu_{i_{n-k+1}}$ from the series $\mu \nu_1 \dots \nu_n$. If all $\nu_i \neq \mu$ the corresponding term is multiplied by a factor 2.

From the charge symmetry we see here too that the series in (53) will contain only even powers of e and hence that n in equation (53a) is an odd integer.

By an argument analogous to that leading from equation (14) to equation (23) but with differential operators instead of γ -matrices it is seen that (for n odd)

$$\begin{aligned} & \left(\frac{1}{2}\right)^n \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} dx' \dots dx^n \varepsilon(01) \dots \varepsilon(n-1, n) A_{\nu_1}(x') \dots A_{\nu_n}(x^n) \\ & \langle [s_{\nu_n}(x^n) [\dots [s_{\nu_1}(x') s_{\mu}(x)] \dots]] \rangle_0 = \frac{i}{2} \left(\frac{1}{2\pi}\right)^{4n+3} \int \dots \\ & \dots \int dp dp' \dots dp^n e^{ix(p-p^n)} A_{\nu_1}(p'-p) \dots A_{\nu_n}(p^n - p^{n-1}) \times \\ & \times P^{(n)} \times \left\{ \frac{\delta(p^2 + m^2)}{(p'^2 - p^2) \dots (p^{n^2} - p^2)} + \frac{\delta(p'^2 + m^2)}{(p^2 - p'^2) \dots (p^{n^2} - p'^2)} + \dots \right. \\ & \left. \dots + \frac{\delta(p^{n^2} + m^2)}{(p^2 - p^{n^2}) \dots (p^{n-1^2} - p^{n^2})} \right\} \end{aligned} \quad (55)$$

where

$$\begin{aligned} P^{(n)} = \pi^{(n)} - \sum_j \delta_{4\nu_i} \delta_{4\nu_{i-1}} (p^{i-1^2} + m^2) \left[\pi_{i, i-1}^{(n)} - \frac{1}{2} \sum_j \delta_{4\nu_j} \delta_{4\nu_{j-1}} \times \right. \\ \left. \times (p^{j-1^2} + m^2) [\pi_{i, i-1, j, j-1}^{(n)} - \dots] \right] \end{aligned} \quad (56)$$

and

$$\pi^{(n)} = (p'_{\nu_1} + p_{\nu_1}) (p''_{\nu_2} + p'_{\nu_2}) \dots (p^n_{\nu_n} + p^{n-1}_{\nu_n}) (p_{\mu} + p^n_{\mu}) \quad (57)$$

$$\pi_{i, j, \dots}^{(n)} = \frac{\pi^{(n)}}{(p^i_{\nu_i} + p^{i-1}_{\nu_i}) (p^j_{\nu_j} + p^{j-1}_{\nu_j}) \dots} \quad (58)$$

$$\pi_{i, j, \dots}^{(n)} = 0 \text{ if two indices are equal.}$$

We here have also used the following formula given by JOST and RAYSKI

$$\varepsilon(x) \frac{\partial^2 \Delta(x)}{\partial x_{\mu} \partial x_{\nu}} = \frac{\partial^2 (\varepsilon(x) \Delta(x))}{\partial x_{\mu} \partial x_{\nu}} - 2 \delta(x) \delta_{4\mu} \delta_{4\nu} \quad (59)$$

The more general commutators in (54) can be evaluated in the same way. Putting

$$\begin{aligned} & A_{\nu_{n+i}}(n-k+i) A_{\nu_{n-k+i}}(n-k+i) = \\ & = \int \delta(n-k+i, n+i) A_{\nu_{n-k+i}}(n-k+i) A_{\nu_{n+i}}(n+i) dx^{n+i} \end{aligned} \quad (60)$$

we get a formula, which is similar to (56) but with ξ (ij) substituted for $\delta_4 v_i$ $\delta_4 v_{i-1}$ in some of the terms. The expression (53a) may now be written

$$\begin{aligned} \langle j_\mu(x) \rangle^{(2n+1)} &= \frac{1}{2} \left(\frac{1}{2\pi} \right)^{8n+7} (-1)^n \int \dots \\ &\dots \int dp dp' \dots dp^{2n+1} e^{ix(p-p^{2n+1})} \\ &\times \bar{P}_{\mu v_1 \dots v_{2n+1}}^{(2n+1)} \left\{ \frac{\delta(p^2+m^2)}{(p'^2-p^2) \dots (p^{n^2}-p^2)} + \dots + \frac{\delta(p^{n^2}+m^2)}{(p^2-p^{n^2}) \dots (p^{n-1^2}-p^{n^2})} \right\} \times \\ &\times A_{v_1}(p'-p) \dots A_{v_n}(p^n-p^{n-1}) \end{aligned} \quad (61)$$

where

$$\begin{aligned} \bar{P}_{\mu v_1 \dots v_n}^{(n)} &= \pi^{(n)} - \sum_i f_{v_i v_{i-1}}^{(1)} (p^{i-1^2} + m^2) \pi_{i, i-1}^{(n)} + \frac{1}{2} \sum_{i,j} f_{v_i v_{i-1} v_j v_{j-1}}^{(2)} \times \\ &\times (p^{i-1^2} + m^2) (p^{j-1^2} + m^2) \pi_{i, i-1, j, j-1}^{(n)} - \frac{1}{6} \sum \dots \end{aligned} \quad (62)$$

and $f_{v_i v_{i-1} v_j v_{j-1}}^{(k)}$ is some function of the $\delta_4 v_i$ and $\delta_{v_i v_{i-1}}$. As will be shown in the appendix, f is actually a tensor and equal to

$$f_{v_i v_{i-1} v_j v_{j-1}}^{(k)} = \delta_{v_i v_{i-1}} \delta_{v_j v_{j-1}} \dots \quad (63)$$

The formulae (62) and (63) express the lorentz invariance of equation (61).

To prove the gauge invariance (supposing the integrals in (61) to converge) we need an identity of the same type as equation (27). We have

$$\begin{aligned} \bar{P}_{\mu v_1 \dots v_n}^{(n)} (pp' \dots p^n) (p'_{v_1} - p_{v_1}) &= (p'^2 - p^2) \times \\ &\times [\pi_{1,0}^{(n)} - \sum_i \delta_{v_i v_{i-1}} (p^{i-1^2} + m^2) \pi_{1,i,i-1}^{(n)} + \dots] - (p^2 + m^2) \times \\ &\times (p'_\mu - p_\mu) [\pi_{1,0}^{(n)} - \sum_i \delta_{v_i v_{i-1}} (p^{i-1^2} + m^2) \pi_{1,0,i,i-1}^{(n)} + \dots] - \\ &- (p'^2 + m^2) (p'_{v_2} - p_{v_2}) [\pi_{1,2}^{(n)} - \sum_i \delta_{v_i v_{i-1}} (p^{i-1^2} + m^2) \times \\ &\times \pi_{1,2,i,i-1}^{(n)} + \dots] = (p'^2 + m^2) \bar{P}_{\mu v_2 \dots v_n}^{(n-1)} (pp'' \dots p^n) - \\ &- (p^2 + m^2) \bar{P}_{\mu v_2 \dots v_n}^{(n-1)} (p' p'' \dots p^n). \end{aligned} \quad (64)$$

We can now repeat the calculation from equation (29) to equation (35) but start from (61) instead of (25) and use (64) instead of (27). The result is obviously that, from this formal point of view, (61) is gauge invariant.

Writing (61) as (n odd)

$$\langle j_\mu(x)^{(n)} \rangle = \frac{1}{2} (-1)^{\frac{n+1}{2}} \left(\frac{1}{2\pi} \right)^{4n+3} \int \cdots \int dq' \cdots dq^n e^{-ix(q' + \cdots + q^n)} \times \\ \times \bar{K}_{\mu\nu_1 \cdots \nu_n}^{(n)}(q' \cdots q^n) A_{\nu_1}(q) \cdots A_{\nu_n}(q^n) \quad (65)$$

we have

$$\bar{K}_{\mu\nu_1 \cdots \nu_n}^{(n)}(q' \cdots q^n) = \left(-\frac{1}{2} \right)^{\frac{n(n+1)}{2}} \int_{-1}^{+1} du_1 (1-u_1)^{n-1} \cdots \int_{-1}^{+1} du_n \times \\ \times \int dq \delta^{(n)}(q^2 + m^2 + \varphi) \bar{P}_{\mu\nu_1 \cdots \nu_n}^{(n)}(q + a_1, q + a_2, \cdots) \quad (66)$$

(compare equation (44)).

Formula (66) converges or diverges as

$$\int dq \delta^{(n)}(q^2 + \alpha^2) q_\mu q_{\nu_1} \cdots q_{\nu_n} \quad (67)$$

which is the same equation as (45). We thus get exactly the same convergent cases as for the spinor field.

The divergent term of (66) in the case $n = 3$ may according to (62) and (46) be written

$$\frac{1}{64} \int_{-1}^{+1} du_1 (1-u_1)^2 \int_{-1}^{+1} du_2 (1-u_2) \int_{-1}^{+1} du_3 \int_{-\infty}^{+\infty} x^3 dx \int dq e^{ix(q^2 + m^2 + \varphi)} \times \\ \times \{ 16 q_\mu q_{\nu_1} q_{\nu_2} q_{\nu_3} - 4 q^2 (\delta_{\mu\nu_3} q_{\nu_1} q_{\nu_2} + \delta_{\nu_1\mu} q_{\nu_2} q_{\nu_3} + \delta_{\nu_2\nu_1} q_{\nu_3} q_\mu + \\ + \delta_{\nu_3\nu_2} q_{\nu_1} q_\mu) + q^4 (\delta_{\mu\nu_3} \delta_{\nu_1\nu_2} + \delta_{\nu_3\nu_2} \delta_{\mu\nu_1}) \}. \quad (68)$$

Let us now consider a mixture of N scalar fields with masses M_i and n spinor fields with masses m_i . From equations (50), (51) and (68) we get the divergent term in the kernel of the e^4 approximation (after a suitable symmetrization of (51))

$$\frac{1}{64} \int_{-1}^{+1} du_1 (1-u_1)^2 \int_{-1}^{+1} du_2 (1-u_2) \int_{-1}^{+1} du_3 \int_{-\infty}^{+\infty} x^3 dx \left[\left\{ 2 \sum_{i=1}^n e^{ixm_i^2} - \sum_{i=1}^N e^{ixM_i^2} \right\} \times \right. \\ \times \int dq e^{ixq^2} \{ 16 q_\mu q_{\nu_1} q_{\nu_2} q_{\nu_3} - 4 q^2 (\delta_{\mu\nu_3} q_{\nu_1} q_{\nu_2} + \delta_{\nu_1\mu} q_{\nu_2} q_{\nu_3} + \\ + \delta_{\nu_2\nu_1} q_\mu q_{\nu_3} + \delta_{\nu_3\nu_2} q_{\nu_1} q_\mu) + q^4 (\delta_{\mu\nu_3} \delta_{\nu_1\nu_2} + \delta_{\nu_3\nu_2} \delta_{\mu\nu_1}) \} + \\ + (e^{ix\varphi} - 1) \times \left[\sum_{i=1}^n e^{ixm_i^2} Sp[\gamma_\mu \gamma_q \gamma_{\nu_1} \gamma_q \gamma_{\nu_2} \gamma_q \gamma_{\nu_3} \gamma_q] - \right. \\ \left. - \sum_{i=1}^N e^{ixM_i^2} (\pi^{(3)}(qqqq) - q^2 \sum_i \delta_{\nu_i \nu_{i-1}} \times \right. \\ \left. \left. \times \pi_{i, i-1}^{(3)}(qq) + \frac{1}{2} \sum_{i \neq j} \delta_{\nu_i \nu_{i-1}} \delta_{\nu_j \nu_{j-1}}) \right] \right] \quad (69)$$

The last term in (69) is convergent as $e^{i\varphi x} - 1$ vanishes linear at the origin. The first term can be made convergent too by the aid of the assumption that

$$\lim_{x \rightarrow 0} \frac{1}{x} \left[2 \sum_{i=1}^n e^{ixM_i^2} - \sum_{i=1}^N e^{ixM_i^2} \right] \quad (70)$$

is finite. This is the case if

$$2n = N \quad (71)$$

which is one of the conditions of JOST and RAYSKI. Their other condition

$$2 \sum_{i=1}^n m_i^2 = \sum_{i=1}^N M_i^2 \quad (72)$$

makes (70) vanish, but this assumption is not needed to make (69) converge.

The symmetrisation of (51) that is necessary to get (69) is certainly allowed. In fact, the expressions given by (7) can always be made symmetric in the variables and the whole calculation can be carried through in this way. We have purposely destroyed this symmetry (equations (22) and (55)) to get formulae, which are more easy to handle.

All the integrals appearing in the kernel of the e^4 approximation are now convergent and hence our formal proof of the gauge-invariance may be applied in this case too. The only remaining divergent (but actually gauge invariant) expression in the phenomena is the charge renormalisation term in the e^2 approximation.

I want to express my respectful gratitude to professor W. PAULI, Zürich, who has suggested this investigation to me, and to thank him and Dr. R. JOST and Dr. J. LUTTINGER for many helpful discussions. I am further indebted to professor T. GUSTAFSON, LUND, who has arranged my stay at the Swiss Federal Institute of Technology, and to the Swedish Atomic Committee for financial support.

Appendix.

Proof of equation (23).

From the formulae (17), (20) and (21) the symbols $E_k^{(n)}$ in (22) may be computed. The result is (for n an odd number)

$$\begin{aligned} E_{(n+1-k)}^{(n)} = & (-1)^k [\varepsilon(0n) \varepsilon(n, n-1) \dots \varepsilon(n-k, 1) \varepsilon(12) \varepsilon(23) \dots \\ & \dots \varepsilon(n-k-2, n-k-1) + \varepsilon(0n) \varepsilon(n, n-1) \dots \\ & \dots \varepsilon(n-k+1, 1) \varepsilon(1, n-k) \varepsilon(n-k, 2) \varepsilon(23) \dots \\ & \dots \varepsilon(n-k-2, n-k-1) + \dots] \end{aligned} \quad (73)$$

Denoting

$$(-1)^k [E_{(n+1-k)}^{(n)} + \varepsilon(01) \varepsilon(12) \dots \varepsilon(n-k-1, n-k) \varepsilon(n-k+1, n-k+2) \dots \dots \varepsilon(n-1, n) \varepsilon(n0)]$$

with

$$S^{(n)(k)}(01 \dots n-k-1; n-k \dots n)$$

we get from formula (73) the following recursion formulae

$$\begin{aligned} (k \neq 0) S^{(n)(k)}(01 \dots n-k-1; n-k \dots n) = & \varepsilon(0n) S^{(n-1)(k-1)}(n12 \dots \\ & \dots n-k-1; n-k \dots n-1) + \varepsilon(01) \varepsilon(1n) S^{(n-2)(k-1)}(n23 \dots \\ & \dots n-k-1; n-k \dots n-1) + \dots + \varepsilon(01) \varepsilon(12) \dots \\ & \dots \varepsilon(n-k-2, n) S^{(k+1)(k-1)}(n \dots n-k-1; n-k \dots n-1) + \\ & + (-1)^k \varepsilon(n-k, n-k+1) \dots \varepsilon(n-1, n) \times \\ & \times S^{(n-k)(0)}(012 \dots n-k-1; n) \end{aligned} \quad (74)$$

and

$$\begin{aligned} S^{(n)(0)}(01 \dots n-1, n) = & \varepsilon(01) S^{(n-1)(0)}(12 \dots n-1; n) + \varepsilon(12) \dots \\ & \dots \varepsilon(n-2, n-1) S^{(n-k)(0)}(012 \dots n-k-1; n) \end{aligned} \quad (75)$$

From the identity

$$S^{(2)(0)}(01; 2) = \varepsilon(02) \varepsilon(21) + \varepsilon(01) \varepsilon(12) + \varepsilon(01) \varepsilon(20) = -1 \quad (76)$$

and equation (75) it is immediately seen, that $S^{(n)(0)}$ can be expressed as a sum of terms, each of which do not contain more than $n-2$ factors ε . From (74) the same statement is seen to be true also for the general expression $S^{(n)(k)}$. Using the property that no real pairs can be created, we now get (23) from (22).

Proof of equation (63).

The symbol $\overline{P}_{\mu\nu_1 \dots \nu_n}^{(n)}$ of equation (62) can be computed from equations (54)–(60). The result is (for n odd)

$$\begin{aligned}
 \overline{P}_{\mu\nu_1 \dots \nu_n}^{(n)} = & \pi^{(n)} - \sum_i \delta_4 \nu_i \delta_4 \nu_{i-1} (p^{i-1^2} + m^2) \left[\pi_{i, i-1}^{(n)} - \right. \\
 & - \frac{1}{2} \sum_j \delta_4 \nu_j \delta_4 \nu_{j-1} (p^{j-1^2} + m^2) \left[\pi_{i, i-1, j, j-1}^{(n)} - \frac{1}{3} \sum [\dots] \right] + \\
 & + \sum_i \xi(i, i-1) (p^{i-1^2} + m^2) \left[\pi_{i, i-1}^{(n)} - \sum_j \delta_4 \nu_j \delta_4 \nu_{j-1} (p^{j-1^2} + m^2) \times \right. \\
 & \times \left. \left[\pi_{i, i-1, j, j-1}^{(n)} - \frac{1}{2} \sum [\dots] \right] \right] + \frac{1}{2!} \sum_{i, j} \xi(i, i-1) \xi(j, j-1) \times \\
 & \times (p^{i-1^2} + m^2) (p^{j-1^2} + m^2) \left[\pi_{i, i-1, j, j-1}^{(n)} - \sum [\dots] \right] + \\
 & + \frac{1}{3!} \sum [\dots] + \dots
 \end{aligned} \tag{77}$$

We here use the definition (3c) of $\xi(ij)$ and write down the coefficient for a term consisting of $2l$ factors $\delta_4 \nu_i \delta_4 \nu_{i-1}$ and λ factors $\delta_{\nu_i \nu_{i-1}}$. This coefficient is

$$\frac{(-1)^\lambda}{\lambda!} \sum_{s=0}^l \frac{(-1)^s}{s! (l-s)!} = \begin{cases} \frac{(-1)^\lambda}{\lambda!} \frac{(1-1)^l}{l!} = 0 & l \neq 0 \\ \frac{(-1)^\lambda}{\lambda!} & l = 0 \end{cases} \tag{78}$$

The only non-vanishing terms in (77) thus consist only of factors $\delta_{\nu_i \nu_{i-1}}$ with coefficients given by (78). This is formula (63).