

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 23 (1950)  
**Heft:** [3]: Supplementum 3. Internationaler Kongress über Kernphysik und Quantenelektrodynamik  
  
**Artikel:** Magnetic analysis of disintegration products  
**Autor:** Buechner, W.W.  
**DOI:** <https://doi.org/10.5169/seals-422242>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

### III.

## Experimentelle Ergebnisse der Kernphysik



## Magnetic analysis of Disintegration products

by **W. W. Buechner** (M. I. T. Cambridge, Mass.).

An annular magnet, in which use is made of 180-degree focusing, has been employed to study the charged particles from several nuclear reactions. The region of uniform field, in which the particles are deflected, is approximately 70 centimeters in mean diameter and 5 centimeters wide with a spacing of 1.5 centimeters between the pole faces. The magnetic field strength is measured with a sensitive null-type flux meter calibrated in terms of deflection of polonium alpha-particles. The energy of the incident particles is also determined in terms of this standard by measuring their deflection after they have been elastically scattered from various thin targets. Nuclear track plates have been used to detect the analysed particle groups. As the apparatus is normally employed, each plate is exposed at a different magnetic field strength, the incident bombarding voltage being held constant. Thus, each plate covers a certain interval in the energy spectrum of the particles resulting from the reaction, the width of the interval for a particular plate depending upon the field strength at which it was exposed. The high resolution of apparatus when used in conjunction with an electrostatic accelerator makes it particularly suitable for measuring reaction energies and for searching for particle groups that provide information regarding nuclear energy levels<sup>1</sup>).

Thus far, attention has been given mainly to proton groups from reactions produced by deuterons having energies in the range of 1.5 MeV. The energies of the first energy levels in  $\text{Li}^7$ ,  $\text{Be}^{10}$ ,  $\text{C}^{13}$ , and  $\text{O}^{17}$ , expected when these nuclei are produced in such reactions, have been found to be  $0.483 \pm 0.003$ ;  $3.375 \pm 0.015$ ;  $3.098 \pm 0.008$ ; and  $0.876 \pm 0.009$  MeV, respectively. A particular search has been made for particle groups that would indicate additional lower energy levels in  $\text{Be}^{10}$  and  $\text{C}^{13}$  to see whether any exist that might be correlated with known levels in the mirror nuclei  $\text{B}^{10}$  and  $\text{N}^{13}$ . While such correspondence would be expected for equal neutron-neutron and proton-proton forces, no such evidence for additional levels has been found. This lack of correlation between the energy levels of mirror nuclei appears also to be the case for  $\text{Li}^7$  and  $\text{Be}^7$ . Studies of the

various reactions in which  $\text{Be}^7$  is produced<sup>2)3)4)</sup> show no indications of an excited state in  $\text{Be}^7$  that would correspond to the well-known level in  $\text{Li}^7$  at 480 MeV<sup>5)</sup>.

The energy levels in  $\text{N}^{13}$  have been studied by the  $\text{C}^{12}(p, \gamma) \text{N}^{13}$  reaction. A resonance has been found for both gamma-ray emission and for positron activity at a proton energy of  $1.697 \pm 0.012$  MeV, the half-width of the resonance being  $74 \pm 9$  KeV. This indicates that, in addition to the one at  $2.383 \pm 0.018$  MeV<sup>6)</sup>, there is an energy level in  $\text{N}^{13}$  at  $3.523 \pm 0.019$  MeV. These measurements show that, in the range of proton energies from 0.6 to 2.1 MeV, there are no additional resonances with a peak intensity so large as 0.12 that of the one at 1.70 MeV. The existence of these levels has been confirmed by the recent work of GROSSKREUTZ on the energy of the neutron groups from the  $\text{C}^{12}(d, n) \text{N}^{13}$  reaction<sup>7)</sup>.

The energies evolved in a number of reactions have been measured. The following values are for the reactions that lead to the formation of the residual nuclei in their ground states, the values being in MeV:

$\text{Li}^6(d, p)\text{Li}^7$ :	$5.006 \pm 0.014$	$\text{Li}^7(d, p)\text{Li}^8$ :	$-0.193 \pm 0.008$
$\text{Be}^9(d, \alpha)\text{Li}^7$ :	$7.145 \pm 0.024$	$\text{Be}^9(d, p)\text{Be}^{10}$ :	$4.576 \pm 0.012$
$\text{C}^{12}(d, p)\text{C}^{13}$ :	$2.729 \pm 0.009$	$\text{O}^{16}(d, p)\text{O}^{17}$ :	$1.925 \pm 0.008$

This work has been assisted by the joint program of the Office of Naval Research and the Atomic Energy Commission.

#### *Bibliography.*

<sup>1)</sup> The apparatus is described in more detail in an article to appear in the December 1, 1949 issue of The Physical Review.

<sup>2)</sup> FREIER, LAMPI and WILLIAMS, Phys. Rev. **75**, 901 (1949).

<sup>3)</sup> BURCHAM and FREEMAN, Nature **163**, 167 (1949).

<sup>4)</sup> MANDEVILLE, SWANN and SNOWDEN, Phys. Rev. **76**, 980 (1949).

<sup>5)</sup> Note added in proof: Recent work has indicated the existence of excited states in  $\text{Be}^7$  at 0.205, 0.470, and 0.745 MeV. (W. A. FOWLER, private communication; GROSSKREUTZ and MATHER, Bul. Amer. Phys. Soc., **24**, November 25, 1949.)

<sup>6)</sup> FOWLER and LAURITSEN, Phys. Rev. **76**, 314 (1949).

<sup>7)</sup> J. C. GROSSKREUTZ, Phys. Rev. **76**, 482 (1949).