

Zeitschrift: Helvetica Physica Acta
Band: 30 (1957)
Heft: I

Artikel: High resolution proton magnetic resonance in water and the concept of hydrogen bonding
Autor: Ogg, Richard A.
DOI: <https://doi.org/10.5169/seals-112806>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

High Resolution Proton Magnetic Resonance in Water and the Concept of Hydrogen Bonding

by Richard A. Ogg

Physikalische Anstalt der Universität Basel.

(14. III. 1957)

The proton magnetic resonance spectrum of liquid water is commonly employed as a reference standard for high resolution proton resonance spectra in other compounds. The development of a thermally insulated "head" containing the transmitter and receiver coils has made possible the study of proton resonance in gaseous water. A liquid water sample of quantity corresponding to about ten atmospheres pressure when vaporized was contained in the conventional slender Pyrex glass tube, previously evacuated and sealed off. The sample tube was heated in an electric furnace to cause complete volatilization of the liquid and was then rapidly transferred to the Dewar vessel insert of a thirty megacycle nuclear magnetic resonance spectrometer. The signal to noise ratio was adequate to demonstrate an appreciable line breadth (of the order of five cycles per second) for the proton resonance. Most interesting was the displacement relative to liquid water. The average of fifteen measurements yielded a figure of 150 cycles per second, the proton resonance in the vapor state falling at the higher magnetic field.

The author had previously demonstrated*) a similar effect in the case of gaseous and liquid ammonia, the proton resonance in the gas falling at higher magnetic field, with a displacement of some 50 cycles per second at a frequency of 30 megacycles. The explanation offered for the present results is precisely that previously proposed, namely that so-called "Hydrogen bonding" is merely close classical electrostatic interaction between molecular electric dipoles. The considerably greater effect observed for water is due to the fact that the permanent electric dipole moment of water molecules is considerably greater than that of ammonia.

*) R. A. Ogg, Jr., J. Chem. Physics, **22**, 560 (1954).

It should be noticed that the 150 cycles per second shift observed for a simple phase transition is greater than the usual magnitudes of "chemical shift" associated with differences in chemical structure. For purpose of relating nuclear magnetic resonance spectra to molecular structure, the vapor state is ideal. In the accompanying letter by Mr. P. DIEHL, it is seen, that the widely held concepts about the "classical" case of ethyl alcohol are due to the previous restriction of the experimental studies of the proton magnetic resonance spectrum to the pure liquid. The above results with water would suggest that (in comparison with the vapor) the spectrum is strongly distorted by electrostatic interaction in the pure liquid. This appears to be in fact the case. The author wishes to acknowledge the assistance of Dr. J. SHOOLERY of Varian Associates, Palo Alto, California, in the experimental work. Financial support of the Research Corporation and of the National Science Foundation is gratefully acknowledged.