

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 31 (1958)  
**Heft:** VII

**Artikel:** Theory of the scattering operator. II, Multichannel scattering  
**Autor:** Jauch, J.M.  
**DOI:** <https://doi.org/10.5169/seals-112928>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Theory of the Scattering Operator II Multichannel Scattering\*)

by **J. M. Jauch\*\*)**

CERN, Theoretical Study Division, Geneva

(10. VIII. 58)

---

*Abstract.* The mathematical theory of the scattering operator is developed for the general scattering systems involving an arbitrary number of channels. It includes as a special case the theory for 'simple scattering systems' given in an earlier paper. The scattering system is defined as a quantum mechanical system which satisfies certain asymptotic and completeness conditions given in Section 4. The existence of the  $S$ -operator as well as its unitary property is then a rigorous mathematical consequence of this property. A crucial step in these deductions is the orthogonality theorem for the left projections of the wave operators which is proved in Section 5. In the last Section 7, we discuss the various ways of introducing the 'in' and 'out'-operators and their relation to the  $S$ -operator.

## 1. Introduction

In a recent paper<sup>1)</sup> the author has developed the theory of scattering on a mathematically rigorous foundation for the so-called 'simple scattering systems'. This paper is an extension of this work to the case of multichannel scattering. The basic philosophy adopted in this paper is much the same as in the previous one. Only mathematically well defined concepts, symbols and operations are employed. All steps are mathematically rigorous. The space of the state vectors is the classical Hilbert space and, in view of recent discussions we may add, this implies a positive definite metric.

The main result of this paper is the precise formulation of the concept of a multichannel scattering system and the proof of the existence of the scattering operator as well as its unitary property in a certain subspace of the Hilbert space. As was already pointed out by EKSTEIN<sup>2),3)</sup> the operator defined as  $S$  in our earlier paper, does not have a generalization in the multichannel case. The suitable operator for this purpose is another one defined and discussed in Section 2.

---

\*) Supported in part by the National Science Foundation.

\*\*) On leave of absence from the State University of Iowa, Iowa City, Iowa, U. S. A.

The results obtained are equally valid whether we are dealing with the relativistic or non-relativistic form of the theory. Furthermore, the formalism is developed in such a manner that the cases with bound states are also included. The number of channels need not be finite as it is the case for instance in nuclear reactions.

The formulation is sufficiently broad so as to include all known types of scattering and reaction processes. In particular, we feel confident that the formulation presented here will serve as a sufficiently precise mathematical framework within which it should be possible to accommodate the theories dealing with fundamental particles and their interactions.

## 2. The operators $S$ and $S'$ for simple scattering systems

A 'simple scattering system' was previously defined<sup>1)</sup> as a quantum mechanical system with the following properties:

For every  $f \in \mathfrak{H}$  the strong limits

$$\lim_{t \rightarrow \mp \infty} V_t^* U_t f = f_{\pm} = \Omega_{\pm} f \quad (2.1)$$

exist and the range of the operators  $\Omega_{\pm}$  is equal to the subspace  $N$  of the continuum states of  $V_t$ .

In these expressions  $V_t = e^{-iHt}$  ( $-\infty < t < +\infty$ ) is the transformation group of the system and  $H$  is the total energy operator. The group  $U_t = e^{-iH_0 t}$  represents the free motion of the system in the absence of the interaction.  $H_0$  is the kinetic energy operator for the particles participating in the scattering process.

The conditions (2.1) have a direct interpretation in terms of the actual scattering process and they entail the existence of a unitary scattering operator

$$S = \Omega_-^* \Omega_+ \quad (2.2)$$

The 'wave operators'  $\Omega_{\pm}$  are isometries in all of  $\mathfrak{H}$  and they satisfy for both signs

$$\left. \begin{aligned} \Omega^* \Omega &= I \\ \Omega \Omega^* &= E_N \end{aligned} \right\} \quad (2.3)$$

where  $E_N$  is the projection operator into the subspace of continuum states of  $H$ .

All these statements were proved in reference<sup>1)</sup> and are here briefly repeated for convenience.

The definition (2.2) of the scattering operator is the one which is usually implied in much of the current literature on this subject. It is possible to

define a different scattering operator, which we shall temporarily denote by  $S'$ :

$$S' = \Omega_+ \Omega_-^* . \quad (2.4)$$

This operator has properties very similar to  $S$  which we shall enumerate here briefly.

The interest in this operator arises from the fact that in a multi-channel theory it has a proper generalization while the operator (2.2) cannot be so generalized.

We observe first that the operator product in (2.4) is well defined since the  $\Omega$  and  $\Omega^*$  are bounded operators and are therefore defined in the entire space  $\mathfrak{H}$ . We can further form without restriction the products

$$S'S'^* = \Omega_+ \Omega_-^* \Omega_- \Omega_+^* = \Omega_+ \Omega_+^* = E_N , \quad (2.5)$$

and similarly

$$S'^* S' = E_N . \quad (2.6)$$

Furthermore

$$E_N S' = \Omega_+ \Omega_+^* \Omega_+ \Omega_-^* = \Omega_+ \Omega_-^* = S' , \quad (2.7)$$

and similarly

$$S' E_N = S' . \quad (2.8)$$

The operator  $S'$  annihilates therefore all the elements in  $M = N^\perp$  and in the invariant subspace  $N$  it is unitary. We shall say  $S'$  is 'quasi-unitary'.

The following relations between the two operators  $S$  and  $S'$  are direct consequences of the definitions (2.2), (2.4) and the relations (2.3).

$$\Omega_+^* S' \Omega_+ = S = \Omega_-^* S' \Omega_- , \quad (2.9)$$

$$\Omega_+^* S' \Omega_- = I , \quad (2.10)$$

$$\Omega_-^* S' \Omega_+ = S^2 . \quad (2.11)$$

The operator  $S'$  commutes with  $V_t$  but not with  $U_t$ . This follows from the intertwining property of  $\Omega$  which was proved in reference<sup>1</sup>):

$$V_t \Omega = \Omega U_t . \quad (2.12)$$

For instance one obtains

$$S' V_t = \Omega_+ \Omega_-^* V_t = \Omega_+ U_t \Omega_-^* = V_t \Omega_+ \Omega_-^* = V_t S' .$$

Since  $U_t$  does not commute with  $S'$  the operator  $S'(t) = U_t^* S U_t$  depends on  $t$ . It satisfies the following relations

$$S'(-\infty) = S'(+\infty) = S . \quad (2.13)$$



These limits are understood in the strong operator topology. This means for every  $f \in \mathfrak{H}$  we have

$$\| (S'(t) - S) f \| \rightarrow 0 \quad \text{for } t \rightarrow \pm \infty. \quad (2.14)$$

One may verify this by using the relations proved in reference<sup>1</sup>):

$$\begin{aligned} \Omega_+(+\infty) &= S = \Omega_-(-\infty) \\ \Omega_+(-\infty) &= I = \Omega_-(+\infty), \end{aligned} \quad (2.15)$$

where  $\Omega(t) = U_t^* \Omega U_t$ . Thus for the limit  $t \rightarrow \infty$  we write the decomposition

$$S'(t) - S = \Omega_+(t) (\Omega_-^*(t) - I) + (\Omega_+(t) - S).$$

Using the boundedness of the  $\Omega(t)$ , we find

$$\| (S'(t) - S) f \| \leq \| (\Omega_-^*(t) - I) f \| + \| (\Omega_+(t) - S) f \|. \quad (2.16)$$

Since both terms on the right hand side tend to zero with  $t \rightarrow +\infty$  by (2.15) the left hand side does so too and one of the relations (2.13) is established. The other is verified similarly.

### 3. The channel energy

The most characteristic feature of the general scattering process is the occurrence of 'free' particles before and after the collision. Every set of free particles is characterized by a set of parameters which expresses the values of the rest masses, spin, charge and whatever other variables are needed for a full description of the free particles. Each possible set of values for these parameters determines a different 'channel' of the system.

A free particle is one which moves exactly the same as if other particles were not present. The Hamiltonian for the free motion of the particles in a channel (also called the channel Hamiltonian) is therefore merely the sum of the kinetic energy of all the particles in the channel. This Hamiltonian is in general not obtained from the total Hamiltonian by the mere omission of certain terms, as it was the case in the simple scattering systems.

We shall illustrate this on one of the simplest examples of multi-channel scattering. Let us consider three fundamental particles labelled 1, 2 and 3.

Let the total Hamiltonian be of the form

$$H = \frac{p_1^2}{2m_1} + \frac{p_2^2}{2m_2} + \frac{p_3^2}{2m_3} + V_{12} + V_{23} + V_{31}$$

where the potentials  $V_{rs}$  describe the three possible two-body interactions of these particles. Let us assume the interaction between 1 and 2 is such

that there exists one or more bound states, represented by square integrable solutions of the Schrödinger equation

$$\left( \frac{p_1^2}{2m_1} + \frac{p_2^2}{2m_2} + V_{12} \right) \Psi = E \Psi$$

The state vector  $\Psi$  describes then a fragment which can enter a scattering process or which may be produced as a final state. The total mass of the fragment is

$$M = m_1 + m_2 - |E|,$$

and the total kinetic energy is given by

$$H_0 = \begin{cases} \frac{1}{2M} \mathbf{P}^2 & \text{non relativistic} \\ \sqrt{M^2 + \mathbf{P}^2} & \text{relativistic} \end{cases}$$

where  $\mathbf{P} = (\mathbf{p}_1 + \mathbf{p}_2)$  is the momentum of the centre of mass.

It is conceivable that the interaction between another pair, say 2 and 3, is sufficiently strong so as to produce also a bound state. This will give rise to a new fragment of the mass

$$M' = m_2 + m_3 - |E'|$$

where  $E'$  is the binding energy of particle 2 and 3. The corresponding expression for the kinetic energy of this fragment is then given by

$$H_0 = \begin{cases} \frac{1}{2M'} \mathbf{P}'^2 & \text{non relativistic} \\ \sqrt{M'^2 + \mathbf{P}'^2} & \text{relativistic} \end{cases}$$

where  $\mathbf{P}' = (\mathbf{p}_2 + \mathbf{p}_3)$ .

In this description every bound state which leads to a different binding energy is in a different channel. For a degenerate bound state we obtain a fragment with additional internal degrees of freedom.

It is clear from this example how one would obtain the most general channel Hamiltonian for the fragments composed of any number of particles. We shall not elaborate this approach however. Instead we shall formulate the properties of channel Hamiltonians which we believe to be essential for the mathematical formulation of multichannel scattering theory.

These properties are purposely formulated sufficiently general so that they would presumably be satisfied for elementary particles as well as stable fragments. From the point of view of scattering theory the distinction between elementary particle and composite fragments is an entirely superficial one and this is as it should be. The physical picture of an

elementary particle breaks down as soon as internal structure reveals itself in certain collision processes. On the other hand, a stable fragment may in every respect behave like a fundamental particle as long as the collisions are sufficiently slow to leave the fragment intact. In a good scattering theory therefore this distinction should not appear as an essential element.

In agreement with this requirement we shall describe the properties of the channel Hamiltonian which are derived from the description of channels in terms of fragments but which are presumed to have general validity for any kind of particle composite or elementary. Instead of working with the channel Hamiltonian  $H$  we shall immediately express the characteristic properties in terms of the 'channel operators'

$$U_t^{(\alpha)} \equiv e^{-iH_\alpha t}.$$

Channel operators  $U_t^{(\alpha)}$  are a set of continuous unitary representations of the additive group of real numbers ( $-\infty < t < +\infty$ ) in the underlying Hilbert space of state vectors with the following three properties:

(1) All  $U_t^{(\alpha)}$  commute with one another for all  $\alpha$  and all  $t$ :

$$[U_t^{(\alpha)}, U_{t'}^{(\beta)}] = 0, \text{ all } \alpha, \beta, t, t'. \quad (3.1)$$

(2) The family of spectral projections of the infinitesimal generators of the group are continuous. There are no discrete eigenvalues.

(3) The operators  $U_t^{(\alpha)}$  are 'essentially different' for different indices. With this we mean

$$U_t^{(\alpha)} f = U_t^{(\beta)} f \text{ for all } t \text{ implies } f = 0 \text{ unless } \alpha = \beta. \quad (3.2)$$

#### 4. The definition of a scattering system

The channel operator which we have defined in the preceding section are rarely known explicitly. Instead one usually knows only the total energy operator  $H$  or its corresponding unitary group

$$V_t = e^{-iHt}. \quad (4.1)$$

It is clear that all the information as to the possible occurrence of various particles as well as their physical properties should be contained in the structure of the group (4.1). The different particles form the different channels and are described by certain channel operators  $U_t^{(\alpha)}$ . We shall now formulate the condition which is needed in order that the group (4.1) describes particles associated with certain channel operators  $U_t^{(\alpha)}$ :

A unitary group  $U_t^{(\alpha)}$  with the three properties described in the preceding section is a channel of the system defined by  $V_t$  if there exists at

least one element  $f \in \mathfrak{H}$ ,  $f \neq 0$ , for which the strong limits

$$\lim_{t \rightarrow \mp \infty} V_t^* U_t^{(\alpha)} f = f_{\pm}^{(\alpha)} \quad (4.2)$$

exist.

The condition (4.2) may be considered the defining property of a channel energy. It extracts from the total transformation operator the kind and energy of the stable fragments which can be associated with it. Since this condition is an asymptotic property the definition of particles which are obtained in this way includes already all the possible self interactions. In the language of field theory: The particles are 'dressed particles'.

On the other hand, we see that in general it is not possible of introducing one single free-particle Hamiltonian. Every channel, that is every kind of free particles, has its own energy operator associated with it.

As a first consequence of the defining property (4.2) we shall show that if there exists one single element  $f$  for which (4.2) is satisfied, then there exists an infinite-dimensional subspace of  $\mathfrak{H}$  with the same property.

Let  $D_\alpha$  be the set of elements  $f$  such that the limit (4.2) exists. We first observe that  $D_\alpha$  is a closed linear manifold, that is a subspace of  $\mathfrak{H}$ . The linear property is obvious. In order to prove the closure property we consider an arbitrary sequence  $f_n \in D_\alpha$  for which  $f_n \rightarrow f$  with  $n \rightarrow \infty$ . We must show that the limit (4.2) exists for the element  $f$ .

Let us define  $W_{t_1 t_2} \equiv V_{t_1}^* U_{t_1} - V_{t_2}^* U_{t_2}$ , then

$$\|W_{t_1 t_2} f\| \leq \|W_{t_1 t_2} f_n\| + \|W_{t_1 t_2} (f - f_n)\|.$$

The second term may further be estimated by

$$\|W_{t_1 t_2} (f - f_n)\| \leq \|W_{t_1 t_2}\| \|f - f_n\| \leq 2 \|f - f_n\|.$$

The last step is a consequence of the triangle inequality

$$\|A + B\| \leq \|A\| + \|B\|$$

for the bounds of bounded operators.

We have therefore in all

$$\|W_{t_1 t_2} f\| \leq \|W_{t_1 t_2} f_n\| + 2 \|f - f_n\|.$$

We choose first a fixed  $n$  such that, independently of  $t_1$  and  $t_2$  the last term is  $\leq \epsilon/2$  for an arbitrary  $\epsilon > 0$ . We then determine a  $T$ , such that for  $t_1 > T$  and  $t_2 > T$  the first term is  $\leq \epsilon/2$  too. This is possible because  $f_n \in D_\alpha$ . For such values of  $t_1$  and  $t_2$  we have then

$$\|W_{t_1 t_2} f\| \leq \epsilon \quad \text{all } t_1 > T, t_2 > T.$$

The elements of the form  $V_t^* U_t f$  satisfy therefore the Cauchy criterion and because of the completeness of  $\mathfrak{H}$   $\lim_{t \rightarrow +\infty} V_t^* U_t f$  exists. The limit  $t \rightarrow -\infty$  is established in the same manner. Thus  $D_\alpha$  is closed, q.e.d.

We now show that the subspace  $D_\alpha$  is invariant under the group  $U_t^{(\alpha)}$ . This means if  $f \in D_\alpha$  then  $U_\tau^{(\alpha)} f \in D_\alpha$ . Indeed

$$\lim_{t \rightarrow \pm \infty} V_t^* U_t^{(\alpha)} U_\tau^{(\alpha)} f = \lim_{t \rightarrow \pm \infty} V_\tau V_t^* U_t^{(\alpha)} f = V_\tau f_\mp, \quad (4.3)$$

and  $D_\alpha$  is seen to be invariant under  $U_t^{(\alpha)}$ .

We can now easily complete the proof that  $D_\alpha$  is infinite dimensional. Assume to the contrary that  $D_\alpha$  is finite dimensional. Then the reduction of  $U_t^{(\alpha)}$  to the invariant subspace  $D_\alpha$  furnishes a finite dimensional unitary representation of the group of real numbers. Such a representation has only discrete eigenvalues, this contradicts the basic property (2) of the previous section. Thus  $D_\alpha$  is infinite dimensional, q. e. d.

It was shown previously<sup>1)</sup> that the mapping  $f \rightarrow f_\pm^{(\alpha)}$  is a linear isometry. The ranges  $R_\pm^{(\alpha)}$  of this correspondence are therefore closed linear manifolds, that is subspaces.

The set of subspaces  $R_\pm^{(\alpha)}$  as  $\alpha$  runs through all the channels of the system span a linear manifold which we denote by  $\{R_\pm^{(\alpha)}\}$ . Its closure shall be denoted by

$$R_\pm \equiv \overline{\{R_\pm^{(\alpha)}\}}. \quad (4.4)$$

We shall further write  $N$  for the subspace of continuum states of  $V_t$ , that is, the orthogonal complement of the subspace  $M$  of proper elements of  $V_t$ .

With these preliminaries out of the way we define the general scattering system as follows:

*A quantum mechanical system, described by the unitary group  $V_t$ , is a scattering system if there exists a set of channels  $\alpha$ , together with their unitary groups  $U_t^{(\alpha)}$ , such that*

$$R_+ = N = R_-. \quad (4.5)$$

The physical meaning of the requirement (4.5) is simply this: Every continuum state must be a superposition of scattering states, that is, states which are in the ranges  $R_\pm^{(\alpha)}$ .

It is readily seen that this definition of the scattering system is a generalization of the definition<sup>1)</sup> for a 'simple scattering system' to which it reduces if there exists only one channel.

The number of channels may be finite or infinite. We shall see in the next section, however, that in case it is infinite it is necessarily countably infinite.

This definition of a scattering system embodies the minimum requirements which must be placed on a scattering system. Yet there are some far reaching conclusions which can be drawn from it as we shall see in the following sections.

### 5. The wave operators

We shall now assume that  $V_t$  describes a general scattering system as defined in the preceding section. There exists then a set of commuting unitary groups  $U_t^{(\alpha)}$  such that for all  $f \in D_\alpha$

$$\lim_{t \rightarrow \mp \infty} V_t^* U_t^{(\alpha)} f = f_{\pm}^{(\alpha)}. \quad (5.1)$$

Let  $R_{\pm}^{(\alpha)}$  be the set of elements  $f_{\pm}^{(\alpha)}$ . According to the previous paper<sup>1)</sup> the mapping of  $D_\alpha$  onto  $R_{\pm}^{(\alpha)}$  is a linear isometry, and  $R_{\pm}^{(\alpha)}$  is a subspace (= closed linear manifold). This mapping defines therefore a bounded linear operator  $\Omega_{\pm}^{(\alpha)}$  on  $D_\alpha$  by the condition

$$\Omega_{\pm}^{(\alpha)} f = f_{\pm}^{(\alpha)} \quad f \in D_\alpha. \quad (5.2)$$

Such an operator can always be extended in a continuous manner to the whole space by, for instance, the following procedure: Let  $f \in \mathfrak{H}$  and  $f = g + h$ ,  $g \in D_\alpha$ ,  $h \in D_\alpha^\perp$ . Define

$$\Omega_{\pm}^{(\alpha)} f = \Omega_{\pm}^{(\alpha)} g. \quad (5.3)$$

We shall denote this extended operator with the same symbol.

To every bounded operator  $\Omega_{\pm}^{(\alpha)}$  can be associated uniquely<sup>4)</sup> an adjoint operator  $\Omega_{\pm}^{(\alpha)*}$  by the condition

$$(\Omega_{\pm}^{(\alpha)*} f, g) = (f, \Omega_{\pm}^{(\alpha)} g) \quad (5.4)$$

for all  $f, g \in \mathfrak{H}$ . It follows from this definition that  $\Omega_{\pm}^{(\alpha)*}$  vanishes on the orthogonal complement  $R_{\pm}^{(\alpha)\perp}$  of the range of  $\Omega_{\pm}^{(\alpha)}$ .

In order to survey some of the general properties of the wave operators we shall introduce a few definitions and deduce from them some elementary consequences.

#### Definition 1:

A bounded linear operator  $\Omega$  is a partial isometry if  $E \equiv \Omega^* \Omega$  is a projection.

It follows that if  $\Omega$  is a partial isometry then  $F \equiv \Omega \Omega^*$  is also a projection.

In order to see this we note that

$$(\Omega(E - I))^* \Omega(E - I) = (E - I) \Omega^* \Omega (E - I) = (E - I) E (E - I) = 0.$$

Therefore for any  $f \in \mathfrak{H}$

$$\| \Omega(E - I) f \|^2 = (f, (\Omega(E - I))^* \Omega(E - I) f) = 0,$$

that is  $\Omega(E - I) f = 0$  for any  $f \in \mathfrak{H}$ ,

$$\text{or} \quad \Omega(E - I) = 0, \quad (5.5)$$

$$\text{or also} \quad \Omega E = \Omega. \quad (5.6)$$

From this we obtain

$$F^2 = \Omega \Omega^* \Omega \Omega^* = \Omega E \Omega^* = \Omega \Omega^* = F$$

$$\text{and} \quad F^* = F.$$

The last two relations express that  $F$  is a projection.

*Definition 2:*

If  $\Omega$  is a partial isometry then

$E = \Omega^* \Omega$  is called the right-projection and

$F = \Omega \Omega^*$  the left-projection of  $\Omega$ .

It follows obviously that conversely  $E$  is the left-projection and  $F$  the right-projection of  $\Omega^*$ .

Left- and right-projections have a maximal property as follows:

The projection  $E$  is the smallest projection with the property

$$\Omega E = \Omega$$

and  $F$  is the smallest projection with the property

$$F \Omega = \Omega.$$

We recall the partial ordering of projections: The projection  $E$  is smaller than  $E_1$ , if for every  $f \in \mathfrak{H}$

$$\| E f \| \leq \| E_1 f \|.$$

This will be written as  $E \leq E_1$ . An equivalent formulation of this relation is  $E_1 E = E E_1 = E$ .

In order to verify the maximal property of the projection  $E$  let us assume that  $E_1$  is another projection such that

$$\Omega E_1 = \Omega.$$



We have then

$$\Omega^* \Omega E_1 = \Omega^* \Omega = E$$

or

$$E E_1 = E$$

Since both  $E$  and  $E_1$  are projections we obtain also by taking the adjoint

$$E_1 E = E$$

which means  $E \leq E_1$ . The second half of the assertion is proved in a similar manner.

The subspaces which are the ranges of the projections  $E$  and  $F$  are respectively the ranges of  $\Omega^*$  and of  $\Omega$ . For instance if  $f$  is in the range of  $F$  then  $Ff = f = \Omega \Omega^* f$ , hence it is in the range of  $\Omega$ . Conversely if  $f$  is in the range of  $\Omega$  there exists a  $g$  such that  $f = \Omega g = F \Omega g$ . Therefore  $f$  is also in the range of  $F$ . The two ranges are identical. In a similar way one verifies the second half of the statement.

We shall now apply these concepts to the operators  $\Omega_{\pm}^{(\alpha)}$ . From the earlier paper<sup>1)</sup> and the extension (5.3) it follows that they are partial isometries. We define

$$E_{\alpha} = \Omega_{\pm}^{(\alpha)*} \Omega_{\pm}^{(\alpha)} \quad (5.7)$$

$$F_{\pm}^{(\alpha)} = \Omega_{\pm}^{(\alpha)} \Omega_{\pm}^{(\alpha)*}. \quad (5.8)$$

For the projections  $E_{\alpha}$  the distinction between the two cases  $\pm$  is unnecessary since the two projections are identical.

We shall now prove the main result of this section:

*Theorem:*

The projections  $F_{\pm}^{(\alpha)}$  are orthogonal for different channels, in the sense

$$F_{+}^{(\alpha)} F_{+}^{(\beta)} = F_{-}^{(\alpha)} F_{-}^{(\beta)} = 0 \quad \text{for} \quad \alpha \neq \beta. \quad (5.9)$$

*Proof:*

We shall prove the case with the minus sign and omit the sign index.

A fully equivalent statement of the theorem is the following: Let  $f^{(\alpha)}$  and  $g^{(\beta)}$  be any two elements of  $\mathfrak{H}$  with the property

$$F^{(\alpha)} f^{(\alpha)} = f^{(\alpha)}, \quad F^{(\beta)} g^{(\beta)} = g^{(\beta)} \quad (5.10)$$

then

$$(f^{(\alpha)}, g^{(\beta)}) = 0 \quad \text{for} \quad \alpha \neq \beta. \quad (5.11)$$

We shall prove the theorem in this form.

According to the preceding remarks the elements  $f^{(\alpha)}$  and  $g^{(\beta)}$  are in the ranges of the operators  $\Omega^{(\alpha)}$  and  $\Omega^{(\beta)}$  respectively. This means there



exists element  $f \in D_\alpha$  and  $g \in D_\beta$  such that

$$f_t^{(\alpha)} \equiv V_t^* U_t^{(\alpha)} f \rightarrow f^{(\alpha)} \quad \text{for } t \rightarrow \infty \quad (5.12)$$

$$g_t^{(\beta)} \equiv V_t^* U_t^{(\beta)} f \rightarrow g^{(\beta)} \quad \text{for } t \rightarrow \infty. \quad (5.13)$$

We shall first show that for any pair of elements  $f$  and  $g$  the operators

$$W_t \equiv U_t^{(\alpha)*} U_t^{(\beta)} \quad (5.14)$$

which are also a unitary group, converge weakly to zero in the limit  $t \rightarrow \infty$ . This means

$$(f, W_t g) \rightarrow 0, \quad \text{for } t \rightarrow \infty; f \in D_\alpha, g \in D_\beta. \quad (5.15)$$

We shall first show that the limit exists and then that it is zero.

That it exists may be seen from

$$(f, W_t g) = (f_t^{(\alpha)}, g_t^{(\beta)}). \quad (5.16)$$

Both  $f_t^{(\alpha)}$  and  $g_t^{(\beta)}$  strongly converge to  $f^{(\alpha)}$  and  $g^{(\beta)}$  respectively and their scalar product converges to  $(f^{(\alpha)}, g^{(\beta)})$ :

$$\begin{aligned} & | (f_t^{(\alpha)}, g_t^{(\beta)}) - (f^{(\alpha)}, g^{(\beta)}) | \\ & \leq | ((f_t^{(\alpha)} - f^{(\alpha)}), g_t^{(\beta)}) | + | (f^{(\alpha)}, (g_t^{(\beta)} - g^{(\beta)})) | \\ & \leq \| f_t^{(\alpha)} - f^{(\alpha)} \| \| g^{(\beta)} \| + \| f^{(\alpha)} \| \| g_t^{(\beta)} - g^{(\beta)} \|. \end{aligned}$$

We have made use of Schwartz's inequality and  $\| g_t^{(\beta)} \| = \| g^{(\beta)} \|$ . Since each of the  $t$ -dependent factors goes to zero (Equation (5.12) and (5.13)) the assertion that the limit  $(f, W_t g)$  exists for  $t \rightarrow \infty$  follows. Moreover, it is shown that this limit is equal to  $(f^{(\alpha)}, g^{(\beta)})$ .

We shall next show that this limit is zero. This is easily established by using a corollary of the mean ergodic theorem of v. NEUMANN<sup>5</sup>). According to this

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T (f, W_t g) dt = (f, P g) \quad (5.17)$$

where  $P$  is the projection operator into the subspace of proper elements of  $W_t$  with eigenvalue  $+1$ . On the other hand, since the integrand has the limit  $(f^{(\alpha)}, g^{(\beta)})$  for  $t \rightarrow \infty$  we also have

$$(f, P g) = (f^{(\alpha)}, g^{(\beta)}). \quad (5.18)$$

It remains to show that  $P$  is the zero operator.

Suppose  $f$  is in the range of  $P$  then by definition of  $P$

$$W_t f = f \quad \text{for all } t$$

or

$$U_t^{(\alpha)} f = U_t^{(\beta)} f \quad (\alpha \neq \beta). \quad (5.19)$$

According to property (3) of the channel operators (Section 3) this implies  $f = 0$ . Consequently  $P = 0$  and therefore by (5.18)  $(f^{(\alpha)}, g^{(\beta)}) = 0$ . Since  $f^{(\alpha)}$  and  $g^{(\beta)}$  were arbitrary elements in the ranges  $F^{(\alpha)}$  and  $F^{(\beta)}$  we have

$$F^{(\alpha)} F^{(\beta)} = 0 \quad \text{for } \alpha \neq \beta. \text{ q.e.d.}$$

We shall now mention a few useful consequences of this theorem.

First we observe that the number of channels is either finite or countably infinite. Indeed, since  $\mathfrak{H}$  is separable and the projections  $F^{(\alpha)}$  orthogonal, their number is at most countably infinite and it is equal to the number of channels.

Furthermore since by the definition of the scattering system the closed linear manifold spanned by the ranges  $R_{\pm}^{(\alpha)}$  is the whole of the space of continuum states  $N$ , we have

$$E_N = \sum_{\alpha} F_{+}^{(\alpha)} = \sum_{\beta} F_{-}^{(\beta)}, \quad (5.20)$$

where  $E_N$  is the projection operator with range  $N$ .

As a further consequence of the orthogonality relations (5.9) we have

$$\Omega_{+}^{(\alpha)*} \Omega_{+}^{(\beta)} = E_{\alpha} \delta_{\alpha\beta} = \Omega_{-}^{(\alpha)*} \Omega_{-}^{(\beta)}. \quad (5.21)$$

For instance the first of these equations is obtained from

$$\Omega_{+}^{(\alpha)*} \Omega_{+}^{(\beta)} = \Omega_{+}^{(\alpha)*} F_{+}^{(\alpha)} F_{+}^{(\beta)} \Omega_{+}^{(\beta)} = \Omega_{+}^{(\alpha)*} F_{+}^{(\alpha)} \Omega_{+}^{(\beta)} \delta_{\alpha\beta} = \Omega_{+}^{(\alpha)*} \Omega_{+}^{(\beta)} \delta_{\alpha\beta} = E_{\alpha} \delta_{\alpha\beta}.$$

Finally we note that the operators  $\Omega^{(\alpha)}$  have the intertwining property

$$V_t \Omega^{(\alpha)} = \Omega^{(\alpha)} U_t^{(\alpha)} \quad (5.22)$$

from which follows, among other things, that the projections  $F_{\pm}^{(\alpha)}$  commute with  $V_t$

$$[F_{\pm}^{(\alpha)}, V_t] = 0 \quad \text{all } t, \text{ all } \alpha. \quad (5.23)$$

There is no orthogonality theorem for the domains  $D_{\alpha}$ , or their projections  $E_{\alpha}$ . In fact it is quite possible that one of the domains  $D_{\alpha}$  may be the entire space  $\mathfrak{H}$ . In any case the different  $E_{\alpha}$  are in general not orthogonal.

\*

## 6. The scattering operator

The operator to be defined is the generalization of the operator  $S'$  of Section 2. In order to avoid a too cumbersome notation, we shall omit the prime.

We define a sequence of operators  $S_n$  by

$$S_n = \sum_{\alpha=1}^n \Omega_+^{(\alpha)} \Omega_-^{(\alpha)*} \quad (6.1)$$

and investigate the limiting properties of these operators as  $n$  increases indefinitely. When the number of channels is finite then the sequence  $S_n$  is finite and stops for some number  $n_0$ . In this case there is no limit problem. We define

$$S = S_{n_0}. \quad (6.2)$$

In case the number of channels is infinite  $S_n$  converges strongly with  $n \rightarrow \infty$  to a limit  $S$  on all of  $\mathfrak{H}$ .

To show this consider an arbitrary element  $f \in \mathfrak{H}$  and the sequence

$$f_n = S_n f. \quad (6.3)$$

We must show  $f_n$  converges to some element  $g$ . Because  $\mathfrak{H}$  is complete it suffices to show that  $f_n$  is a Cauchy sequence.

Assuming  $n \geq m$ , we have

$$\|f_n - f_m\|^2 = \|(S_n - S_m)f\|^2 = \left\| f, \sum_{\alpha, \beta=m}^n \Omega_-^{(\alpha)} \Omega_+^{(\alpha)*} \Omega_+^{(\beta)} \Omega_-^{(\beta)*} f \right\|.$$

The general term under the summation sign can be simplified by using Equation (5.7) and (5.8).

$$\Omega_-^{(\alpha)} \Omega_+^{(\alpha)*} \Omega_+^{(\beta)} \Omega_-^{(\beta)*} = \delta_{\alpha\beta} \Omega_-^{(\alpha)} E_{\alpha} \Omega_-^{(\beta)*} = \delta_{\alpha\beta} \Omega_-^{(\alpha)} \Omega_-^{(\alpha)*} = \delta_{\alpha\beta} F_-^{(\alpha)}. \quad (6.4)$$

Hence

$$\|f_n - f_m\|^2 = \left\| f, \sum_{\alpha=m}^n F_-^{(\alpha)} f \right\|. \quad (6.5)$$

According to Equation (5.9) the projections are all mutually orthogonal. Hence

$$G_n \equiv \sum_{\alpha=1}^n F_-^{(\alpha)}$$

is a sequence of non-decreasing projections. Such a sequence converges strongly to a projection. The sequence of elements

$$g_n \equiv G_n f$$

is therefore a Cauchy sequence and since

$$\|f_n - f_m\|^2 = \|g_n - g_m\|^2$$

$f_n$  is also a Cauchy sequence. The strong limit  $n \rightarrow \infty$  exists

$$g = \lim_{n \rightarrow \infty} f_n$$

and the correspondence

$$f \rightarrow g = Sf$$

defines the linear operator

$$S = \sum_{\alpha=1}^{\infty} \Omega_{+}^{(\alpha)} \Omega_{-}^{(\alpha)*}. \quad (6.6)$$

This is the scattering operator.

Next we prove that  $S$  is quasi-unitary:

$$S^*S = SS^* = E_N. \quad (6.7)$$

For instance, using the identity (6.4), we find with (5.20)

$$S^*S = \sum_{\alpha=1}^{\infty} F_{-}^{(\alpha)} = E_N \quad (6.8)$$

$$SS^* = \sum_{\alpha=1}^{\infty} F_{+}^{(\alpha)} = E_N. \quad (6.9)$$

Since  $V_t$  commutes with all terms in the sum (6.6) (cf. Equation (5.22)) we have also

$$[V_t, S] = 0. \quad (6.10)$$

The following relations are also useful

$$\Omega_{-}^{(\beta)*} S \Omega_{-}^{(\alpha)} = \Omega_{+}^{(\beta)*} S \Omega_{+}^{(\alpha)} = \Omega_{-}^{(\beta)*} \Omega_{+}^{(\alpha)}. \quad (6.11)$$

The physical interpretation of the scattering operator is obtained from the matrix elements. Let the system be at  $t \rightarrow -\infty$  in a channel  $\alpha$ . With this we mean that it approaches in the norm for  $t \rightarrow -\infty$  the state  $U_t^{(\alpha)} f_{\alpha}$ , where  $f_{\alpha} \in D_{\alpha}$  and  $\|f_{\alpha}\| = 1$ . Because of the basic property of scattering systems, this also approaches in the norm  $V_t f_{+}^{(\alpha)}$  as  $t \rightarrow -\infty$ . The probability for finding the system at time  $t$  in the state  $U_t^{(\alpha)} f_{\beta}$  ( $f_{\beta} \in D_{\beta}$ ) of channel  $\beta$  is then

$$P_{\beta\alpha}(t) = |(U_t^{(\beta)} f, V_t f_{+}^{(\alpha)})|^2.$$

In the limit  $t \rightarrow \infty$  we have

$$(U_t^{(\beta)} f_\beta, V_t f_+^{(\alpha)}) = (V_t^* U_t^{(\beta)} f_\beta, f_+^{(\alpha)}) \rightarrow (f_-^{(\beta)}, f_+^{(\alpha)}).$$

We can write the last expression in the equivalent forms (cf. Equation (6.11))

$$\begin{aligned} (f_-^{(\beta)}, f_+^{(\alpha)}) &= (f_-^{(\beta)}, S f_-^{(\alpha)}) = (f_+^{(\beta)}, S f_+^{(\alpha)}) \\ &= (f_\beta, \Omega_-^{(\beta)*} S \Omega_-^{(\alpha)} f_\alpha) \\ &= (f_\beta, \Omega_+^{(\beta)*} S \Omega_+^{(\alpha)} f_\alpha). \end{aligned} \quad (6.12)$$

These expressions lead to the usual formulae for scattering cross-sections and decay time when properly specialized.

## 7. Asymptotic properties of observables

In this section we shall investigate the question how the asymptotic behaviour of observables is related to the  $S$ -operator in a general scattering system. This question is of interest because the asymptotic operators have often been introduced as auxiliary quantities in the definition of the scattering operator<sup>6), 7)</sup>. In fact in more recent works the tendency has been to formulate the scattering theory in such a manner that no explicit reference is made to the free Hamiltonian describing 'bare' particles. Instead one attempts to replace it by the asymptotic properties of a sufficiently complete set of observables (or field operators). The relation between these observables in the past and the future is then used for the definition of the scattering operator<sup>3), 8)</sup>.

We shall examine here the asymptotic properties of observables from the point of view of the rigorous scattering theory.

Let  $A$  be a self-adjoint, bounded operator in  $\mathfrak{H}$ , representing an observable. We shall assume that  $A$  is independent of time in the Schrödinger picture in which state vectors change in time according to

$$f_t = V_t f. \quad (7.1)$$

In the Heisenberg picture the operator  $A$  varies in time according to

$$A_t = V_t^* A V_t \quad (7.2)$$

while the state vectors are constant.

We shall first discuss the case of a 'simple scattering system', that is, a system with one channel only<sup>1)</sup>. The transformation operator for this channel shall be denoted, as before, by  $U_t = e^{-iH_0 t}$ .

The asymptotic condition as usually formulated is to assume that with every operator  $A$  there are associated two operators  $A_{in}$  and  $A_{out}$  which

in some sense, to be made precise, represent the operator  $A$  in the distant past and the remote future.

The  $A_{in}$  and  $A_{out}$  are considered the observables of free particle motion. The definition of the operators  $A_{in}$  and  $A_{out}$  is not unique even for the case of simple scattering systems. The ambiguity is related to the ambiguity in the definition of the scattering operator discussed in section 2 of this paper.

One sense in which the asymptotic condition can be made precise is to require the weak convergence of the operators  $U_t A_t U_t^*$  to the constant operators  $A_{in}$  and  $A_{out}$  for the respective limits  $t \rightarrow -\infty$ , and  $t \rightarrow +\infty$ .

In other words, for any two elements  $f, g \in \mathfrak{H}$  we shall have

$$\begin{aligned} (f, U_t A_t U_t^* g) &\rightarrow (f, A_{in} g) \quad \text{for } t \rightarrow -\infty, \\ \text{and} \quad (f, U_t A_t U_t^* g) &\rightarrow (f, A_{out} g) \quad \text{for } t \rightarrow +\infty. \end{aligned} \quad (7.3)$$

We shall now examine whether this condition is in fact satisfied for simple scattering systems and whether it can be used for the definition of the S-operator.

From the identity

$$U_t V_t^* U_t^* - \Omega_-^* A \Omega_- = U_t V_t^* A (V_t U_t^* - \Omega_-) + (U_t V_t^* - \Omega_-^*) A \Omega_-$$

we find

$$\begin{aligned} &| (f, (U_t A_t U_t^* - \Omega_-^* A \Omega_-) g) | \\ &\leq | (f, U_t V_t^* A (V_t U_t^* - \Omega_-) g) | + | (f, (U_t V_t^* - \Omega_-^*) A \Omega_- g) |. \end{aligned}$$

For the first term we have by Schwartz' inequality

$$\begin{aligned} &| (f, U_t V_t^* A (V_t U_t^* - \Omega_-) g) | \\ &\leq \| f \| \quad \| A \| \quad \| (V_t U_t^* - \Omega_-) g \| \end{aligned}$$

This term tends to zero with  $t \rightarrow -\infty$  for all  $f, g \in \mathfrak{H}$  as a consequence of the basic property of a simple scattering system.

For the second term we have

$$\begin{aligned} &| (f, (U_t V_t^* - \Omega_-^*) A \Omega_- g) | \\ &\leq \| f \| \quad \| A \| \quad \| (V_t U_t^* - \Omega_-) g \| \end{aligned}$$

and this is again convergent to zero for  $t \rightarrow -\infty$ . We have thus verified the weak convergence of

$$U_t A_t U_t^* \text{ to } A_{in} \equiv \Omega_-^* A \Omega_- \text{ for } t \rightarrow -\infty.$$

We remark that we could have asserted strong convergence for operators  $A$  which leave the subspace  $N$  of continuum states invariant. Indeed we have

$$\begin{aligned} \|(U_t A_t U_t^* - A_{\text{in}})f\| &\leq \|U_t V_t^* A(\Omega_- - V_t U_t^*)f\| \\ &+ \|(\Omega_-^* - U_t V_t^*) A \Omega_- f\|. \end{aligned}$$

The first term converges to zero without restriction for  $t \rightarrow -\infty$ . For the second term on the other hand we can only assert convergence if  $A \Omega_- f \in N$ . Since  $\Omega_- f \in N$  this is assured if  $N$  is invariant under  $A$ .

If  $N = \mathfrak{S}$ , that is if there are no bound states, this is no restriction and strong convergence can always be asserted in that case.

In a similar way one can discuss the operators  $A_{\text{out}}$  and the convergence of the corresponding expressions in the limit  $t \rightarrow +\infty$ . The relation between  $A_{\text{in}}$  and  $A_{\text{out}}$  is then obtained from the two defining equations

$$\begin{aligned} A_{\text{in}} &= \Omega_-^* A \Omega_- \\ A_{\text{out}} &= \Omega_+^* A \Omega_+. \end{aligned} \tag{7.4}$$

If we multiply the second Equation (7.4) from the left with  $\Omega_+$  and from the right with  $\Omega_+^*$ , we obtain first

$$\Omega_+ A_{\text{out}} \Omega_+^* = F A F \tag{7.5}$$

where  $F$  is the left projection of  $\Omega_+$  with range  $N$ . Multiplying now (7.5) from the left with  $\Omega_-^*$  and from the right with  $\Omega_-$  we obtain

$$\Omega_-^* \Omega_+ A_{\text{out}} \Omega_+^* \Omega_- = A_{\text{in}}. \tag{7.6}$$

Here we have used the fact that the left projection of  $\Omega_{\pm}$  are the same for simple scattering systems. This leads to the definition of the  $S$ -operator according to

$$A_{\text{out}} = S^* A_{\text{in}} S \tag{7.7}$$

with  $S = \Omega_-^* \Omega_+$ , in agreement with Equation (9) reference<sup>8</sup>). The  $S$ -operator obtained in this manner is the  $S$ -operator which does not allow a generalization to the multichannel case. The definition of  $S$  through Equations (7.3) and (7.5) is therefore only suitable for the 'simple scattering systems'.

A different definition of  $A_{\text{in}}$  and  $A_{\text{out}}$  which is used by EKSTEIN<sup>3</sup>), allows a generalization to the multichannel case. We shall again use the same notation as before even though the operators to be defined now differ from the ones discussed so far.

We verify first that for  $f, g \in N$

$$(f, A_t g) \rightarrow (f, \Omega_- A(t) \Omega_-^* g) \quad \text{for } t \rightarrow +\infty. \quad (7.8)$$

Where

$$A_t = V_t^* A V_t$$

$$A(t) = U_t^* A U_t.$$

Indeed

$$| (f, A_t g) - (f, \Omega_- A(t) \Omega_-^* g) |$$

$$\leq | (f, V_t^* A (V_t - U_t \Omega_-^*) g) | + | (f, (V_t^* - \Omega_- U_t^*) A U_t \Omega_-^* g) |.$$

For the first term we have

$$| (f, V_t^* A (V_t - U_t \Omega_-^*) g) | \leq \| f \| \| A \| \| (I - V_t^* U_t \Omega_-^*) g \|.$$

Since  $g \in N$

$$\lim_{t \rightarrow +\infty} V_t^* U_t \Omega_-^* g = \Omega_- \Omega_-^* g = F g = g.$$

Therefore the right hand side vanishes with  $t \rightarrow +\infty$ . For the second term we obtain

$$| (f, (V_t^* - \Omega_- U_t^*) A U_t \Omega_-^* g) |$$

$$\leq \| (I - V_t^* U_t \Omega_-^*) f \| \| A \| \| g \|$$

and this vanishes for  $t \rightarrow \infty$  because  $f \in N$ . Thus relation (7.8) is established. A similar reasoning for  $t \rightarrow -\infty$  leads to a corresponding relation. We can then define two time dependent observables  $A_{in}(t)$  and  $A_{out}(t)$  according to

$$\left. \begin{aligned} A_{out}(t) &\equiv \Omega_- A(t) \Omega_-^* \\ A_{in}(t) &= \Omega_+ A(t) \Omega_+^* \end{aligned} \right\} \quad (7.9)$$

As was pointed out by EKSTEIN<sup>3)</sup> these are not 'free' operators since their time dependence is only apparently governed by the free transformation operator  $U_t$  as may be seen from

$$\Omega_- A(t) \Omega_-^* = \Omega_- U_t^* A U_t \Omega_-^* = V_t^* \Omega_- A \Omega_-^* V_t.$$

By eliminating from (7.9) the operators  $A(t)$ , we obtain

$$A_{out}(t) = \Omega_- \Omega_+^* A_{in}(t) \Omega_+ \Omega_-^* \quad (7.10)$$

or

$$A_{out}(t) = S^* A_{in}(t) S, \quad (7.11)$$

with

$$S = \Omega_+ \Omega_-^*. \quad (7.12)$$



The similarity of (7.11) with (7.7) is superficial and misleading (as is the notation) since all the quantities involved in (7.11) are quite different from the ones occurring in (7.7). We shall not refer to (7.7) any more in the rest of the paper.

The generalization of these considerations to the multichannel case are now fairly straightforward. We need to show for instance the weak convergence for  $f, g \in N$

$$(f, A_t g) \rightarrow (f, A_{\text{out}}(t) g) \quad \text{for } t \rightarrow +\infty, \quad (7.13)$$

with

$$A_{\text{out}}(t) = \sum_{\alpha} \Omega_{-}^{(\alpha)} U_t^{(\alpha)*} A \sum_{\beta} U_t^{(\beta)} \Omega_{-}^{(\beta)*}. \quad (7.14)$$

To show this we use formula (5.20), and obtain for instance

$$g = E_N g = \sum_{\alpha} F_{-}^{(\alpha)} g, \quad (7.15)$$

and a similar decomposition for  $f$ . We shall write  $f^{(\beta)} = F_{-}^{(\beta)} f$  and  $g^{(\beta)} = F_{-}^{(\beta)} g$ , so that

$$(f, A_t g) = \sum_{\alpha, \beta} (f^{(\alpha)}, A_t g^{(\beta)}), \quad (7.16)$$

and show that the general term under the summation sign converges to  $(f^{(\alpha)}, A_{\text{out}}(t) g^{(\beta)})$  for  $t \rightarrow +\infty$ . This last expression may also be written as

$$(f^{(\alpha)}, \Omega_{-}^{(\alpha)} U_t^{(\alpha)*} A U_t^{(\beta)} \Omega_{-}^{(\beta)}) \quad \text{since } F_{-}^{(\alpha')} \Omega_{-}^{(\alpha)} = \delta_{\alpha' \alpha} \Omega_{-}^{(\alpha)}.$$

The stated convergence is now proved by writing

$$\begin{aligned} & | (f^{(\alpha)}, (A_t - A_{\text{out}}(t)) g^{(\beta)}) | \\ & \leq | (f^{(\alpha)}, V_t^* A (V_t - U_t^{(\beta)} \Omega_{-}^{(\beta)}) g^{(\beta)}) | \\ & + | (f^{(\alpha)}, (V_t^* - \Omega_{-}^{(\alpha)} U_{+}^{(\alpha)*}) A U_t^{(\beta)} \Omega_{-}^{(\beta)*} g^{(\beta)}) |. \end{aligned}$$

The vanishing of the first factor in the limit  $t \rightarrow +\infty$  follows now from

$$\| (V_t - U_t^{(\beta)} \Omega_{-}^{(\beta)}) g^{(\beta)} \| \rightarrow 0 \quad \text{for } t \rightarrow +\infty.$$

Similarly one concludes the vanishing of the second factor.

We summarize the result for convenience:

$$(f, A_t g) \rightarrow (f, A_{\text{out}}^{\text{in}}(t) g) \quad \text{for } t \rightarrow \pm\infty \quad (7.17)$$

and  $f, g \in N$ .

$$A_{\text{out}}^{\text{in}}(t) = \sum_{\alpha} \Omega_{\mp}^{(\alpha)} U_t^{(\alpha)*} A \sum_{\beta} U_t^{(\beta)} \Omega_{\mp}^{(\beta)*}. \quad (7.18)$$

The relation between  $A_{out}(t)$  and  $A_{in}(t)$  is exactly as in the single channel case

$$A_{out}(t) = S^* A_{in}(t) S, \quad (7.19)$$

with  $S$  now defined as

$$S = \sum_{\alpha} \Omega_{+}^{(\alpha)} \Omega_{-}^{(\alpha)*}. \quad (7.20)$$

To prove this we eliminate  $A$  from the two relations (7.18). For instance we obtain by multiplying from the left with  $\Omega_{+}^{(\alpha)*}$  and from the right with  $\Omega_{+}^{(\beta)}$  using (5.21)

$$\Omega_{+}^{(\alpha)*} A_{in}(t) \Omega_{+}^{(\beta)} = E_{\alpha} U_t^{(\alpha)*} A U_t^{(\beta)} E_{\beta}.$$

This relation we multiply next from the left with  $\Omega_{-}^{(\alpha)}$  and from the right with  $\Omega_{-}^{(\beta)*}$  and sum over the indices  $\alpha$  and  $\beta$ . Using the fact that  $E_{\alpha}$  is right projection of  $\Omega_{\alpha}$  we find

$$S^* A_{in}(t) S = A_{out}(t)$$

with  $S$  given by (7.20).

The question needs to be examined whether this procedure can be used for the definition of the scattering operator. The answer is yes if the asymptotic operators  $A_{out}(t)$  are known for a sufficiently large set of operators in  $\mathcal{A}$  so that they generate in  $N$  an irreducible operator ring. The  $S$ -operator is then uniquely determined by (7.19) in the subspace  $N$  up to a numerical factor of magnitude 1.

It is also possible to introduce time-independent operators  $A_{in}$  and  $A_{out}$  by defining

$$A_{out, in}(t) = V_t^* A_{out, in} V_t. \quad (7.21)$$

By using the intertwining property (5.22) one obtains

$$A_{out, in} = \sum_{\alpha} \Omega_{+}^{(\alpha)} A \sum_{\beta} \Omega_{+}^{(\beta)*}, \quad (7.22)$$

and

$$A_{out} = S^* A_{in} S. \quad (7.23)$$

### Appendix

In reference<sup>1)</sup> we have stated the following lemma: let  $V_t = e^{-iHt}$  and  $U_t = e^{-iH_0 t}$  be the unitary transformation groups associated with a 'simple scattering system'. Let  $\Omega$  denote either  $\Omega_{+}$  or  $\Omega_{-}$  and let  $\Omega_1$  be any bounded intertwining operator such that

$$V_t \Omega_1 = \Omega_1 U_t. \quad (1)$$

Denote by  $R$  the range of  $\Omega$  (which is common to  $\Omega_+$  and  $\Omega_-$ ) and by  $R_1$  the range of  $\Omega_1$  then

$$R_1 \subseteq R. \quad (2)$$

The proof given in reference<sup>1)</sup> was incorrect as indicated in the footnote. We shall here give a correct proof.

We remark first of all that it is sufficient to prove the theorem for intertwining operators which are isometries. This simplification is possible because if  $T$  is any bounded intertwining operator, then there exists another one  $W$  which is an isometry and which has exactly the same range as  $T$ .

This is a simple consequence of the well-known theorem on the polar decomposition of bounded linear operators<sup>9)</sup>. According to this theorem one can associate with every bounded linear operator  $T$  a unique linear isometry  $W$  with the same left and right projections as  $T$ . It is related to  $T$  by the formulas

$$T = W R \quad (3)$$

and

$$R = (T^* T)^{\frac{1}{2}}. \quad (4)$$

The operator  $W$  is defined as follows: on all elements  $g$  of the form  $g = Rf$ , we have  $Wg = Tf$ . On the linear manifold  $R(\mathfrak{H})$  of such elements  $W$  is homogeneous, additive and isometric. Such a transformation can be uniquely extended by continuity to the closed linear manifold  $\overline{R(\mathfrak{H})}$ , preserving these properties. On the orthogonal complement of this subspace  $W$  is zero by definition.  $W$  is then a partial isometry. Its left and right projections are the same as the left and right projections of  $T$ .

To apply this theorem for our case we assume that  $T$  is an intertwining operator such that

$$V_t T = T U_t. \quad (5)$$

We aim to show that then also

$$V_t W = W U_t \quad (6)$$

where  $W$  is the partial isometry just defined. In order to see this we evaluate the left-hand side of (6) as follows. Let the arbitrary  $f$  be decomposed  $f = g + h$  with  $g \in R(\mathfrak{H})$  and  $h \in \overline{R(\mathfrak{H})}^\perp$ . We have then  $Wf = Wg$ . Since  $g$  is of the form  $Ru$  with  $u \in \mathfrak{H}$ , or a limit point of such elements we have

$$Wf = Tu \text{ or the corresponding limiting relation.}$$

Consequently

$$V_t Wf = V_t Tu = T U_t u \quad (7)$$

On the other hand we have

$$U_t f = U_t g + U_t h, \quad U_t g \in \overline{R(\mathfrak{H})} \quad \text{and} \quad U_t h \in R(\mathfrak{H})^\perp$$

because  $R$  commutes with  $U_t$ .

Therefore

$$WU_t f = WU_t g = WU_t Ru = WR U_t u = T U_t u. \quad (8)$$

Combining (7) and (8) we find for all  $f$

$$V_t W f = W U_t f. \quad (9)$$

This is the content of Equation (6). It is thus sufficient to prove the theorem for partial isometries.

Let now  $\Omega$ , be a partial isometry with the intertwining property

$$V_t \Omega_1 = \Omega_1 U_t. \quad (1)$$

The relation (1) can be extended to a certain class of bounded functions of  $V_t$  and  $U_t$  by a standard procedure\*). In particular if  $F_\lambda$  and  $E_\lambda$  are the spectral resolutions associated with  $V_t$  and  $U_t$  such that

$$U_t = \int_{-\infty}^{+\infty} e^{-t\lambda i} dE_\lambda \quad (10)$$

$$V_t = \int_{-\infty}^{+\infty} e^{-i\lambda t} dF_\lambda. \quad (11)$$

Then we have also

$$F_\lambda \Omega_1 = \Omega_1 E_\lambda \quad (12)$$

from which follows

$$F_\lambda F_1 = \Omega_1 E_\lambda \Omega_1^* = F_1 F_\lambda \quad (13)$$

where

$$F_1 = \Omega_1 \Omega_1^* \quad (14)$$

is the left-projection of  $\Omega_1$ .

Consider now an element  $f$  in the range of  $\Omega_1$  such that

$$f = F_1 f. \quad (15)$$

We decompose it according to

$$f = g + h$$

with  $g \in R \equiv N$  and  $h \in N^\perp = M$ .

We wish to show that  $h = 0$  for arbitrary choice of  $f \in R_1$ .

\*) See for instance ref. 4), p. 341.

Let  $\lambda_i$  be the eigenvalues of the operators  $V_t$  and

$$P_i \equiv F_{\lambda_i+0} - F_{\lambda_i-0}$$

the corresponding projection operators. The total projection  $P$  corresponding to the subspace  $M$  of bound states is then the sum

$$P = \sum_i P_i$$

Since  $h \in M$  we have  $Ph = h$  and since  $h = Pf$  and  $[P, F_1] = 0$ , we also have

$$F_1 h = F_1 Pf = P F_1 f = Pf = h \quad (16)$$

Thus  $h$  is also in the range of  $\Omega_1$ .

We express now the norm of  $h$  in the following manner

$$\begin{aligned} \|h\|^2 &= (h, Ph) = (h, \sum_i P_i h) = (h, \sum_i P_i F_1 h) \\ &= (h, \Omega_1 \sum_i (E_{\lambda_i+0} - E_{\lambda_i-0}) \Omega_1^* h), \text{ or finally} \\ \|h\|^2 &= (\Omega_1^* h, \sum_i (E_{\lambda_i+0} - E_{\lambda_i-0}) \Omega_1^* h) \end{aligned}$$

The right-hand side is zero, because  $U_t$  has a spectral revolution with no discontinuities. Therefore  $h = 0$  for all  $f$ , or  $f \in R$ , or finally  $R_1 \subseteq R$ , q.e.d.

### Literaturverzeichnis

- 1) J. M. JAUCH, *Helv. Phys. Acta*, **31**, 127 (1958).
- 2) H. EKSTEIN, *Phys. Rev.* **101**, 880 (1956).
- 3) H. EKSTEIN, *Nuovo Cimento* **4**, 1017 (1956).
- 4) See for instance F. RIESZ and B. SZ.-NAGY, *Functional Analysis*, F. UNGAR, Publishing Co., New York 1955, p. 201.
- 5) J. v. NEUMANN, *Proc. Nat. Acad.* **18**, 70 (1932).
- 6) C. N. YANG and D. FELDMAN, *Phys. Rev.* **79**, 972 (1950).
- 7) G. KÄLLÈN, *Ark. Fys.* **2**, 187, 371 (1952).
- 8) H. LEHMANN, K. SYMANZIK, and W. ZIMMERMANN, *Nuovo Cim.* **1**, 205 (1955).
- 9) See reference<sup>4)</sup>, p. 286.