

Zeitschrift: Helvetica Physica Acta
Band: 34 (1961)
Heft: VI-VII

Artikel: Quantum theory in real Hilbert space. III, Fields of the first kind (linear field operators)
Autor: Stueckelberg, E.C.G. / Guenin, M. / Piron, C.
DOI: <https://doi.org/10.5169/seals-113192>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Quantum Theory in Real Hilbert Space III: Fields of the 1st kind (Linear Field Operators)

by E. C. G. Stueckelberg, M. Guenin, C. Piron and H. Ruegg*)

(Universities of Geneva and Lausanne)

(3. V. 1961)

Abstract: The method of RHS (real Hilbert space, see^{1) 2)}) is applied to the free scalar and spinor fields. We remark, that *two kinds of fields exist*:

Fields of the 1st kind commute with \bar{J} ($\rightarrow i = \sqrt{-1}$ in CHS (complex Hilbert space)). In CHS, they are linear operators.

Fields of the 2nd kind anti-commute with \bar{J} . In CHS, they are anti-linear operators.

The general formulas of this article are valid for both cases. In this publication, only the fields of the 1st kind are explicitly discussed. The relation between *statistics* and *strong time reflection* (CT) are clarified. Furthermore, a concise formulation of *contragredient four-spinors* is given. Some well known formulas are explicitly restated in RHS in order to show the difference between fields of the 1st kind and fields of the 2nd kind³⁾.

§ 1. Field Operators

We consider the *scalar field* $w(x)$ ($\neq w^T(x)$) and the *N-component spinor field* $\psi^A(x)$ ($\neq \psi^{TA}(x)$, $AB \dots = 12 \dots N$). Field operators satisfy the *wave equation*

$$(\square - M^2) w(x) = (\square - M^2) \psi^A(x) = 0, \quad (1.1)$$

$$\square = -\text{sig}(g^{nn}) g^{\alpha\beta} \partial_\alpha \partial_\beta = \Delta - \partial_t^2,$$

$$\Delta = \sum_1^d (\partial_i)^2; \quad d = n - 1; \quad \partial_n = \partial_t. \quad (1.2)$$

Observables $F^{\alpha \dots}(x)$ are *bi-linear forms* in $w(x)$ and $w^T(x)$ (or $\psi^A(x)$ and $\psi^{TA}(x)$) and their derivatives (involving numbers or \bar{J} -dependent operators):

$$w_\alpha(x) = \partial_\alpha w(x); \quad \psi_\alpha^A(x) \equiv \partial_\alpha \psi^A(x). \quad (1.3)$$

There exist two kinds of observables, $F^{(1)\alpha \dots}(x)$ and $F^{(2)\alpha \dots}(x)$, depending on whether the transposed field operator operates *after* ($F^{(1)}$) or *before* ($F^{(2)}$)

*) Supported by the Swiss National Research Fund.

the untransposed operators. Observables are *symmetric operators* in RHS ($F^T = F$), which *commute* with \check{J} . From

$$[A B, C] = A [B, C]_{\mp} \pm [A, C]_{\mp} B \quad (1.4)^*)$$

follows, that fields either *commute* (fields of the 1st kind)

$$[\check{J}, w(x)] = 0; \quad [\check{J}, \psi^A(x)] = 0 \quad (1.5, 1^{\text{st}} \text{ k.})$$

or *anti-commute* (fields of the 2nd kind) with \check{J} .

$$(\check{J}, \bar{w}(x)) = 0; \quad (\check{J}, \bar{\psi}^A(x)) = 0. \quad (1.5, 2^{\text{nd}} \text{ k.})^*)^{**})$$

Using the operators

$$\check{J} = j \times 1, \quad \bar{K} = k \times 1, \quad \check{L} = l \times 1 \quad (1.6)$$

where j , k and l are the pseudoquaternions (see I A-4.9), one concludes from (we write w , for w and ψ^A)

$$w = 1 \times w_{(r)} + j \times w_{(i)} + k \times w_{(k)} + l \times w_{(l)} \quad (1.7)$$

(a form analogous to (I A-2.3)) and its transposed, that only

$$w = 1 \times w_{(r)} + j \times w_{(i)} \quad (1.8, 1^{\text{st}} \text{ k.})$$

or

$$\bar{w} = k \times w_{(k)} + l \times w_{(l)} = (1 \times w_{(k)} + j \times w_{(l)}) (k \times 1) = w \bar{K} \quad (1.8, 2^{\text{nd}} \text{ k.})$$

can occur in the bilinear forms defining observables. *The present article is essentially restricted to fields of the 1st kind.* However some formulas, valid for either kind of fields are included. The discussions of fields of the 2nd kind is reserved to a later publication (see ³)).

For either kind of field, a *phase transformation*

$$'w(x) = e^{\lambda \check{J}} w(x); \quad 'w^T(x) = w^T(x) e^{-\lambda \check{J}} \quad (1.9)$$

leaves the observables invariant.

§ 2. Quantization of the scalar field

We look for an observable $\theta^{\alpha\beta}(x)$, from which \check{H}_{μ} and $\check{M}_{\mu\nu}$ may be constructed (see (I 0.25), (I 0.27)). In principle, bilinear observables, for example the scalar

$$F^{(1)}(x) = w^T(x) w(x) \quad (2.1)$$

*) $[A, B] = [A, B]_- = AB - BA$; $(A, B) = [A, B]_+ = AB + BA$.

**) Operators with a $-$ ($\bar{K}, \bar{L}, \bar{w}, \dots$) anti-commute with \check{J} .

transform according to

$$\begin{aligned} {}'F^{(1)}({}'x) &= (O_{(L)}^{-1} w^T({}'x) O_{(L)}) (O_{(L)}^{-1} w({}'x) O_{(L)}) = \\ &= F^{(1)}(L^{-1} {}'x) = F^{(1)}(x) \end{aligned} \quad (2.2)$$

if

$${}'w({}'x) = O_{(L)}^{-1} w({}'x) O_{(L)} = e^{\lambda \check{J}} w(L^{-1} {}'x). \quad (2.3)$$

For infinitesimal transformations, the phase λ must be zero. The proper Lorentz-group $\{L_{(\text{cont})}\}$, is therefore generated in RHS by (I 5.6) with

$$[\check{J} \check{H}_\mu, w(x)] = -w_\mu(x), \quad (2.4)$$

$$[\check{J} \check{M}_{\mu\nu}, w(x)] = -[x_\mu, \partial_\nu] w(x). \quad (2.5)$$

Thus, fields of the 1st kind transform like scalar observables. (For fields of the 2nd kind, it is essential that \check{J} is *inside* of the commutators (2.4) and (2.5)!) For pseudochronous and pseudochorous transformations, a phase factor may occur.

We have, for the most general $\theta^{\alpha\beta}$, the form:

$$\theta^{\alpha\beta} = \alpha_1 \theta^{(1)\alpha\beta} + \alpha_2 \theta^{(2)\alpha\beta}, \quad (2.6)$$

$$\theta^{(1)\alpha\beta}(x) = (w^{T\alpha} w^\beta + w^\beta w^{T\alpha} - g^{\alpha\beta} (w_\rho^T w^\rho + M^2 w^T w)) (x), \quad (2.7^{(1)})$$

$$\theta^{(2)\alpha\beta}(x) = (w^\alpha w^{T\beta} + w^\beta w^{T\alpha} - g^{\alpha\beta} (w_\rho w^{T\rho} + M^2 w w^T)) (x) \quad (2.7^{(2)})$$

and (2.4) takes the form

$$\begin{aligned} & - [\check{J} \check{H}_\mu, w(y')] = \\ &= - \int d\check{\sigma}_\alpha(y) \{ \alpha_1 [\check{J}(w^{T\alpha} w_\mu) (y), w(y')] + \alpha_2 [\check{J}(w^\alpha w_\mu^T) (y), w(y')] \\ & \quad + \alpha_1 [\check{J}(w_\mu^T w^\alpha) (y), w(y')] + \alpha_2 [\check{J}(w_\mu w^{T\alpha}) (y), w(y')] \} \\ & + \int d\check{\sigma}_\mu(y) \{ \alpha_1 [\check{J}(w_\alpha^T w^\alpha) (y), w(y')] + \alpha_2 [\check{J}(w^\alpha w_\alpha^T) (y), w(y')] \} \\ & + M^2 \int d\check{\sigma}_\mu(y) \{ \alpha_1 [\check{J}(w^T w) (y), w(y')] + \alpha_2 [\check{J}(w w^T) (y), w(y')] \} \\ & \equiv w_\mu(y'). \end{aligned} \quad (2.8)$$

We chose $\tau(y) = \tau(y') = 0$ i.e. y and y' are events on the same hypersurface with a time like normal $\check{\nu}^\alpha(y)$ ($d\check{\sigma}^\alpha(y) = \check{\nu}^\alpha(y) d\sigma(y)$; $(\check{\nu}^\alpha \check{\nu}_\alpha)(y) = \text{sig}(g^{nn})$). w , w_μ , w_μ^T and w^T being linearly independent, all terms, except

the first and fourth terms, have to cancel out. In particular, the last integral proportional to M^2 has to vanish. This leads to:

$$\alpha_1 [\check{J}(w^T w)(y), w(y')] + \alpha_2 [\check{J}(w w^T)(y), w(y')] = 0, \quad (2.9)$$

$$\begin{aligned} & - \alpha_1 [\check{J}(w^{T\alpha} w_\mu)(y), w(y')] - \alpha_2 [\check{J}(w_\mu w^{T\alpha})(y), w(y')] = \\ & = \check{\delta}^\alpha(y y') w_\mu(y). \end{aligned} \quad (2.10)$$

Where $\check{\delta}^\alpha(y y')$ is the (pseudochronous) δ -function on the surface $\tau(y) = 0$:

$$\check{\delta}^\alpha(y y') = \check{\delta}^\alpha(y' y); \quad \int d\check{\sigma}_\alpha(y) \check{\delta}^\alpha(y' y) f(y) = f(y'). \quad (2.11)$$

In order that the 2nd and 3rd term cancel against the second integral, the symmetry condition, compatible with (2.11)

$$\int d\check{\sigma}_\mu(y) \check{\delta}_\alpha(y y') f(y) = \text{sig}(g^{nn}) (\check{v}_\mu \check{v}_\alpha)(y') f(y') \quad (2.12)$$

must be satisfied. (2.9) and (2.10) are necessary conditions for (2.4). We may integrate these two conditions, using the *pseudochronous invariant number*, defined by

$$\check{D}^0(x y) = \check{D}^0(x - y) = -\check{D}^0(y x), \quad (2.13)$$

$$(\square_x - M^2) \check{D}^0(x y) = 0, \quad (2.14)$$

$$\check{D}^0(y y') = 0; \quad \partial_\alpha^y \check{D}^0(y y') = -\check{\delta}_\alpha(y y') \quad (2.15) *$$

obtaining

$$\alpha_1 [\check{J} w^T(x) w(z), w(y')] + \alpha_2 [\check{J} w(z) w^T(x), w(y')] = \check{D}^0(x y') w(z). \quad (2.16)$$

A somewhat lengthy calculation shows, that (2.16) is a sufficient condition for (2.5). The CR (commutation relation) (2.16) (although more general) is analogous to the CR proposed by GREEN⁴) and VOLKOV⁵) for spinor fields (see (8.8)).

Let us consider fields of the 1st kind. Applying (1.4), we find:

$$\begin{aligned} & \check{J} \alpha_1 \{w^T(x) [w(z), w(y)]_\mp \pm [w^T(x), w(y)]_\mp w(z)\} + \\ & + \check{J} \alpha_2 \{w(z) [w^T(x), w(y)]_\mp \pm [w(z), w(y)]_\mp w^T(x)\} = \\ & = \check{D}^0(x y) w(z). \end{aligned} \quad (2.17, 1^{\text{st}} \text{ k.})$$

*) The particular choice $\tau(y) = y^n - y'^n = 0$, $d\check{\sigma}_\alpha(y) = (00 \dots 0 d^d y)$; $\check{\delta}^\alpha(y y') = (00 \dots 0 \delta(\vec{y} - \vec{y}'))$ leads to the usual definition of $\check{D}^0(x y)$ (metric (6.14)).

The most simple solution is

$$\boxed{[w(x), w(y)]_{\mp} = 0} \quad (2.18, 1^{\text{st}} \text{ k.})$$

$$\check{J} [w^T(x), w(y)]_{\mp} = \check{D}^0(x, y) \quad (2.19, 1^{\text{st}} \text{ k.})$$

$$\boxed{\pm \alpha_1 + \alpha_2 = 1} \quad (2.20, 1^{\text{st}} \text{ k. } \mp)$$

It will be more convenient to write (2.19) in the more usual form:

$$\boxed{\check{J} [w(x), w^T(y)]_{\mp} = \pm \check{D}^0(x, y)} \quad (2.19, 1^{\text{st}} \text{ k.})$$

§ 3. The Charge Operator for Scalar Fields

The observables

$$\check{j}^{(1)\alpha}(x) = ((\check{J} w)^T w^{\alpha} + w^{T\alpha} (\check{J} w)) (x), \quad (3.1^{(1)}) *$$

$$\check{j}^{(2)\alpha}(x) = -(\check{J} w w^{T\alpha} + w^{\alpha} (\check{J} w)^T) (x) \quad (3.1^{(2)}) *$$

satisfy the continuity equation. We form:

$$\check{j}^{\alpha}(x) = \beta_1 \check{j}^{(1)\alpha}(x) + \beta_2 \check{j}^{(2)\alpha}(x) \quad (3.2)$$

defining thus a $\tau(y)$ -independent scalar

$$Q = \beta_1 Q^{(1)} + \beta_2 Q^{(2)} = \int d\sigma_{\alpha}(y) \check{j}^{\alpha}(y) \quad (3.3)$$

called the *charge* Q . If we require, that the phase transformation (1.9) is an orthogonal transformation in RHS

$$'w(x) = O^{-1}(\lambda) w(x) O(\lambda) = e^{\lambda \check{J}} w(x), \quad (3.4)$$

$$O(\lambda) = e^{\check{J} \lambda Q}. \quad (3.5)$$

We need the CR:

$$\begin{aligned} & -[\check{J} Q, w(y')] = \\ & = -\int d\sigma_{\alpha}(y) \{ \beta_1 [(\check{J}(\check{J} w)^T w^{\alpha})(y), w(y')] + \beta_1 [(\check{J} w^{T\alpha} \check{J} w)(y), w(y')] + \\ & + \beta_2 [(w w^{T\alpha})(y), w(y')] - \beta_2 [(\check{J} w^{\alpha} (\check{J} w)^T)(y), w(y')] \} \equiv \check{J} w(y'). \end{aligned} \quad (3.6)$$

*) The signes have been chosen so as to give, for fields of the 1st kind:

$$\check{j}^{(1)\alpha}(x) = \check{J}^{-1} (w^T w^{\alpha} - w^{T\alpha} w) (x), \quad (3.1^{(1)}, 1^{\text{st}} \text{ k.})$$

$$\check{j}^{(2)\alpha}(x) = \check{J}^{-1} (w w^{\alpha T} - w^{\alpha} w^T) (x). \quad (3.1^{(2)}, 1^{\text{st}} \text{ k.})$$

For fields of the 1st kind, this condition reduces to

$$\begin{aligned}
 & - [Q, w(y')] = \\
 & = \int d\sigma_\alpha(y) \{ \beta_1 [\check{J}(w^T w^\alpha)(y), w(y')] - \beta_2 [\check{J}(w^\alpha w^T)(y), w(y')] - \\
 & \quad - \beta_1 [\check{J}(w^T w^\alpha)(y), w(y')] + \beta_2 [\check{J}(w w^T)(y), w(y')] = \\
 & = \int d\sigma_\alpha(y) \check{\delta}^\alpha(y, y') w(y) = w(y'). \quad (3.6, 1^{\text{st}} \text{ k.})
 \end{aligned}$$

Comparison with (2.16) shows, that the first two terms do not contribute if $\beta_1 = \lambda \alpha_1$ and $\beta_2 = -\lambda \alpha_2$, while the second two terms equal the integral in the third member, if

$$\beta_1 = \alpha_1, \quad \beta_2 = -\alpha_2. \quad (3.7, 1^{\text{st}} \text{ k.})$$

Thus, even for the general CR (2.16), Q defined by (3.7) is the generator of the infinitesimal phase transformation for fields of the 1st kind.

§ 4. Charge Conjugation for the Scalar Field of the 1st kind

For fields of the 1st kind and for the most simple CR's (2.18₋) and (2.19₋), follows that:

$$'w(x) = O_{(C)}^{-1} w(x) O_{(C)} = w^T(x), \quad (4.1)$$

$$'w^T(x) = O_{(C)}^{-1} w^T(x) O_{(C)} = w(x) \quad (4.1)^T$$

is an orthogonal transformation in RHS. The CR's lead to BE-statistics. In other words: $O_{(C)}$ -covariance requires BE-statistics. We have further:

$$O_{(C)}^{-1} \theta^{(1)\alpha\beta}(x) O_{(C)} = \theta^{(2)\alpha\beta}(x), \quad (4.2^{(1)})$$

$$O_{(C)}^{-1} \theta^{(2)\alpha\beta}(x) O_{(C)} = \theta^{(1)\alpha\beta}(x). \quad (4.2^{(2)})$$

Thus, choosing $\alpha_1 = \alpha_2 = 1/2$ in (2.20₋), we find, that

$$\theta^{\alpha\beta}(x) = \frac{1}{2} (\theta^{(1)\alpha\beta} + \theta^{(2)\alpha\beta})(x) \quad (4.3)$$

is invariant with respect to $O_{(C)}$. Furthermore, from

$$O_{(C)}^{-1} \check{j}^{(1)\alpha}(x) O_{(C)} = \check{j}^{(2)\alpha}(x), \quad (4.4^{(1)})$$

$$O_{(C)}^{-1} \check{j}^{(2)\alpha}(x) O_{(C)} = \check{j}^{(1)\alpha}(x) \quad (4.4^{(2)})$$

follows, that on account of (3.7) and (3.3)

$$O_{(C)}^{-1} Q O_{(C)} = -Q \quad (4.5) *$$

i.e. \check{j}^α and Q change sign under $O_{(C)}$.

We may now define two kinds of time reversal

(1) $T \rightarrow O_{(T)}$, *weak time reversal* (T)

$$'w('x) = O_{(T)}^{-1} w('x) O_{(T)} = w(T^{-1} 'x) \quad (4.7) **$$

with respect to which $\check{j}^\alpha(x)$ is a pseudochronous vector, and Q a scalar;

(2) $T \rightarrow O_{(C)} O_{(T)} \equiv O_{(CT)}$, *strong time reversal* (CT)

$$'w('x) = O_{(CT)}^{-1} w('x) O_{(CT)} = w^T(T^{-1} 'x) \quad (4.8)$$

with respect to which $j^\alpha(x)$ is an (ortho)vector and \check{Q} a pseudochronous scalar.

The second definition seems more appropriate because *classical particle theory* (see STUECKELBERG⁸) defines

$$\begin{aligned} j^\alpha(x) &= \int_{-\infty}^{+\infty} d\lambda \dot{z}^\alpha(\lambda) \delta(x - z(\lambda)), \\ \dot{z}^\alpha(\lambda) &= \frac{d}{d\lambda} z^\alpha(\lambda); \quad (\dot{z}_\alpha \dot{z}^\alpha)(\lambda) = \text{sig}(g^{nn}), \\ \check{Q} &= \int d\sigma_\alpha(y) \check{j}^\alpha(y) = \text{sig}(\dot{z}^n(\lambda)). \end{aligned}$$

Thus $O_{(C)}$ -covariance or strong time reversal ($O_{(CT)}$)-covariance decide for BE-statistics in the case of scalar fields (see SCHWINGER⁶) and PAULI⁷).

§ 5. The Development of the Scalar Field (1st kind) in Terms of Positive Frequency Wave Packet Operators

Let us define integral operators Ω and $\Omega^{1/2}$, operating on space functions $f(\vec{x})$ which vanish sufficiently strong for $|\vec{x}| \rightarrow \infty$:

$$\Omega f(\vec{x}) = (M^2 - \Delta)^{1/2} f(\vec{x}) = \int d^d y \Omega(|\vec{x} - \vec{y}|) f(\vec{y}), \quad (5.1)$$

$$\Omega^{1/2} f(\vec{x}) = (M^2 - \Delta)^{1/4} f(\vec{x}) = \int d^d y \Omega^{1/2}(|\vec{x} - \vec{y}|) f(\vec{y}). \quad (5.2)$$

$$*) \quad O_{(C)}^{-1} \check{j}^\alpha(x) O_{(C)} = -\check{j}^\alpha(x). \quad (4.6)$$

**) $'x = Tx \rightarrow \{ 'x^i = x^i; 'x^n = -x^n \}$. $T^{-1} = T$. The arbitrary phase factor may be left out.

The kernels $\Omega(|\vec{z}|)$ are essentially Hankel functions decreasing $\propto \exp(-|M||\vec{z}|)$ for $|\vec{z}| \gg |M|^{-1}$. We note especially

$$\int d^d x g(\vec{x}) \cdot \Omega f(\vec{x}) = \int d^d x g(\vec{x}) \Omega \cdot f(\vec{x}) = \int d^d x (\Omega^{1/2} g) (\Omega^{1/2} f)(\vec{x}). \quad (5.3)^*$$

In terms of Ω , we define *two denumerable sets* $\{u', u'', \dots, u^{(\theta)}, \dots\}$ and $\{v', v'', \dots, v^{(\theta)}, \dots\}$ of *positive frequency wave packet operators* (PFWP's) depending but on the operator \check{J} and vanishing for $|\vec{x}| \rightarrow \infty$. They satisfy:

$$-\partial_t u'(\vec{x}, t) = \partial^n u'(x) = \check{J} \Omega u'(\vec{x}, t). \quad (5.4)$$

These sets are *normalised* in terms of 'matrix elements'

$$j^\alpha(u', v')(x) = -j^\alpha(v', u')(x) = \frac{1}{2} \check{J}^{-1} (u' (\cdot \partial^\alpha - \partial^\alpha \cdot) v')(x), \quad (5.5)^*$$

$$\begin{aligned} Q(u', v') &= \int (d\check{\sigma}_\alpha j^\alpha(u', v'))(y) = \frac{1}{2} \check{J}^{-1} \int d^d y (u' (\cdot \partial^n - \partial^n \cdot) v')(y) = \\ &= \frac{1}{2} \int d^d y (u' (\cdot \Omega - \Omega \cdot) v')(\vec{y}, t) = 0, \end{aligned} \quad (5.6)$$

$$\begin{aligned} Q(u''^T, u') &= -Q(u', u''^T) = \frac{1}{2} \int d^d y (u''^T (\cdot \Omega + \Omega \cdot) u')(\vec{y}, t) = \\ &= \int d^d y ((\Omega^{1/2} u''^T) (\Omega^{1/2} u'))(\vec{y}, t) \equiv \delta_{u'' u'} \geq 0. \end{aligned} \quad (5.7)$$

Furthermore, each set is *complete* if the \check{J} dependent operators:

$$\mathbf{S}_{u'} u'(x) u'^T(y) = D^+(x y) = D^+(x - y), \quad (5.8^+)$$

$$\mathbf{S}_{u'} u'^T(x) u'(y) = D^-(x y) = D^-(x - y) = D^+(x y) = D^{+T}(x y) \quad (5.8^-)$$

depend but on $x - y$ and are invariant with respect to the subgroup $\{L_{(\text{ochr})}\}$. A PF-solution $f^+(x)$ of the wave equation may be expanded in terms of one of the sets:

$$f^+(x) = \mathbf{S}_{u'} u'(x) f_u^+ = \frac{1}{2} \check{J}^{-1} \int d\check{\sigma}_\alpha(y) D^+(x y) (\cdot \partial_y^\alpha - \partial_y^\alpha \cdot) f^+(y). \quad (5.9^+)$$

For NF-solutions:

$$f^-(x) = \mathbf{S}_{v'} v'^T(x) f_v^- = -\frac{1}{2} \check{J}^{-1} \int d\check{\sigma}_\alpha(y) D^-(x y) (\cdot \partial_y^\alpha - \partial_y^\alpha \cdot) f^-(y) \quad (5.9^-)$$

*) Operators $\cdot \partial^\alpha$, $\cdot \Omega$ (with point on the left) operate, in the usual way, *to the right*. Operators $\partial^\alpha \cdot$, $\Omega \cdot$ (with point on the right) operate *to the left*.

holds. The general solution of the wave equation may be written as

$$\begin{aligned} w(x) &= 2^{-1/2} \left(\mathbf{S}_u a_u u'(x) + \mathbf{S}_v b_v^T v'^T(x) \right) = \\ &= - \int \check{d}\sigma_\alpha(y) \check{D}^0(x y) (\partial_y^\alpha - \partial_y^\alpha) w(y) = \\ &= \int \check{d}\sigma_\alpha(y) (-\partial_x^\alpha \check{D}^0(x y) \cdot w(y) - \check{D}^0(x y) w^\alpha(y)) \end{aligned} \quad (5.10)$$

with

$$\check{D}^0(x y) = \frac{1}{2} \check{J} (D^+ - D^-) (x y) = -\check{D}^0(y x) = \check{D}^0(x y). \quad (5.11)$$

A comparison, for $x = y'$, shows, that the *pseudochronous number* (5.11) is identical with the number defined by (2.13), (2.14) and (2.15).

The CR's, resp. ACR's (2.18) and (2.19) imply

$$[a_u, a_u]_{\mp} = [b_v^T, b_v^T]_{\mp} = [a_u, b_v^T]_{\mp} = 0, \quad (5.12)$$

$$\begin{aligned} [a_u, a_u^T]_{\mp} &= \pm \delta_{u'u''}; & [b_v^T, b_v]_{\mp} &= \mp \delta_{v'v''}; \\ [a_u, b_v]_{\mp} &= 0. \end{aligned} \quad (5.13)$$

As a_u, a_u^T and a_u^T, a_u are positive operators, (5.13₊) contains an algebraic contradiction. This is PAULI's⁹⁾ argument for excluding ACR's and FD-statistics for scalar fields.

The CR's can be satisfied in terms of the creation- (a^T) and annihilation-(a)-operators:

$$a = \begin{pmatrix} 0 & 1 & 0 & 0 & . \\ 0 & 0 & \sqrt{2} & 0 & . \\ 0 & 0 & 0 & \sqrt{3} & . \\ . & . & . & . & . \end{pmatrix}; \quad a^T = \begin{pmatrix} 0 & 0 & 0 & . \\ 1 & 0 & 0 & . \\ 0 & \sqrt{2} & 0 & . \\ . & . & . & . \end{pmatrix}; \quad N = a^T a = \begin{pmatrix} 0 & 0 & 0 & . \\ 0 & 1 & 0 & . \\ 0 & 0 & 2 & . \\ . & . & . & . \end{pmatrix} \quad (5.14)$$

writing

$$\left. \begin{aligned} a_{u(q)} &= 1 \times (1 \times 1) \times (1 \times 1) \times \cdots \times (a \times 1) \times (1 \times 1) \times \cdots \\ b_{v(q)} &= 1 \times (1 \times 1) \times (1 \times 1) \times \cdots \times (1 \times a) \times (1 \times 1) \times \cdots \end{aligned} \right\} \quad (5.15)$$

The eigenvalues of $N_u = a_u^T a_u$ are the non-negative integers and the charge has the form

$$Q = \mathbf{S}_u \left(N_u + \frac{1}{2} \right) - \mathbf{S}_v \left(N_v + \frac{1}{2} \right) = \mathbf{S}_u N_u - \mathbf{S}_v N_v. \quad (5.16)$$

No 'zero-point charge' appears because a 1 to 1 correspondence between the sets $\{u'\}$ and $\{v'\}$ can be established.

Charge conjugation can be written explicitly, if the sets $\{u'\}$ and $\{v'\}$ are chosen *identical*:

$$\begin{aligned} 'a_{u'} &= O_{(C)}^{-1} a_{u'} \quad O_{(C)} = b_{v'} , \\ 'b_{v'} &= O_{(C)}^{-1} b_{v'} \quad O_{(C)} = a_{u'} \end{aligned} \quad (5.17)$$

from which:

$$O_{(C)} = O_{(C)}^{-1} = O_{(C)}^T \quad (5.18)$$

follows. The operator is

$$O_{(C)} = \prod_{u'} e^{-\pi/2 (a_{u'}^T b_{v'} - b_{v'}^T a_{u'})} e^{\pm \check{J} \pi N_{u'}}. \quad (5.19)$$

In order to quantize explicitly $\check{\Pi}_\mu$, denumerable sets, satisfying

$$\begin{aligned} \partial^\alpha u'(x) &\cong \check{J} \check{k}'^\alpha u'(x), \\ \check{k}'_\alpha \check{k}'^\alpha + M^2 &= 0; \quad \check{k}'^n \geq |M| \end{aligned} \quad (5.20)$$

have to be chosen. Taking in account some additional orthogonality relations, one finds

$$\check{\Pi}^\mu \cong \mathbf{S}_{u'} \left(N_{u'} + \frac{1}{2} \right) \check{k}'^\mu + \mathbf{S}_{v'} \left(N_{v'} + \frac{1}{2} \right) \check{k}'^\mu. \quad (5.21)$$

(5.21) is symmetric for particle and antiparticle states. Aside from the infinite 'zero-point contribution', the energy spectrum has a lower limit: Thus, thermo-statistics with a positive absolute temperature can be applied¹⁰⁾.

§ 6. Dual Spin-Spaces

In this section we deal exclusively with real and complex numbers: we distinguish between φ^A *contravariant* (ctr) and χ_A *covariant* (cov) *spinors*.

φ^A and χ_A are two dual, N-dimensional *spin spaces* (SS). We shall distinguish between (real) RSS and (complex) CSS. The general case being CSS, all symbols φ^A , χ_A , ... should be written as $\widehat{\varphi}^A$, $\widehat{\chi}_A$, ... For any complex number

$$\widehat{\varrho} = \varrho_{(r)} + i \varrho_{(i)} \rightleftharpoons \varrho = \varrho(\check{J}) = \varrho_{(r)} + \check{J} \varrho_{(i)} \quad (6.1) *$$

gives the relation between the complex number $\widehat{\varrho}$ and the \check{J} -dependent operator $\varrho = \varrho(\check{J})$ (see I, Annex 2).

*) We shall use $\widehat{\varrho}$, $\widehat{\sigma}$, ... for complex numbers (resp. for \check{J} -dependent operators) and λ , μ , ... for real numbers.

SS or A -space is related to physical space-time or α -space by a *mixed A -space bi-spinor α -space vector* $\gamma^{\alpha A}_B$, satisfying

$$\gamma^{\alpha A}_C \gamma^{\beta C}_B + \gamma^{\beta A}_C \gamma^{\alpha C}_B = 2 g^{\alpha\beta} \gamma^{0A}_B \quad (6.2)$$

or, in matrix notation

$$(\gamma^\alpha, \gamma^\beta) = 2 g^{\alpha\beta} \gamma^0, \quad (6.2 M) *$$

where $\gamma^{0A}_B = \delta^A_B$ is the identity bi-spinor. (6.2) with $\hat{\gamma}^{\alpha A}_B$ is an *algebraic equation* and, with $\gamma^{\alpha A}_B$, it is an *operator equation* depending but on J . Two contragredient spinors allow to define a scalar $\chi \varphi = \chi_A \varphi^A$ and a vector $\chi \gamma^\alpha \varphi = \chi_A \gamma^{\alpha A}_B \varphi^B$ in α -space.

We resume, without proof and references, a number of well known theorems:

(1) The ring $\Gamma = \{\gamma^r\}$, $r = 0, \alpha, [\alpha_1 \alpha_2], \dots, [\alpha_1 \alpha_2 \dots \alpha_n] = 1, 2, \dots, 2^n$ is formed from the totally antisymmetric α -space tensors

$$\begin{aligned} \gamma^{\alpha_1 \dots \alpha_\nu} &= \gamma^{[\alpha_1 \dots \alpha_\nu]} = (\nu!)^{-1} \sum_P (-1)^P P(\gamma^{\alpha_1} \dots \gamma^{\alpha_\nu}) = \\ &= \gamma^{\alpha_1} \dots \gamma^{\alpha_\nu} \text{ if all } \alpha_1 \dots \alpha_\nu \text{ are different.} \end{aligned} \quad (6.3 M) **$$

The subsets $\Gamma^{(\nu)} = \{\gamma^{\alpha_1 \dots \alpha_\nu}\}$ contain $\binom{n}{\nu}$ independent elements.

(2) Two irreducible representations of (6.2) γ^α and γ^α are related by

$$' \gamma^\alpha S = S \gamma^\alpha \quad (6.4 M)$$

where S is either the zero matrix or a non-singular square matrix. In particular, for $n = 2m$ (even), all irreducible representations are *equivalent* i.e. we have

$$' \gamma^\alpha = S \gamma^\alpha S^{-1}; \quad S = \{S'^A_A\}; \quad 'N = N. \quad (6.5 M)$$

For $n = 2m + 1$, two non equivalent representations exist. A matrix S commuting with the $n \gamma^\alpha$'s of an irreducible representation is always of the form:

$$[S, \gamma^\alpha] = 0; \quad S = \varrho \gamma^0. \quad (6.6 M)$$

Furthermore if S and S' satisfy (6.5 M), we have

$$S' = \varrho S. \quad (6.7 M)$$

*) Matrix multiplication is always a *contraction over contragredient spinor indices*. Formulas with (... M) are matrix identities.

**) \sum_P implies summation over all $\nu!$ permutations P of $\alpha_1 \dots \alpha_\nu$ with $(-1)^P = +(-)1$ for even (odd) permutations.

(3) For $n = 2m$, the 2^n elements of the ring $I = \{\gamma^r\}$ are linearly independent.

(4) The relation

$$\gamma^r \gamma^s = \varepsilon^{rs} \gamma^t; \quad t = t(r s); \quad \varepsilon^{rs} = \pm 1 \quad (6.8 M)$$

holds.

(5) If an irreducible representation of (6.2) for $n = 2m$ is found, we may form

$$\gamma_{n+1} = \varrho \gamma^{12 \dots n}, \quad \varrho = 1 \text{ or } \check{J} \quad (6.9 M)$$

and thus obtain an irreducible representation for $'n = n + 1 = 2m + 1$, because

$$(\gamma_{n+1}, \gamma^\alpha) = 0; \quad \alpha = 12 \dots 2m; \quad (\gamma_{n+1})^2 = \gamma^0. \quad (6.10 M)$$

(6) The number ξ^r in

$$(w. s.) \gamma_r \gamma^r = \xi^r \gamma^0 = \xi^{(v)} \gamma^0 \quad (6.11 M) *$$

depends but on the set $I^{(v)}$ and takes the values

$$\xi^{(v)} = (-1)^{v/2} \quad \text{for } v = 2\mu, \quad (6.12)$$

$$\xi^{(v)} = (-1)^{(v-1)/2} \quad \text{for } v = 2\mu + 1 \quad (6.13)$$

if

$$\text{signat}(g^{\alpha\beta}) = + (11 \dots 1 - 1). \quad (6.14)$$

(7) If the representation is given, to each L corresponds a transformation $S_{(L)}$

$$L \rightarrow \varrho S_{(L)}; \quad S \rightarrow L_{(L)} \quad (6.15)$$

(defined up to a factor ϱ) leaving $\gamma^{\alpha A}_B$ invariant:

$$\gamma'^{\alpha A}_B = L'^\alpha_\alpha S'^A_{(L)A} \gamma^{\alpha A}_B S^{-1B}_{(L)'B}, \quad (6.16)$$

$$\gamma'^\alpha = L'^\alpha_\alpha S_{(L)} \gamma^\alpha S_{(L)}^{-1}. \quad (6.16 M)$$

This relation in SS is perfectly analogous to (I, 3.9) in RHS; instead of the arbitrary phase factor in (I, 3.12), the arbitrary $\hat{\varrho}$ appears in (6.15).

(8) To the infinitesimal transformation

$$L'^\alpha_\alpha = \delta'^\alpha_\alpha + \frac{1}{2} \delta \omega^{\mu\nu} \Sigma_{\mu\nu}{}'^\alpha_\alpha \quad (6.17)$$

*) (w. s.) means 'without summation over indices in contragredient positions'.

corresponds

$$S_{(L)} = \gamma^0 + \frac{1}{4} \delta \omega^{\mu\nu} \gamma_{\mu\nu}. \quad (6.18 \text{ M})$$

(9) To the transformations $L = (P, T, PT)^*$ corresponds, for $n = 2m$ ($d = n - 1$)**)

$$\begin{aligned} S_P &= \varrho \gamma^n; & S_T &= \varrho \gamma^{12\dots d}; & S_{PT} &= \varrho \gamma_{n+1}; \\ S_P^{-1} &= \varrho^{-1} \gamma_n; & S_T^{-1} &= \varrho^{-1} \gamma_{d\dots 21}; & S_{PT}^{-1} &= \varrho^{-1} \gamma^{n+1}. \end{aligned} \quad (6.19 \text{ M})$$

(10) If the Dirac-equations are written as

$$(\gamma^{\alpha A}{}_B \partial_\alpha + M \gamma^{0 A}{}_B) \varphi^B(x) = (\gamma^\alpha \partial_\alpha + M \gamma^0) \varphi(x) = 0, \quad (6.20 \text{ ctr})$$

$$\chi_A(x) (\gamma^{\alpha A}{}_B \partial_\alpha - M \gamma^{0 A}{}_B) = \chi(x) (\gamma^\alpha \partial_\alpha - M \gamma^0) = 0, \quad (6.20 \text{ cov})$$

the momentum-energy-density tensor is

$$\theta^{\alpha\beta}(x) = T^{\alpha\beta}(x) + \frac{1}{2} \partial_\varrho s^{\varrho\alpha\beta}(x), \quad (6.21)$$

$$\partial_\varrho s^{\varrho\alpha\beta}(x) = (T^{\beta\alpha} - T^{\alpha\beta})(x), \quad (6.22)$$

$$T^\alpha{}_\beta(x) = \frac{1}{2} (\chi \gamma^\alpha (\cdot \partial_\beta - \partial_\beta \cdot) \varphi)(x), \quad (6.23)$$

$$\partial_\alpha T^\alpha{}_\beta(x) = 0, \quad (6.24)$$

$$s^{\alpha\beta\gamma}(x) = \frac{1}{2} (\chi \gamma^{\alpha\beta\gamma} \varphi)(x) \quad (6.25)$$

and the current-charge-density vector

$$j^\alpha(x) = (\chi \gamma^\alpha \varphi)(x), \quad (6.26)$$

$$j^\alpha(x) = j^\alpha_{(\text{convection})}(x) - \partial_\varrho m^{\varrho\alpha}(x), \quad (6.27)$$

$$j^\alpha_{(\text{convection})}(x) = - (2M)^{-1} (\chi (\cdot \partial^\alpha - \partial^\alpha \cdot) \varphi)(x), \quad (6.28)$$

$$m^{\alpha\beta}(x) = - (2M)^{-1} (\chi \gamma^{\alpha\beta} \varphi)(x). \quad (6.29)$$

In order that $\varphi^A(x)$ and $\chi_A(x)$ satisfy the wave equation (1.1), M must be a *real number* and the signature must have the form (6.14).

*) $P \rightarrow \{x^i = -x^i, x^n = x^n\}$, $T \rightarrow \{x^i = x^i, x^n = -x^n\}$.

$PT \rightarrow \{x^\alpha = -x^\alpha\}$ for $n = 2m$.

**) For $n = 2m + 1$, only $S_{(PT)}$ can be defined. As we require the full L -group, our discussion is limited to $n = 2m = \text{even}$, $d = n - 1 = \text{odd}$.

In addition to these well known theorems, we add the two following theorems which follow from a theorem of FROBENIUS and SCHUR^{11) 12)} (given in Annex):

(11) For $n = 2m$, the number of dimensions of an irreducible representation is

$$N = 2^{n/2} = 2^m. \quad (6.30)$$

(12) For

$$n = 2, 4 \pmod{8} = 2, 4; 10, 12; \dots \quad (6.31)$$

all representations are equivalent to a *real representation*. For

$$n = 6, 8 \pmod{8} = 6, 8; 14, 16; \dots \quad (6.32)$$

all representations are *necessarily complex*. However a matrix C , the *Pauli-Matrix*⁹⁾, exists, relating

$$\widehat{\gamma}^{*r} = \widehat{C} \widehat{\gamma}^r \widehat{C}^{-1} \quad \text{or} \quad \gamma^{Tr} = C \gamma^r C^{-1}. \quad (6.33)$$

(13) Every IMG (irreducible matrix group, see Annex) $\{\pm \widehat{\gamma}^r\}$ is equivalent to a unitary representation $\{\pm \widehat{\gamma}^r\}$. In RHS, this implies for every $\{\pm \widehat{\gamma}^r\}$, there exists an S

$$\gamma^r = S \gamma^r S^{-1} \quad (6.34 \text{ M})$$

so as to have

$$\gamma^{\sim Tr} \equiv \gamma^{\times r} \stackrel{*}{=} (\gamma^r)^{-1} = (w. s.) \xi^r \gamma_r. \quad (6.35 \text{ M})^*)$$

(14) For $n = 2, 4 \pmod{8}$, where $\{\widehat{\gamma}^r\}$ may be chosen real, the unitary representation may be chosen *real and orthogonal*:

$$\gamma^{\sim r} \stackrel{*}{=} (\gamma^r)^{-1} = (w. s.) \xi^r \gamma_r. \quad (6.36 \text{ M})^*)$$

) $\stackrel{}{=}$ signifies 'equal in a particular representation'.

\sim is the *transposed matrix* in SS (T being reserved for transposed operator in RHS, see I).

$$\gamma_A^{\sim r B} = \gamma^r B_A. \quad (6.37)$$

\sim interchanges the spinor-indices 'left \rightleftharpoons right', leaving their covariance unchanged!

\times is the *hermitian conjugate* matrix in CSS

$$\widehat{\gamma}_A^{\times r B} = \widehat{\gamma}^{*r B}_A. \quad (6.38)$$

To it corresponds, in a \check{J} -dependent representation, the operator relation in RHS

$$\gamma_A^{\times r B} = \gamma^{Tr B}_A. \quad (6.39)$$

§ 7. The fundamental spinors $\overset{\circ}{\eta}_{AB}$ and $\hat{\eta}_{AB}$

In order to form observables $\theta^{\alpha\beta}$, T^α_β , $s^{\alpha\beta\gamma}$ and j^α ((6.21)–(6.25)) from a contravariant field operator $\psi^A(x)$, we need a *non-singular fundamental spinor* η_{AB} from which we may form a *covariant operator* $\psi_A^T(x) = \eta_{AB} \psi^{TB}(x)$

$$\eta^{-1A}{}^C \eta_{CB} = \eta_{BC} \eta^{-1C}{}^A = \gamma^0{}^A{}_B, \quad (7.1)$$

$$\eta^{-1} \eta = \eta \eta^{-1} = \gamma^0. \quad (7.1 \text{ M})$$

We shall assume a *real representation* of $\gamma^{\alpha A}{}_B$ i.e. restrict our considerations to $n = 2, 4 \pmod{8}$. However the theory can also be written for $\overset{\circ}{J}$ -dependent $\gamma^{\alpha A}{}_B$ (thus for $n = 6, 8 \pmod{8}$), but the calculations are much longer*).

Analogous to (6.16), we require invariance of η with respect to the group $S_{(L)}$ (i.e. $c(L) = 1$ or $= \text{sig}(\det(L'^i{}_i))$ or $= \text{sig}(L'^n{}_n)$)

$$\eta_{A'B} = c(L) \eta_{AB} S_{(L)}^{-1A}{}_{A'} S_{(L)}^{-1B}{}_{B'}, \quad (7.2)$$

$$\eta = c(L) S_{(L)}^{-1} \sim \eta S_{(L)}^{-1}. \quad (7.2 \text{ M})$$

For $L_{(\text{cont})}$ (cf. (6.18)) we have

$$\gamma^{\sim\mu\nu} \eta + \eta \gamma^{\sim\mu\nu} = 0, \quad (7.3 \text{ M})$$

which allows the *two possibilities*:

$$\gamma^{\sim\alpha} = \mp \overset{\circ}{\eta} \gamma^\alpha \overset{\circ}{\eta}^{-1} \quad \text{i. e.} \quad \gamma^{\sim\alpha} \overset{\circ}{\eta} = \mp \overset{\circ}{\eta} \gamma^\alpha. \quad (7.4 \text{ M} \circ)$$

We shall now show that $\overset{\circ}{\eta}$ is pseudochronous while $\hat{\eta}$ is pseudochorous. From (7.2), with (6.19 M) ($\varrho = \lambda$), follows

$$\overset{\circ}{\eta} = c(P) \lambda^{-2} \gamma_n^{\sim} \overset{\circ}{\eta}_n \gamma_n = \mp c(P) \lambda^{-2} \overset{\circ}{\eta} (\gamma_n)^2 = \pm c(P) \lambda^{-2} \overset{\circ}{\eta}. \quad (7.5 \text{ M})$$

Thus we have $\lambda^2 = 1$ and $c(P) = + (-) 1$ for $\overset{\circ}{\eta}$ ($\hat{\eta}$). $\overset{\circ}{\eta}$ is thus orthochorous and $\hat{\eta}$ is pseudochorous. Further

$$\begin{aligned} \overset{\circ}{\eta} &= c(T) \lambda^{-2} \gamma_{d\dots 1}^{\sim} \overset{\circ}{\eta} \gamma_{d\dots 1} = \\ &= \mp c(T) \lambda^{-2} \overset{\circ}{\eta} (\gamma_d)^2 \dots (\gamma_1)^2 = \mp c(T) \lambda^{-2} \overset{\circ}{\eta} \end{aligned} \quad (7.6 \text{ M})$$

leads again to $\lambda^2 = 1$, and $c(T) = - (+) 1$ for $\overset{\circ}{\eta}$ ($\hat{\eta}$). $\overset{\circ}{\eta}$ is thus pseudochronous and $\hat{\eta}$ orthochorous.

*) The statement given in a previous communication¹³⁾, that only real representations can be used, is therefore erroneous!

Transposing (7.4), we find

$$\gamma^\alpha = \mp \overset{\smile}{\eta}^{-1} \sim \gamma^\alpha \overset{\smile}{\eta} \sim; \quad \gamma^\alpha \sim = \mp \overset{\smile}{\eta} \sim \gamma^\alpha \overset{\smile}{\eta} \sim^{-1}. \quad (7.7 \text{ M})$$

(7.4) and (7.7 M) are both changes of representation of the type (6.5 M) (because $\gamma^\alpha = -\gamma^\alpha$ is a change of representation). Thus, on account of (6.7 M), we have

$$\overset{\smile}{\eta} = \lambda \overset{\smile}{\eta}. \quad (7.8 \text{ M})$$

Transposing this equation, we find $\lambda^2 = 1$; $\overset{\smile}{\eta}$ and $\overset{\smile}{\eta}$ are either symmetric or antisymmetric matrices. Furthermore we have

$$\overset{\smile}{\eta}^{-1} \overset{\smile}{\eta} = S_{(PT)} = \pm \gamma^{12 \dots n} = \pm \gamma_{n+1}. \quad (7.9 \text{ M})$$

In order to find the symmetry of $\overset{\smile}{\eta}$, we choose an orthogonal representation (6.36):

$$\gamma^{\sim i} \equiv (\gamma^i)^{-1} = \gamma_i = \gamma^i, \quad (7.10 \text{ M})$$

$$\gamma^{\sim n} \equiv (\gamma^n)^{-1} = \gamma_n = -\gamma^n. \quad (7.11 \text{ M})$$

From the ACR's (6.2 M) follows that $\overset{\smile}{\eta} \equiv \lambda \gamma^n$ satisfies (7.4 M), $\overset{\smile}{\eta}$ is therefore an *anti-symmetric matrix* in this particular representation. Now, in analogy to (6.34 M), we have in a general representation

$$\overset{\smile}{\eta}_{A'B} = \overset{\smile}{\eta}_{AB} S^{-1A}{}_{A'} S^{-1B}{}_{B'} \quad (7.12)$$

conservation of this antisymmetry: $\overset{\smile}{\eta}_{AB}$ is, in all (real) representations, *antisymmetric*:

$$\overset{\smile}{\eta}_{AB} = \overset{\smile}{\eta}_{BA} = -\overset{\smile}{\eta}_{AB}; \quad \overset{\smile}{\eta} \sim = -\overset{\smile}{\eta}. \quad (7.13); (7.13 \text{ M})$$

We choose, in particular

$$\overset{\smile}{\eta}_{AB} \equiv \gamma_B^A \quad (7.14)$$

in order to have for the charge-density

$$\begin{aligned} j^{(1)n} &= \overset{\smile}{\psi}^T \gamma^n \psi = \overset{\smile}{\psi}_C^T \gamma^{nC}{}_B \psi^B = \psi^{TA} (-\overset{\smile}{\eta}_{AC} \gamma^{nC}{}_B) \psi^B \equiv \\ &\equiv \psi^{TA} \psi^B (-\overset{\smile}{\gamma}_{AB}^n) \equiv \sum_A \psi^{TA} \psi^A > 0; \quad -\overset{\smile}{\gamma}_{AB}^n \equiv \delta_B^A \end{aligned} \quad (7.15)$$

a *positive operator*. We may now determine the symmetries of the *pseudo-chronous covariant bi-spinors*:

$$\overset{\smile}{\gamma}_{AB}^r = \overset{\smile}{\eta}_{AC} \gamma^{rC}{}_B; \quad \overset{\smile}{\gamma}^{r(\text{cov})} = \overset{\smile}{\eta} \gamma^r. \quad (7.16); (7.16 \text{ M})$$

We have, on account of (7.4 M),

$$\overset{\circ}{\gamma}^{\alpha_1 \dots \alpha_p} = \overset{\circ}{\gamma}^{\alpha_p} \dots \overset{\circ}{\gamma}^{\alpha_1} \overset{\circ}{\eta} = - \overset{\circ}{\gamma}^{\alpha_p} \dots \overset{\circ}{\gamma}^{\alpha_1} \overset{\circ}{\eta} = - (-1)^p \overset{\circ}{\eta} \overset{\circ}{\gamma}^{\alpha_p} \dots \overset{\circ}{\gamma}^{\alpha_1}. \quad (7.17)$$

This gives the following table:

sym.:		$\overset{\circ}{\gamma}^\alpha$	$\overset{\circ}{\gamma}^{\alpha_1 \alpha_2}$			$\overset{\circ}{\gamma}^{\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5}$	(7.18)
antisym.:	$\overset{\circ}{\gamma}^0 = \overset{\circ}{\eta}$			$\overset{\circ}{\gamma}^{\alpha_1 \alpha_2 \alpha_3}$	$\overset{\circ}{\gamma}^{\alpha_1 \alpha_2 \alpha_3 \alpha_4}$...

From (7.9 M) and (7.17) follows:

$$\overset{\circ}{\eta} = \pm \overset{\circ}{\eta} \quad \text{for} \quad n = \begin{cases} 2 \pmod{8} \\ 4 \pmod{8} \end{cases}. \quad (7.19)$$

§ 8. Quantization of the Spinor Field

If $\psi^A(x)$ satisfies (6.20 ctr), it follows from the symmetries of $\overset{\circ}{\gamma}_{AB}^\alpha = \overset{\circ}{\gamma}_{(AB)}^\alpha$ and $\overset{\circ}{\gamma}_{AB}^0 = \overset{\circ}{\gamma}_{[AB]}^0$, that

$$\overset{\circ}{\psi}_A(x) = \overset{\circ}{\eta}_{AB} \overset{\circ}{\psi}^B(x) = - \overset{\circ}{\psi}^B(x) \overset{\circ}{\eta}_{BA} \quad (8.1)$$

satisfies (6.20 cov). The same is true for $\psi^{TA}(x)$ and $\psi_A^T(x)$, because we consider but real representations ($n = 2, 4 \pmod{8}$). We form the tensors $T^\alpha_\beta(x)$ and $s^{\alpha\beta\gamma}(x)$, either posing

$$\begin{aligned} \chi_A &= (\overset{\circ}{J} \overset{\circ}{\psi}_A)^T; & \varphi^A &= \psi^A: \\ T^{(1)\alpha}_\beta(x) &= \frac{1}{2} ((\overset{\circ}{J} \overset{\circ}{\psi})^T \gamma^\alpha \psi_\beta - (\overset{\circ}{J} \overset{\circ}{\psi}_\beta)^T \gamma^\alpha \psi) = \\ &= \frac{1}{2} (\psi^{TA} \overset{\circ}{J}^T \psi_\beta^B + \psi_\beta^{TB} \overset{\circ}{J} \psi^A) (x) (-\overset{\circ}{\gamma}_{AB}^\alpha) \end{aligned} \quad (8.2^{(1)})$$

or, posing

$$\begin{aligned} \chi_A &= \overset{\circ}{\psi}_A; & \psi^A &= (\overset{\circ}{J} \varphi^A)^T: \\ T^{(2)\alpha}_\beta(x) &= \frac{1}{2} (\overset{\circ}{\psi} \gamma^\alpha (\overset{\circ}{J} \psi_\beta)^T - \overset{\circ}{\psi}_\beta \gamma^\alpha (\overset{\circ}{J} \psi)^T) = \\ &= \frac{1}{2} (\psi^A \psi_\beta^{TB} \overset{\circ}{J}^T + \overset{\circ}{J} \psi_\beta^B \psi^A) (x) (-\overset{\circ}{\gamma}_{(AB)}^\alpha). \end{aligned} \quad (8.2^{(2)})$$

The second form of both equations shows a) that $T^{(1)\alpha}_\beta$ and $T^{(2)\alpha}_\beta$ are *observables* (on account of the symmetry of $\overset{\circ}{\gamma}_{(AB)}^\alpha$) and b) that they are *orthochronous*, if we define:

$$\psi'^A(x) = O_{(L)}^{-1} \psi^A(x) O_{(L)} = e^{\check{J} \lambda} S_{(L)A}^A \psi^A(L^{-1} x). \quad (8.3)^*$$

The same is true for $s^{(1)\alpha\beta\gamma}$ and $s^{(2)\alpha\beta\gamma}$, on account of the antisymmetry of $\gamma_{AB}^{\alpha\beta\gamma} = \gamma_{[A}^{\alpha\beta\gamma} \delta_{B]}$. Thus, $\theta^{(1)\alpha\beta}$ and $\theta^{(2)\alpha\beta}$ are *orthochronous observables*.

It can be shown, using the Dirac-equations and partial integrations, that one may write

$$\check{\Pi}_\mu^{(1)} = \int d\check{\sigma}_\alpha(y) ((\check{J}\check{\psi})^T \gamma^\alpha \psi_\mu)(y), \quad (8.4^{(1)})$$

$$\check{\Pi}_\mu^{(2)} = - \int d\check{\sigma}_\alpha(y) (\check{\psi}_\mu \gamma^\alpha (\check{J}\check{\psi})^T)(y). \quad (8.4^{(2)})$$

With

$$\check{\Pi}_\mu = \alpha_1 \check{\Pi}_\mu^{(1)} + \alpha_2 \check{\Pi}_\mu^{(2)},$$

the relation

$$- [\check{J} \check{\Pi}_\mu, \psi^{B'}(y')] = \psi_\mu^{B'}(y') \quad (8.5)$$

is satisfied, if

$$\begin{aligned} & -\alpha_1 [(\check{J} (\check{J}\check{\psi})^T \gamma^\alpha \psi_\mu)(y), \psi^{B'}(y')] + \alpha_2 [(\check{J}\check{\psi}_\mu \gamma^\alpha (\check{J}\check{\psi})^T)(y), \psi^{B'}(y')] \\ & \equiv \check{\delta}^\alpha(y' y) \psi_\mu^{B'}(y). \end{aligned} \quad (8.6)$$

This relation can be integrated, using the *invariant pseudochronous (real) number* (function)

$$\check{S}^{0A}_B(x y) = (-\gamma^{\alpha A}_B \partial_\alpha^x + \gamma^{0A}_B M) \check{D}^0(x y) \quad (8.7)^{**}$$

in the form

$$\begin{aligned} & -\alpha_1 [\check{J} (\check{J}\check{\psi}_A)^T(x) \gamma^{\alpha A}_B \psi^B(z), \psi^{B'}(y')] + \\ & + \alpha_2 [\check{J}\check{\psi}_A(z) \gamma^{\alpha A}_B (\check{J}\check{\psi}^B)^T(x), \psi^{B'}(y')] \equiv \check{S}^{0B'}_A(y' x) \gamma^{\alpha A}_B \psi^B(z). \end{aligned} \quad (8.8)$$

In order to verify (8.6), we operate on (8.8) with ∂_μ^z and pose $x = z = y$. Further we use (2.15)

$$\check{S}^{0B'}_A(y' y) = \gamma^{\beta B'}_A \check{\delta}_\beta(y' y) \quad (8.9)$$

and (2.12)

$$\begin{aligned} & \int d\check{\sigma}_\alpha(y) \check{\delta}_\beta(y' y) (\gamma^\beta \gamma^\alpha)^{B'}_B f^B(y) = -\text{sig}(g_{nn}) (\check{\nu}_\alpha \check{\nu}_\beta)(y') (\gamma^\beta \gamma^\alpha)^{B'}_B f^B(y') = \\ & = f^{B'}(y') = \int d\check{\sigma}_\alpha(y) \check{\delta}^\alpha(y' y) f^{B'}(y). \end{aligned} \quad (8.10)^{***}$$

*) We may, as in (2.3), introduce the arbitrary phase factor for pseudochorous and for pseudochronous transformations.

**) The function $\check{S}^A_B(xy)$ is not to be confounded with the transformation matrix $S'^A_{(L)A}$ or $S'^A_{(L)A}$.

***) $(\gamma^\beta \gamma^\alpha)^{B'}_B \check{\delta}_\beta(y' y) = \gamma^{0B'}_B \check{\delta}^\alpha(y' y).$ (8.10 a)

So far, all relations are valid for fields of the 1st and of the 2nd kind. Leaving the discussion of fields of the 2nd kind for a later publication³⁾, we rewrite (8.3) for fields of the 1st kind ($[J, \psi^A] = 0$) using (1.4):

$$\begin{aligned} & -\alpha_1 \{ \overset{\circ}{\psi}_A^T(x) \gamma^{\alpha A}{}_B [\psi^B(z), \psi^{B'}(y')]_{\mp} \pm [\overset{\circ}{\psi}_A^T(x), \psi^{B'}(y')]_{\mp} \gamma^{\alpha A}{}_B \psi^B(z) \} + \\ & + \alpha_2 \{ \overset{\circ}{\psi}_A(z) \gamma^{\alpha A}{}_B [\psi^{TB}(x), \psi^{B'}(y')]_{\mp} \pm [\overset{\circ}{\psi}_A(z), \psi^{B'}(y')]_{\mp} \gamma^{\alpha A}{}_B \psi^{TB}(x) \} = \\ & = \overset{\circ}{S}^{0B'}{}_A(y' x) \gamma^{\alpha A}{}_B \psi^B(z). \end{aligned} \quad (8.11, 1^{\text{st}} \text{ k.})$$

This is a generalised form of the CR's discussed by GREEN⁴⁾ and VOLKOV⁵⁾. The most simple solution is

$$\boxed{[\psi^A(x), \psi^B(y)]_{\mp} = 0} \quad (8.12, 1^{\text{st}} \text{ k.})$$

and

$$\mp [\overset{\circ}{\psi}_A^T(x), \psi^{B'}(y')]_{\mp} = \overset{\circ}{S}^{0B'}{}_A(y' x) \quad (8.13, 1^{\text{st}} \text{ k.})$$

which give to the 1st term the right kind of structure. In the 2nd term, we use the identity, following from (8.1) and (7.4 M)

$$\overset{\circ}{\psi}_A(z) \gamma^{\alpha A}{}_B \psi^{TB}(x) = \psi^B(z) \gamma^{\alpha A}{}_B \overset{\circ}{\psi}_A^T(x).$$

Using again (8.13) and (8.12), we find that (8.11) holds, provided

$$\boxed{\alpha_1 \mp \alpha_2 = 1.} \quad (8.14, 1^{\text{st}} \text{ k. } \mp)$$

We may rewrite (8.13) in the more usual form

$$\boxed{[\psi^A(x), \overset{\circ}{\psi}_B(y)]_{\mp} = \overset{\circ}{S}^{0A}{}_B(x y).} \quad (8.13, 1^{\text{st}} \text{ k.})$$

§ 9. The Charge Operator for Spinor Fields

The two expressions for j^α (6.26) are

$$\overset{\circ}{j}^{(1)\alpha}(x) = (\overset{\circ}{\psi}^T \gamma^\alpha \psi)(x) = (\psi^{TA} \psi^B)(x) (-\overset{\circ}{\gamma}^{\alpha}{}_{AB}), \quad (9.1^{(1)})$$

$$\overset{\circ}{j}^{(2)\alpha}(x) = (\psi \gamma^\alpha \psi^T)(x) = (\psi^A \psi^{TB})(x) (-\overset{\circ}{\gamma}^{\alpha}{}_{AB}). \quad (9.1^{(2)})$$

Forming again $Q = \beta_1 Q^{(1)} + \beta_2 Q^{(2)}$ (see (3.2)), we have, for fields of the 1st kind, analogous to (3.6, 1st k.),

$$\begin{aligned} & - [Q, \psi^{B'}(y')] = \\ & = - \int d\sigma_\alpha(y) \{ \beta_1 [(\overset{\circ}{\psi}^T \gamma^\alpha \psi)(y), \psi^{B'}(y')] + \beta_2 [(\overset{\circ}{\psi} \gamma^\alpha \psi^T)(y), \psi^{B'}(y')] \} \equiv \\ & \equiv \int d\sigma_\alpha(y) \overset{\circ}{\delta}^\alpha(y' y) \psi^{B'}(y) = \psi^{B'}(y'). \end{aligned} \quad (9.2, 1^{\text{st}} \text{ k.})$$

For fields of the 1st kind, we compare (9.2) with (8.6), (the $\overset{\circ}{J}$'s cancel out and we omit the index $_\mu$ of differentiation*). (9.2) results, if we pose again (3.7, 1st k.) i.e.

$$\beta_1 = \alpha_1; \quad \beta_2 = -\alpha_2. \quad (9.3, 1^{\text{st}} \text{ k.})$$

§ 10. Charge Conjugation for the Spinor Field (1st kind)

We see at once, rising the index of (8.13)

$$[\psi^A(x), \psi^{TB}(y)]_\mp = S^{0AB}(x y) \quad (10.1)$$

that the invariant number (function)

$$\begin{aligned} S^{0AB}(x y) &= (-\overset{\circ}{\gamma}^{\alpha AB} \partial_\alpha^x + \overset{\circ}{\gamma}^{0AB} M) \overset{\circ}{D}^0(x y) = \\ &= S^{0BA}(y x) = \overset{\circ}{\eta}^{-1BC} \overset{\circ}{S}^{0A}_C(x y) \end{aligned} \quad (10.2) **$$

is symmetric with respect to $A, x \rightleftharpoons B, y$. Therefore the substitution

$$\begin{aligned} {}'\psi^A(x) &= O_{(C)}^{-1} \psi^A(x) O_{(C)} = \psi^{TA}(x), \\ {}'\psi^{TA}(x) &= O_{(C)}^{-1} \psi^{TA}(x) O_{(C)} = \psi^A(x) \end{aligned} \quad (10.3)$$

is again an orthogonal transformation, if we choose the ACR's (8.12₊), (8.13₊), (10.1₊). Thus *FD-statistics for spinor fields is a consequence of $O_{(C)}$ -covariance or of $O_{(CT)}$ -covariance* (in perfect analogy to § 4). Furthermore we have, on account of (8.2), again the relations (4.2). Using (8.14₊), we have again, with $\alpha_1 = \alpha_2 = 1/2$, the expression (4.3): $\theta^{\alpha\beta}(x)$ is in-

*) Omitting the index $_\mu$ means, that we use the integrated equation (8.8)!

$$**) \quad \overset{\circ}{\gamma}^{rAB} = \overset{\circ}{\eta}^{-1BD} \overset{\circ}{\gamma}^{rA}_D = \overset{\circ}{\eta}^{-1AC} \overset{\circ}{\eta}^{-1BD} \overset{\circ}{\gamma}^r_{CD}, \quad (10.4)$$

$$\overset{\circ}{\gamma}^{r(\text{ctr})} = -\overset{\circ}{\gamma}^r \overset{\circ}{\eta}^{-1} = -\overset{\circ}{\eta}^{-1} \overset{\circ}{\gamma}^{r(\text{cov})} \overset{\circ}{\eta}^{-1}. \quad (10.4 \text{ M})$$

variant with respect to $O_{(C)}^*$). The equations (4.4) are also valid. Therefore, using again (3.7, 1st k.) (= (9.3, 1st k.)) we have again (4.5) and (4.6)*): j^α and Q change sign under $O_{(C)}$. The rest of the discussion is analogous to § 4.

§ 11. The Development of the Spinor Field in Terms of Positive Frequency Wave Packet Operators

We introduce again two denumerable sets of PFWP's $\{\varphi'^A, \varphi''^A, \dots \varphi^{(\varrho)A} \dots\}$ and $\{\chi'^A, \chi''^A \dots \chi^{(\varrho)A} \dots\}$ satisfying the Dirac equation (6.20) and (5.4). The sets are normalised in terms of

$$j^\alpha(\varphi', \chi')(x) = j^\alpha(\chi', \varphi')(x) = (\overset{\circ}{\varphi'} \gamma^\alpha \overset{\circ}{\chi'})(x) = (\varphi'^A \chi'^B)(x) (-\gamma_{AB}^\alpha). \quad (11.1)**$$

On account of the decomposition (6.27), one verifies that the *normalisation*

$$Q(\varphi', \chi') = 0; \quad Q(\varphi'^T, \varphi'') = Q(\varphi'', \varphi'^T) = \delta_{\varphi' \varphi''} > 0 \quad (11.2)$$

is possible. *Completeness* is assured, if the $\overset{\circ}{J}$ -dependent operators

$$\overset{\circ}{S}^{+A}_B(x y) \equiv \mathbf{S}_{\varphi'} \varphi'^A(x) \overset{\circ}{\varphi}_B'^T(y) = \overset{\circ}{S}^{+A}_B(x - y), \quad (11.3+)**$$

$$\overset{\circ}{S}^{-A}_B(x y) \equiv \mathbf{S}_{\varphi'} \varphi'^{TA}(x) \overset{\circ}{\varphi}_B'(y) = \overset{\circ}{S}^{-A}_B(x - y), \quad (11.3-)**$$

$$S^{-AB}(x y) = S^{+TAB}(x y) = S^{+BA}(y x) \quad (11.4)$$

are invariant with respect to $\{L_{(\text{ochr})}\}$. Again a PF-solution $f^{+A}(x)$ of (6.20) may be written in the form

$$f^{+A}(x) = \mathbf{S}_{\varphi'} \varphi'^A(x) f_{\varphi'}^+ = \int d\overset{\circ}{\sigma}_\alpha(y) \overset{\circ}{S}^{+A}_B(x y) \gamma^{\alpha B}_C f^{+C}(y) \quad (11.5+)$$

and

$$f^{-A}(x) = \mathbf{S}_{\chi'} \chi'^{TA}(x) f_{\chi'}^- = \int d\overset{\circ}{\sigma}_\alpha(y) \overset{\circ}{S}^{-A}_B(x y) \gamma^{\alpha B}_C f^{-C}(y). \quad (11.5-)$$

*) $\theta^{\alpha\beta}$ and $\overset{\circ}{J}^\alpha$ can be expressed in terms of *commutators*:

$$T^\alpha_\beta(x) = \frac{1}{4} \overset{\circ}{J}^{-1} ([\psi^{TA}, \psi_\beta^B] + [\psi^A, \psi_\beta^{TB}]) (x) (-\gamma_{(AB)}^\alpha), \quad (10.6)$$

$$s^{\alpha\beta\gamma}(x) = \frac{1}{4} \overset{\circ}{J}^{-1} [\psi^{TA}, \psi^B] (x) (-\gamma_{[AB]}^{\alpha\beta\gamma}). \quad (10.7)$$

$$\overset{\circ}{j}^\alpha(x) = \frac{1}{2} [\psi^{TA}, \psi^B] (x) (-\gamma_{(AB)}^\alpha). \quad (10.8)$$

**) The pseudochronous sign in $\overset{\circ}{\varphi}'_A$ and $\overset{\circ}{S}^{+A}_B$ does not imply any covariance property but only indicates lowering of the index.

The general solution of (6.20) may be written as

$$\begin{aligned}\psi^A(k) &= \mathbf{S}_{\varphi'} a_{\varphi'} \varphi'^A(x) + \mathbf{S}_{\chi'} b_{\chi'}^T \chi'^{TA}(x) = \\ &= \int \overset{\circ}{d}\sigma_\alpha(y) \overset{\circ}{S}^{0A}_B(x y) \gamma^{\alpha B}_C \psi^C(y)\end{aligned}\quad (11.6)$$

with

$$\overset{\circ}{S}^{0A}_B(x y) = (\overset{\circ}{S}^{+A}_B + \overset{\circ}{S}^{-A}_B)(x y). \quad (11.8)$$

If $x = y'$ is chosen on $\tau(y') = \tau(y) = 0$, it follows that

$$\overset{\circ}{S}^{0B'}_C(y' y) \gamma^{\alpha C}_B = \gamma^{0B'}_B \overset{\circ}{\delta}^\alpha(y' y). \quad (11.9)$$

Now, this is exactly the condition (8.9), (8.10). Thus $\overset{\circ}{S}^{0A}_B(x y)$, defined by (11.8) is identical with (8.7).

From the development (11.6) and (8.12), (8.13), follow the CR's or ACR's:

$$[a_{\varphi'}, a_{\varphi''}]_{\mp} = [b_{\chi'}^T, b_{\chi''}^T]_{\mp} = [a_{\varphi'}, b_{\chi'}^T]_{\mp} = 0, \quad (11.10_{\mp})$$

$$[a_{\varphi'}, a_{\varphi''}^T]_{\mp} = \delta_{\varphi' \varphi''}; \quad [b_{\chi'}^T, b_{\chi''}^T]_{\mp} = \delta_{\chi' \chi''}; \quad [a_{\varphi'}, b_{\chi'}^T]_{\mp} = 0. \quad (11.11_{\mp})$$

Taking the ACR's (on account of $O_{(C)}$ -covariance), (10.8) leads to

$$\mathcal{Q} = \mathbf{S}_{\varphi'} \left(N_{\varphi'} - \frac{1}{2} \right) - \mathbf{S}_{\chi'} \left(N_{\chi'} - \frac{1}{2} \right) = \mathbf{S}_{\varphi'} N_{\varphi'} - \mathbf{S}_{\chi'} N_{\chi'} \quad (11.12)$$

with $N_{\varphi'} = a_{\varphi'}^T a_{\varphi'}$. The ACR's (11.10) and (11.11) are satisfied in terms of the pseudo-quaternions (I, A-4.8)

$$\begin{aligned}a^T &= \frac{1}{2} (l + j) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}; \quad a = \frac{1}{2} (l - j) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}; \\ N &= a^T a = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{2} (1 - k)\end{aligned}\quad (11.13)$$

with

$$a_{\varphi(q)} = 1 \times (k \times k) \times (k \times k) \times \cdots \times (a \times 1) \times (1 \times 1) \times \cdots \quad (11.14)$$

$$b_{\chi(q)} = 1 \times (k \times k) \times (k \times k) \times \cdots \times (k \times a) \times (1 \times 1) \times \cdots$$

The eigenvalues are $\{N'_{\varphi'}\} = \{0, 1\}$ i.e. FD-statistics holds. The explicit form of $O_{(C)}$ is again given by (5.19), if the sets $\{\varphi'\}$ and $\{\chi'\}$ are chosen to be identical. Quantization of $\overset{\circ}{\Pi}_\mu$ leads to

$$\overset{\circ}{\Pi}^\mu = \mathbf{S}_{\varphi'} \left(N_{\varphi'} - \frac{1}{2} \right) \overset{\circ}{k}'^\mu + \mathbf{S}_{\chi'} \left(N_{\chi'} - \frac{1}{2} \right) \overset{\circ}{k}'^\mu, \quad (11.15)$$

if the PFWP's satisfy (5.20). Thus, the energy is, apart from the negative infinite 'zero point contribution' (which is again symmetric for particle and anti-particle states), a positive definite operator: the spectrum has a lower limit, and statistical thermodynamics with a positive absolute temperature may be applied¹⁰).

The CR's (11.10₋) and (11.11₋) lead, whatever α_1 and α_2 (satisfying (8.14, 1st k.₋)) we choose, to an energy spectrum without lower or upper limit: thus statistical thermodynamics cannot be applied. This is PAULI's⁹) reason for excluding BE-statistics for spinor fields.

Annex I: The Theorem of Frobenius and Schur^{11) 12)},

states that an *irreducible matrix group* (IMG) $\{\widehat{D}_i\}$ ($ik. \dots = 12 \dots h$) of order h belongs to one of the three kinds:

An IMG is of the 1st, 2nd or 3rd kind, depending on whether

$$h^{-1} \sum_i \text{tr}(\widehat{D}_i^2) = \begin{cases} -1 \\ +1 \\ 0 \end{cases} \text{ for an IMG of the } \begin{cases} 1^{\text{st}} \\ 2^{\text{nd}} \\ 3^{\text{rd}} \end{cases} \text{ kind.} \quad (\text{A-1.1})$$

(1) An IMG is of the 1st kind, if it is *equivalent to a group of real matrices*, i.e. if for all representations $\{\widehat{D}_i\}$ a matrix \widehat{S} exists, satisfying

$$\{\widehat{D}_i\} = \{\widehat{S} \widehat{D}_i \widehat{S}^{-1}\} \quad \text{with} \quad \widehat{D}_i^* = \widehat{D}_i. \quad (\text{A-1.2})$$

(2) An IMG is of the 2nd kind, if it is *equivalent to its conjugate complex group*, i.e. if for all representations a matrix \widehat{C} exists, satisfying

$$\{\widehat{D}_i\} = \{\widehat{C} \widehat{D}_i \widehat{C}^{-1}\} \quad \text{with} \quad \widehat{D}_i^* = \widehat{D}_i. \quad (\text{A-1.3})$$

NB.: For an IMG of the 1st kind, a matrix \widehat{C} exists *a fortiori* ($\widehat{C} = \widehat{S}^{*-1} \widehat{S}$).

(3) An IMG is of the 3rd kind, if it is not equivalent to its complex conjugate group i.e. *if no matrix \widehat{C} satisfying (A-1.3) exists*.

Now the set $\{\pm \Gamma\} = \{\pm \gamma^r\}$ forms, for $n = 2m$, on account of (6.8 M) an IMG of order 2×2^n . In order to find out to which kind it belongs, we have to evaluate

$$(2 \times 2^n)^{-1} \sum_r \sum_{(\pm)} \text{tr}((\pm \gamma^r)^2) = 2^{-n} \sum_r \text{tr}((\gamma^r)^2). \quad (\text{A-1.4})$$

To evaluate this sum, we decompose each subset $\Gamma^{(v)}$ into two parts

$$\{\gamma^{\alpha_1 \dots \alpha_n}\} = \{\gamma^{i_1 \dots i_n}\} + \{\gamma^{i_1 \dots i_{n-1} n}\}, \quad (\text{A-1.5})$$

each having ($d = n - 1$)

$$\binom{n}{v} = \binom{d}{v} + \binom{d}{v-1} \quad (\text{A-1.6})$$

elements. Using (6.12) and (6.13), we have

$$\begin{aligned} \sum_r \text{tr} [(\gamma^r)^2] &= \sum_v \left[\binom{d}{v} - \binom{d}{v-1} \right] \begin{Bmatrix} (-1)^{v/2} \\ (-1)^{(v-1)/2} \end{Bmatrix} \text{tr}(\gamma^0) = \\ &= N \text{Re} \sum_v \left[\binom{d}{v} - \binom{d}{v-1} \right] (1 - i) i^v = \\ &= 4 N \text{Re} [(1 + i)^{n-3}] = 2^{(n+1)/2} N \cos\left((n-3) \frac{\pi}{4}\right). \quad (\text{A-1.7}) \end{aligned}$$

We remark first, that our condition (A-1.1), (A-1.4) has the periodicity ' $n = n \pmod{8} = \dots, n-8, n, n+8, \dots$ '. In particular, we find, on account of $\cos(\pi/4) = 2^{-1/2}$, $N = 2^{n/2}$ i.e. (6.30) and furthermore that our IMG is of the 1st kind for $n = 2, 4 \pmod{8}$ (6.31) and of the 2nd kind for $n = 6, 8 \pmod{8}$ (6.32).

Bibliography

- 1) E. C. G. STUECKELBERG, *Helv. Phys. Acta* **33**, 727 (1960); to be referred to as I.
- 2) E. C. G. STUECKELBERG and M. GUENIN, *Helv. Phys. Acta* **34**, 621, (1961); to be referred to as II.
- 3) M. GUENIN and E. C. G. STUECKELBERG, *Helv. Phys. Acta* **34**, 506, (1961) (a preliminary note on fields of the 2nd kind).
- 4) H. S. GREEN, *Phys. Rev.* **90**, 270 (1953).
- 5) D. V. VOLKOV, *JETP* **9** (n° 5), 1107 (1959).
- 6) J. SCHWINGER, *Phys. Rev.* **82**, 914 (1951).
- 7) W. PAULI, *Niels Bohr and the Development of Physics*, pp. 30–31 (1955).
- 8) E. C. G. STUECKELBERG, *Helv. Phys. Acta* **14**, 322 and 588 (1941).
- 9) W. PAULI, *Ann. Inst. Poincaré* **6**, 137 (1936), *Phys. Rev.* **58**, 716 (1940), *Rev. Mod. Phys.* **13**, 203 (1941), *Progr. Theor. Phys.* **5**, 526 (1950).
- 10) See f. expl. E. C. G. STUECKELBERG, *Helv. Phys. Acta* **33**, 605 (1960).
- 11) G. FROBENIUS and I. SCHUR, *Berl. Sitz. Ber.* (1906), pp. 186–208. The theorem of F. and S. is quoted, without reference or outline of proof in:
- 12) J. S. LOMONT, *Application of Finite Groups*, Acad. Press (New York, 1959), p. 51.
- 13) E. C. G. STUECKELBERG, *Helv. Phys. Acta* **32**, 254 (1959).