

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 34 (1961)  
**Heft:** [6]: Supplementum 6. Proceedings of the International Symposium on polarization phenomena of nucleons  
  
**Artikel:** Polarization in p-He4 and p-p scattering at 22 MeV  
**Autor:** Conzett, H.E. / Igo, G. / Nir, A.  
**DOI:** <https://doi.org/10.5169/seals-513275>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Polarization in $p$ -He<sup>4</sup> and $p$ - $p$ Scattering at 22 MeV

By H. E. CONZETT, G. IGO and A. NIR<sup>1)</sup>

Lawrence Radiation Laboratory, University of California, Berkeley

We have developed a beam of 22-MeV protons, using  $p$ -He<sup>4</sup> scattering as the polarizing mechanism. We have followed the method first used by ROSEN and BROLLEY [1]<sup>2)</sup> three years ago, but before describing a few details of the method, I should like to summarize briefly the information available on  $p$ -He<sup>2</sup> scattering as a proton polarizer. This can be done most conveniently in terms of the following table.

Table

$E_{LAB}$ (MeV)	Data		Reference
	( $\theta$ )	$P(\theta)$	
0.95-18	$X$	—	K. W. BROCKMANN, Phys. Rev. 108, 1000 (1957) contains a list of authors whose work is referred to here. M. K. RUSSELL and J. H. WILLIAMS, Phys. Rev. 106, 286 (1957).
39.8	$X$	—	
1.37	—	one angle	M. J. SCOTT, Phys. Rev. 110, 1398 (1958).
2.02	—	one angle	
3.58	—	one angle	
7.8	—	one angle	J. SANADA <i>et al.</i> , J. phys. Soc. Japan, to be published.
11.4	—	one angle	
14.4	—	one angle	
10	—	10 angles	Reference 1.

Phase shifts derived from analyses of the very complete differential cross section data have been used to calculate the polarizations; and finally, description of the  $p$ - $\alpha$  interaction in terms of an effective two-body potential with a central and a spin orbit term has resulted in inter-

<sup>1)</sup> On leave from the Weizmann Institute, Rehovoth, Israel.

<sup>2)</sup> Numbers in brackets refer to References, page 258.

polarizations and predictions of the phase shifts, and thus the polarizations, up to 40 MeV [2]. It should be noted here (table) that the polarization measurements at these energies are sparse; and, since the polarizations are more sensitive than the angular distributions of the differential cross sections to uncertainties in the phase shifts, it is important to extend these measurements in order to provide further checks on theory, and thus enable experimenters to use the calculated values more confidently. The results of GAMMEL and THALER's optical-model potential calculations are shown in figure 1. I want to call your attention to the fact that between 10 and 11 MeV  $P(130^\circ) > +90\%$ , and in fact, ROSEN and BROLLEY's [1] measured value at 10 MeV is  $P(130^\circ) = +99^{+1}_{-2}\%$ . It is precisely this measurement that we use in setting up our polarized proton beam; that is, we have incident on a  $H_2$  gas target a 43.1 MeV  $\alpha$ -particle beam of about  $1 \mu A$  intensity. Protons knocked on at a laboratory angle of  $25^\circ$  are in fact equivalent to 10.8 MeV (lab. energy) protons scattered from  $He^4$  at a centre-of-mass angle of  $130^\circ$ .




Figure 1

Figure 2 shows a sketch of our experimental set up. The region of the first scattering is imbedded in paraffin to reduce the high-energy neutron background that is most troublesome. This is shown by the dotted line. After passing through the  $H_2$  gas target, the  $\alpha$ -beam passes through a shielding wall, is stopped in a Faraday Cup and monitored.

The protons knocked on at  $25^\circ$  with respect to the incident beam are collimated into the second scattering chamber. Measured left-right asymmetries in the second scattering then give us  $P_2$ , the polarization




Figure 2

that would be induced in the second scattering if the proton beam were unpolarized. Figure 3 is a schematic drawing of our scattering geometry; the  $\alpha$ -particle energy of 45.5 MeV is that just on the downstream side of the entrance foil. The following conditions apply:

1. First scattering: 8 atm. of H<sub>2</sub>,  $E_\alpha = 43.1$  MeV,  $E_p = 23$  MeV.
2. Second scattering: 6 atm. of He<sup>4</sup> or H<sub>2</sub>,  $E_p = 22$  MeV, Polarization essentially 100%, and a beam of  $\approx 6 \cdot 10^6$  protons/s.
3.  $\Delta\theta_1$ , the angle subtended at the first scattering volume by the 11/16" collimator defining the proton beam, is about  $1.4^\circ$ . This angular spread is kept small in order to minimize the inherent asymmetry in the second scattering due to non-uniform illumination across the second target.
4.  $\Delta\theta_2$ , the angular resolution of our counters, including beam width, is about  $9^\circ$ .
5. This geometry gives us second-scattered counting rates in each counter of about 3000/h in  $p$ -He<sup>4</sup> scattering at an angle of  $45^\circ$ .
6. We use a simple coincidence counting arrangement to reduce the background to the point where it constitutes no problem, at least for measurements in the forward hemisphere. A pulse from the plastic scintillator-photomultiplier tube is used to gate on the pulse-height



changed, and so the differences in counter-detection efficiencies were not cancelled out.




Figure 4




Figure 5

This raises the question of the importance of a measurement of the  $p$ - $p$  polarization at these energies. MAC GREGOR's phase-shift analyses [5] of the differential cross section data result in four possible solutions, i.e. sets of phase shifts. Two of them predict small negative polarizations in the energy and angular regions of the above results and hence, seem to be ruled out by the polarization measurements. The other two solu-

tions predict small positive polarizations and even a very precise measurement cannot distinguish between them. Hence, in closing, we would welcome any expression of theoretical opinion on this question.

#### REFERENCES

- [1] L. ROSEN and J. E. BROLLEY, Jr., Phys. Rev. *107*, 1454 (1951).
- [2] J. L. GAMMEL and R. M. THALER, Phys. Rev. *109*, 2041 (1958).
- [3] K. W. BROCKMAN's result quoted in reference 2.
- [4] W. A. BLANPIED, Phys. Rev. *116*, 738 (1959).
- [5] M. H. MACGREGOR, Phys. Rev. *113*, 1559 (1959).