

Zeitschrift: Helvetica Physica Acta
Band: 34 (1961)
Heft: [6]: Supplementum 6. Proceedings of the International Symposium on polarization phenomena of nucleons

Artikel: Observation on the reaction Li₇(p,)He₄ using polarized protons
Autor: Bearpark, K. / Hall, I. / Segel, R.E.
DOI: <https://doi.org/10.5169/seals-513282>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Observation on the Reaction $\text{Li}^7(p,\alpha)\text{He}^4$ Using Polarized Protons

By K. BEARPARK, I. HALL, R. E. SEGEL, S. M. SHAFROTH and N. W. TANNER
Clarendon Laboratory, Oxford

WOLFENSTEIN [1]¹⁾ in 1949 predicted that the reaction $\text{Li}^7(p,\alpha)\text{He}^4$ would be sensitive to proton polarization which produces an additional term in the angular distribution of the form $\sin 2\Theta \cos \Phi$. The magnitude of the coefficient associated with this term could not be predicted because of the unknown channel spin phase which could not be deduced from the unpolarized angular distribution. However it was clear that a large azimuthal asymmetry would occur at least at some energies.

A polarized proton beam was obtained by scattering 2 to 4 MeV protons from carbon at 60° (lab) at which angle the polarization had been measured [2] as -25 to -35% depending on scattering energy. The scattered beam was allowed to pass through an absorber foil and impinge on a lithium target as shown in figure 1. Alpha particles from the reaction were detected by four CsI scintillation counters at 45° (c.m. angle approximately) to the scattered beam. Using a primary beam of $2\mu\text{A}$, a carbon target 8 mg/cm^2 thick and a lithium target 0.5 mg/cm^2 thick, about 1000α -counts per hour per counter were recorded for a lithium bombarding energy of 2 MeV. The lithium bombarding energy was varied by appropriate choice of primary bombarding energy, scattering target thickness and carbon absorber thickness. The inherent asymmetry of the chamber was determined by replacing the carbon scatterer with a platinum foil of equivalent thickness.

Figure 2 shows the fractional change r (defined in the same way as WOLFENSTEIN's [1] r) in the intensity of the α -particles at $\Theta = 45^\circ$, $\Phi = 0^\circ$ due to proton polarization (100%).

The immediate consequence of the sign of the asymmetry is that the interpretation of $\text{Li}^7(p,\alpha)\text{He}^4$ in terms of compound states of $J = 0$ and $J = 2$ by INGLIS and by CRITCHFIELD and TELLER [3] is not acceptable. On the other hand two states of $J = 2$ (suggested by CHRISTY and

¹⁾ Numbers in brackets refer to References, page 338.

RUBIN [4] can give the right sign and magnitude. Three curves for the latter case calculated by WOLFENSTEIN [1] for channel spin phase γ are shown in figure 2. Apparently $\gamma \simeq 200^\circ$.




Figure 1

Geometry used for $\text{Li}^7(\mathbf{p}, \alpha)\text{He}^4$ measurement: p , primary beam; \mathbf{p} , polarized beam; S_1, S_2, S_α , collimating slits for primary and polarized beams and α -particles; c , carbon target; Li , lithium target; A , carbon absorber foil; $\alpha_1, \dots, \alpha_4$, CsI counters




Figure 2

Asymmetry r as a function of E_p showing measured points and curves calculated by WOLFENSTEIN [1]

REFERENCES

- [1] L. WOLFENSTEIN, Phys. Rev. 75, 1664 (1949).
- [2] J. E. EVANS and M. A. GRACE, Nucl. Phys. 15, 646 (1960).
- [3] D. R. INGLIS, Phys. Rev. 74, 21 (1948). — C. CRITCHFIELD and E. TELLER, Phys. Rev. 60, 10 (1941).
- [4] R. F. CHRISTY and S. RUBIN, Phys. Rev. 71, 275A (1947).