

Zeitschrift: Helvetica Physica Acta
Band: 34 (1961)
Heft: [6]: Supplementum 6. Proceedings of the International Symposium on polarization phenomena of nucleons

Artikel: Supplementary note : production and detection of a polarized deuteron beam using the atomic beam magnetic resonance method
Autor: Hughes, V.W.
DOI: <https://doi.org/10.5169/seals-541277>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Supplementary note by V. W. HUGHES

**Production and Detection of a Polarized Deuteron Beam Using
the Atomic Beam Magnetic Resonance Method**

21 March 1961

An accelerated beam of completely polarized deuterons has been obtained with the apparatus described. The anisotropy $W(0^\circ)/W(90^\circ)$ obtained from atoms in the state $(F,m) = (3/2, -3/2)$, for which the deuterons have complete polarization transverse to their direction of motion, has been measured to be:

$$\frac{W(0^\circ)}{W(90^\circ)} = 0.45 \pm 0.10$$

in good agreement with the theoretical value of 0.40. The instrumental asymmetry was measured by removing the wire stop and measuring the neutron asymmetry associated with the «flopped-in» atoms, which will include equal numbers in the two states $(F,m) = (3/2, -3/2)$ and $(3/2, -1/2)$, and noting that apart from the instrumental asymmetry the neutrons should have an isotropic angular distribution. As a further check the anisotropy $W(90^\circ)/W(0^\circ)$ obtained from atoms in the state $(F,m) = (3/2, -1/2)$ was measured and found to be:

$$\frac{W(90^\circ)}{W(0^\circ)} = 0.51 \pm 0.28$$

in good agreement with the theoretical value of 0.50.

From the above data we conclude that the accelerated beam of deuterons obtained from atoms flopped into the state $(F,m) = (3/2, -3/2)$ has a polarization of 100%, within the statistical accuracy of the measurement. The intensity of the completely polarized deuteron beam is about 2×10^{-11} ampere. The unpolarized background beam is about 2×10^{-10} ampere and consists primarily of D_2^+ ions.

The ionizer used in these experiments is an electrostatic Pierce-type gun with an ionization efficiency of about 10^{-4} . The only instrumental additions to the apparatus as described are:

- 1) Titanium-wrapped tungsten filaments for additional pumping in the ionizer region.
- 2) Helmholtz coils to produce a small magnetic field in the z -direction in the region of the ionizer and of the injector to the accelerator.