

Zeitschrift: Helvetica Physica Acta
Band: 37 (1964)
Heft: VII-VIII

Artikel: Spatial cluster decomposition properties of the S-matrix
Autor: Hepp, Klaus
DOI: <https://doi.org/10.5169/seals-113507>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Spatial Cluster Decomposition Properties of the S-Matrix

by Klaus Hepp

Seminar für theoretische Physik, Eidgenössische Technische Hochschule, Zürich*)

(11.V.64)

Although no nontrivial rigorous model of a local relativistic quantum field theory is known up to now, the general theory of quantized fields is believed to describe short-range interactions of relativistic particles. The physically reasonable properties of the Wightman framework¹⁾ supporting this are exemplified by the space-like cluster decomposition properties of the vacuum expectation values of products of field operators²⁾³⁾ and by the existence of asymptotic scattering states in strong convergence²⁾⁴⁾.

It is the purpose of this note to prove that the S-matrix in a local relativistic quantum field theory fulfills the spatial cluster decomposition properties, which have recently⁵⁾ been postulated to hold for any reasonable relativistic scattering theory. It will be demonstrated that the outcome of a scattering or production process between massive particles is asymptotically independent of the presence of other particles. This justifies an expansion of the S-matrix in cluster amplitudes⁵⁾ with some analogy to sums of connected Feynman diagrams in perturbation theory.

We consider a Wightman field theory of an at most countable family of local spinor fields $A_\mu^r(x)$. We assume that polynomials B_i^* in the smeared-out fields

$$B_i^* = \sum_{(r)(\mu)}^{\infty} \int_i \varphi_{(\mu)}^{(r)}(x_1, \dots, x_n) A_{\mu_1}^{r_1}(x_1) \dots A_{\mu_n}^{r_n}(x_n) dx_1 \dots dx_n \quad (1)$$

create from the vacuum Ω states belonging to irreducible representations $[m_i, s_i]$, $m_i > 0$, of the covering group of the inhomogeneous proper orthochronous Lorentz group:

$$B_i^* \Omega \in \mathfrak{H}_{[m_i, s_i]}, \quad B_i \Omega = 0. \quad (2)$$

For a smooth positive frequency solution $f \in \mathfrak{S}_+^{m_i}$ of the Klein Gordon equation to the mass m_i ($f_i(x) = (2\pi)^{-3/2} \int d\vec{p} \theta(\vec{p}) \delta(\vec{p}^2 - m_i^2) \hat{f}_i(\vec{p}) e^{-i\vec{p}\cdot x}$, $\hat{f}_i(\vec{p}) \in \mathfrak{S}(R^3)$) we define with $B_i(x) = U(x, 1) B_i U(x, 1)^{-1}$:

$$B_i(f_i, t) = i \int_{x^0=t} d^3x f_i^*(x) \overset{\leftrightarrow}{\partial}_0 B_i(x). \quad (3)$$

Under these assumptions HAAG⁴⁾ and RUELLE²⁾ have proved that

$$s - \lim_{t \rightarrow \pm\infty} B_1^*(f_1, t) \dots B_n^*(f_n, t) \Omega = |\bar{f}_1, \dots, \bar{f}_n^{\text{ex}}\rangle, \text{ ex} = \text{in, out}, \quad (4)$$

exists in the strong topology in \mathfrak{H} and defines an asymptotic scattering state belonging to the wave functions $\bar{f}_i = \{\bar{f}_i(\vec{p})_\alpha, \alpha = -s_i, -s_i + 1, \dots, +s_i\}$. The subspaces \mathfrak{H}^{ex} of scattering states have the structure of Fock spaces over $\{\mathfrak{H}_{[m_i, s_i]}\}$. The scattering amplitudes $(\bar{f}_1, \dots, \bar{f}_m^{\text{out}} | \bar{g}_1 \dots \bar{g}_n^{\text{in}})$ are tensor-valued tempered distributions

$$S_{mn}^{(\alpha)(\beta)}(\vec{p}_1, \dots, \vec{p}_m; -q_1, \dots, -q_n) \in \mathfrak{S}'(R^{3(m+n)}) \quad (5)$$

*) Present address: Institute for Advanced Study, Princeton.

and define under the additional assumption $\mathfrak{H}^{in} = \mathfrak{H}^{out}$ a unitary S -matrix by $(\bar{f}_1, \dots, \bar{f}_m^{out} | S | \bar{g}_1, \dots, \bar{g}_n^{out}) = (\bar{f}_1, \dots, \bar{f}_m^{out} | \bar{g}_1, \dots, \bar{g}_n^{in})$. For a 3-vector \mathbf{a} and $f \in \mathfrak{S}^+$ we set $f^a(x) = f(x^0, \mathbf{x} - \mathbf{a})$. If $B^*(f, t) \Omega = |\bar{f}\rangle$, then, due to the translation invariance of $B(x)$, one has $B^*(f^a, t) \Omega = |\bar{f}^a\rangle$ with $\bar{f}^a(\mathbf{p})_\alpha = e^{i\mathbf{p}\cdot\mathbf{a}} \bar{f}(\mathbf{p})_\alpha$. We are now prepared to prove the

Theorem 1: Under the above assumptions

$$|\langle \bar{f}_1 \dots \bar{f}_k, \bar{f}_{k+1}^a \dots \bar{f}_{k+l}^{out} | \bar{g}_1, \dots, \bar{g}_m, \bar{g}_{m+1}^a, \dots, \bar{g}_{m+n}^{in} \rangle - \langle \bar{f}_1 \dots \bar{f}_k^{out} | \bar{g}_1, \dots, \bar{g}_m^{in} \rangle \langle \bar{f}_{k+1} \dots \bar{f}_{k+l}^{out} | \bar{g}_{m+1}, \dots, \bar{g}_{m+n}^{in} \rangle| \leq c (1 + |\mathbf{a}|)^{-1/2}, \quad (6)$$

for all $\bar{f}_i(\mathbf{p})_\alpha, \bar{g}_j(\mathbf{q})_\beta \in \mathfrak{D}(R^3)$ and arbitrary integers $k, l, m, n \geq 0$, with $c < \infty$ independent of \mathbf{a} . In particular one has

$$|\langle \bar{f}_1 \dots \bar{f}_k, \bar{f}_{k+1}^a \dots \bar{f}_{k+l}^{out} | \bar{g}_1, \dots, \bar{g}_m^{in} \rangle| \leq c (1 + |\mathbf{a}|)^{-1/2}. \quad (7)$$

Proof: If $B^* \Omega \in \mathfrak{H}_{[m, s]}$, $B \Omega = 0$, then there exists²⁾ for every $\bar{f} = \{\bar{f}(\mathbf{p})_\alpha \in \mathfrak{D}(R^3), \alpha = -s, -s+1, \dots, +s\}$ a finite number of quasi-local fields $B_r(x)$, $1 \leq r \leq N$, and $f_r \in \mathfrak{S}_+^m$ such that

$$B_r^*(x) \Omega \in \mathfrak{H}_{[m, s]}, B_r(x) \Omega = 0, \sum_{r=1}^N B_r^*(f_r, t) \Omega = |\bar{f}\rangle. \quad (8)$$

Without loss of generality we assume for (6) that one has already

The C_∞ -function $|\bar{f}_i\rangle = B_i^*(f_i, t) \Omega, |\bar{g}_j\rangle = {}'B_j^*(\bar{g}_j, t) \Omega$.

$$F(s, t, \mathbf{a}) = (B_1^*(f_1, s) \dots B_{k+l}^*(f_{k+l}^a, s) \Omega, {}'B_1^*(g_1, t) \dots {}'B_{m+n}^*(g_{m+n}^a, t) \Omega) - (B_1^*(f_1, s) \dots \Omega, {}'B_1^*(g_1, t) \dots \Omega) (\dots B_{k+l}^*(f_{k+l}, s) \Omega, \dots {}'B_{m+n}^*(g_{m+n}^*, t) \Omega) \quad (9)$$

converges for $s, -t \rightarrow +\infty$ to

$$\begin{aligned} & \langle \bar{f}_1 \dots \bar{f}_k, \bar{f}_{k+1}^a \dots \bar{f}_{k+l}^{out} | \bar{g}_1, \dots, \bar{g}_m, \bar{g}_{m+1}^a, \dots, \bar{g}_{m+n}^{in} \rangle \\ & - \langle \bar{f}_1, \dots, \bar{f}_k^{out} | \bar{g}_1, \dots, \bar{g}_m^{in} \rangle \langle \bar{f}_{k+1} \dots \bar{f}_{k+l}^{out} | \bar{g}_{m+1}, \dots, \bar{g}_{m+n}^{in} \rangle. \end{aligned}$$

In order to show the boundedness of $(1 + |\mathbf{a}|)^{1/2} |\lim_{s, -t \rightarrow \infty} F(s, t, \mathbf{a})|$, we write for an arbitrary $A > 0$:

$$\begin{aligned} \lim_{s, -t \rightarrow \infty} F(s, t, \mathbf{a}) &= - \int_A^\infty ds \int_{-\infty}^{-A} dt \frac{d}{ds} \frac{d}{dt} F(s, t, \mathbf{a}) + \int_A^\infty ds \frac{d}{ds} F(s, -A, \mathbf{a}) + \\ &+ \int_{-\infty}^{-A} dt \frac{d}{dt} F(A, t, \mathbf{a}) - F(A, -A, \mathbf{a}) \equiv \sum_{i=1}^4 F_i(A, \mathbf{a}). \end{aligned} \quad (10)$$

Using the Schwarz inequality $F_1(A, \mathbf{a})$ can be majorized by

$$\int_A^\infty ds \left\| \frac{d}{ds} B_1^*(f_1, s) \dots B_{k+l}^*(f_{k+l}^a, s) \Omega \right\| \left\| \int_{-\infty}^{-A} dt \left\| \frac{d}{dt} {}'B_1^*(g_1, t) \dots {}'B_{m+n}^*(g_{m+n}^a, t) \Omega \right\| \right\| \quad (11)$$

and by four terms of the form

$$\begin{aligned} & \int_A^\infty ds \left\| \frac{d}{ds} B_1^*(f_1, s) \dots B_k^*(f_k, s) \Omega \right\| \left\| B_{k+1}^*(f_{k+1}, s) \dots B_{k+l}^*(f_{k+l}, s) \Omega \right\| \\ & \times \int_{-\infty}^{-A} dt \left\| \frac{d}{dt} {}'B_1^*(g_1, t) \dots {}'B_m^*(g_m, t) \Omega \right\| \left\| {}'B_{m+1}^*(g_{m+1}, t) \dots {}'B_{m+n}^*(g_{m+n}, t) \Omega \right\|. \end{aligned} \quad (12)$$

We expand $\| d/ds B_1^*(f_1, s) \dots B_{k+l}^*(f_{k+l}, s) \Omega \|^2$ in a sum of products of truncated vacuum expectation values (TVEV)²⁾⁴⁾, where the 1-point- and pure 2-point contributions vanish. The TVEV for $r \geq 3$ give rise to factors like:

$$\int d\mathbf{x}_1 \dots d\mathbf{x}_r \langle B_{i_1}^{(*)}(\mathbf{x}, s) \dots B_{i_r}^{(*)}(\mathbf{x}_r, s) \rangle^T h_1^{(a)}(\mathbf{x}_1, s) \dots h_r^{(a)}(\mathbf{x}_r, s) \quad (13)$$

with $h_i^{(a)} \in \mathfrak{S}_+$ possibly translated by \mathbf{a} . (13) can be majorized²⁾⁶⁾ uniformly in \mathbf{a} by

$$\begin{aligned} &\leq c \sup_{\mathbf{x}_1 (-\mathbf{a})} |h_1(\mathbf{x}_1, s)| \dots \sup_{\mathbf{x}_{r-1} (-\mathbf{a})} |h_{r-1}(\mathbf{x}_{r-1}, s)| \dots \\ &\times \int d\mathbf{x}_r |h_r(\mathbf{x}_r, -\mathbf{a}, s)| \leq c'(1 + |s|)^{-3/2(r-2)}. \end{aligned} \quad (14)$$

Since a similar estimate holds for (12), one obtains after s - and t -integration $|F_1(A, \mathbf{a})| < c_1(1 + A)^{-1}$, independently of \mathbf{a} . By expansion in TVEV one sees that terms like $\|B_1^*(f_1, A) \dots B_{k+l}^*(f_{k+l}, A) \Omega\|^2$ are uniformly bounded for all A and \mathbf{a} . Therefore $|F_i(A, \mathbf{a})| \leq c_i(1 + A)^{-1/2}$, $i = 2, 3$, holds equally, uniformly in \mathbf{a} .

In order to majorize $F_4(A, \mathbf{a})$, we first assume that all the B_i, B_j in (9) are localized in a compact region $K = \{\mathbf{x}_0^2 + \mathbf{x}^2 \leq R^2\}$, (i.e. $\text{supp } {}_i \varphi_{(\mu)}^{(*)} \subset K^{\times n}$ in (1)) and that $\text{supp } f_i|_{x_0=0}, \text{supp } g_j|_{x_0=0} \subset \{\mathbf{x}^2 \leq R^2\}$. Then $B(f^a, t)$ is localized in $\bigcup_{|\mathbf{y}| \leq R+t} \{(x_0 - t)^2 + (\mathbf{x} - \mathbf{a} - \mathbf{y})^2 \leq R^2\}$. For $A \leq |\mathbf{a}|/9$ the $B(f_i, A), B(g_j, -A)$ are localized space-like to the $B(f_{k+i}^a, A), B(g_{m+j}^a, -A)$ for sufficiently large $|\mathbf{a}|$. Then the majorization of³⁾, theorem 1, applies and gives for sufficiently large $|\mathbf{a}|$ and $A \leq |\mathbf{a}|/9$ the estimate $|F_4(A, \mathbf{a})| \leq c(A) |\mathbf{a}|^{3/2} e^{-M|\mathbf{a}|/2}$ ($M > 0$: smallest mass in the theory). The constant $c(A)$ is essentially a sum of products of norms like

$$\|{}' B_m(g_m, -A) \dots B_k^*(f_k, A) \Omega\|,$$

which for $A \leq |\mathbf{a}|/9$ are uniformly bounded by a polynomial in $|\mathbf{a}|$, due to the temperedness of the Wightman distributions.

For arbitrary B_i, B_j, f_i, g_j one uses the method of³⁾, section 5, and obtains for any integer $N \geq 0$ a constant c_4^N independent of \mathbf{a}, A such that for $A \leq |\mathbf{a}|/9$:

$$|F_4(A, \mathbf{a})| \leq c_4^N (1 + |\mathbf{a}|)^{-N}. \quad (15)$$

The theorem is proved by setting $A = |\mathbf{a}|/9$.

The theorem can be extended to wave-packets \bar{f}_i, \bar{g}_j from \mathfrak{S} , where the bounds c_i of $F_i(A, \mathbf{a})$, $1 \leq i \leq 3$, and c_4^N can be chosen to stay bounded on bounded sets of the $\{\bar{f}_i, \bar{g}_j\}$ in \mathfrak{S} . Using this one shows as in⁷⁾ that (6) entails the following distribution identity:

Corollary: For all integers $k, l, m, n \geq 0$ one has in the topology of $\mathfrak{S}'(R^{3(k+l+m+n)})$ for $\varkappa < 1/2$:

$$\begin{aligned} &\lim_{\mathbf{a} \rightarrow \infty} |1 + |\mathbf{a}||^{\varkappa}| S_{k+l, m+n}^{(\alpha, \alpha')(\beta, \beta')}(\mathbf{p}_1, \dots, \mathbf{p}_{k+l}, -\mathbf{q}_1, \dots, -\mathbf{q}_{m+n}) e^{i\mathbf{a} \left\{ \sum_{i=1}^k \mathbf{p}_i - \sum_{j=1}^m \mathbf{q}_j \right\}} \\ &- S_{k, m}^{(\alpha)(\beta)}(\mathbf{p}_1, \dots, \mathbf{p}_k, -\mathbf{q}_1 \dots, -\mathbf{q}_m) S_{l, n}^{(\alpha')(\beta')(\mathbf{p}, \mathbf{q})}(\mathbf{p}_{k+1}, \dots, \mathbf{p}_{k+l}, -\mathbf{q}_{m+1}, \dots, -\mathbf{q}_{m+n})| = 0. \end{aligned} \quad (16)$$

The theorem can be trivially extended to the decomposition into several clusters along a space-like plane. Furthermore the limit (6) relies only in a rudimentary way on the covariance of the theory under the homogeneous Lorentz group and on the microscopic locality of the fields. Essential for the proof was the existence of quasi-local

field operators, which create eigenstates of the mass operator from the vacuum and whose TVEV decrease sufficiently strongly for large space-like separations of the arguments. Therefore the spatial cluster decomposition property of the S-matrix holds as well in the Araki-Haag framework of (macroscopically) localized observables⁶⁾.

The approach of the limit as 0 ($|\mathbf{a}|^{-1/2}$) is governed by the decay of the wave-packets f_i, g_j . For non-overlapping wave-packets in velocity-space:

$$\left\{ \frac{\mathbf{p}}{\sqrt{\mathbf{p}^2 + m_i^2}} : \mathbf{p} \in \text{supp } \hat{f}_i \right\} \cap \left\{ \frac{\mathbf{p}}{\sqrt{\mathbf{p}^2 + m_j^2}} : \mathbf{p} \in \text{supp } \hat{f}_j \right\} = \emptyset, \quad i \neq j, \quad (17)$$

the limit (4) is attained faster than any power of $(1 + |t|)^{-1}$ (see also H. ARAKI, R. HAAG, and D. W. ROBINSON, to be published). Then the scattering amplitudes show stronger cluster decomposition properties:

Theorem 2: For non-overlapping $\{f_i\}$ and, separately, $\{g_j\}$ (5) can be majorized by $c_N(1 + |\mathbf{a}|)^{-N}$ ($c_N < \infty$ for all N) in all directions \mathbf{e} satisfying for all $0 \leq a < \infty$:

$$\mathbf{a} = a \mathbf{e} \neq \frac{\mathbf{p}_i}{\sqrt{\mathbf{p}_i^2 + m_i^2}} - \frac{\mathbf{p}_j}{\sqrt{\mathbf{p}_j^2 + m_j^2}} \quad (18)$$

for $\mathbf{p}_i \in \text{supp } \hat{f}_i$ ($1 \leq i \leq k$), $\mathbf{p}_j \in \text{supp } \hat{f}_j$ ($k+1 \leq j \leq k+l$),

or $\mathbf{p}_i \in \text{supp } \hat{g}_i$ ($1 \leq i \leq m$), $\mathbf{p}_j \in \text{supp } \hat{g}_j$ ($m+1 \leq j \leq m+n$).

Proof: Using the notations of theorem 1 we shall show under (18) for \mathbf{e} that $|F_i(a \mathbf{e}, A)|$, $1 \leq i \leq 3$, can be majorized by $c_N^i(1 + A)^{-N}$, with $c_N^i < \infty$ for all N uniformly in a . Then theorem 2 follows by setting $A = a/9$.

Let $\varepsilon > 0$ be so small that the ε -neighbourhoods of the supports of $\{\hat{f}_i\}$ and, separately, of $\{\hat{g}_j\}$ still fulfill (17) and (18). Let $S(h, s) \equiv \{ \mathbf{x} = (\mathbf{p}/\sqrt{\mathbf{p}^2 + m^2}) t, \mathbf{p} \in U_\varepsilon(\text{supp } \hat{h}) \}$. Then the integral (13) over the region $S(h_1, s) \times \dots \times S(h_r, s)$ is strongly decreasing in s uniformly in a , due to the strong decrease of the supremum of $|\langle B_{i_1}^{(*)}(\mathbf{x}_1, s) \dots B_{i_r}^{(*)}(\mathbf{x}_r, s) \rangle^T|$ in this region. For, a purely geometric argument shows that because of (18) the infimum of $|\mathbf{x}_i(-\mathbf{a}) - \mathbf{x}_{i+1}(-\mathbf{a})|$ for $\mathbf{x}_k \in S(h_k, s)$ increases linearly in s uniformly in a , at least for one $1 \leq i \leq r-1$. On the other hand the strong decrease of $|h_k(\mathbf{x}_k, s)|$ in s uniformly for $\mathbf{x}_k \in R^3 - S(h_k, s)^2$ guarantees a similar majorization of (13) integrated over $R^3 - S(h_k, s)$.

Again, the results can be generalized to more complicated configurations along a space-like plane. No non-trivial temporal cluster decomposition properties of the S-matrix have been obtained.

In conclusion I wish to thank Professor R. JOST for stimulating discussions.

Bibliography

- ¹⁾ A. S. WIGHTMAN, Phys. Rev. 101, 860 (1956).
- ²⁾ D. RUELLE, Helv. Phys. Acta 35, 147 (1962).
- ³⁾ H. ARAKI, K. HEPP and D. RUELLE, Helv. Phys. Acta 35, 164 (1962).
- ⁴⁾ R. HAAG, Phys. Rev. 112, 669 (1958).
- ⁵⁾ E. H. WICHMANN and J. H. CRICHTON, Phys. Rev. 132, 2788 (1963).
- ⁶⁾ H. ARAKI, *Vorlesungen über axiomatische Quantenfeldtheorie* (Zürich 1961/62).
- ⁷⁾ R. JOST and K. HEPP, Helv. Phys. Acta 35, 34 (1962).