Zeitschrift:	Helvetica Physica Acta
Band:	38 (1965)
Heft:	I
Artikel:	Acoustic scattering from an inhomogeneity of the medium
Autor:	Morse, Philip M.
DOI:	https://doi.org/10.5169/seals-113573

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Acoustic Scattering from an Inhomogeneity of the Medium\*)

## by Philip M. Morse

Massachusetts Institute of Technology, Cambridge, Massachusetts

## (27. VIII. 64)

Abstract. The wave equation is written in its dimensionless form, which makes it possible to compute separately the effects of an inhomogeneity of density or of compressibility in a small region, on the pressure and velocity of the region where irradiated by an incident plane wave. Specific formulas are given for these quantities, and for the resulting scattered wave from a sphere of radius  $a \ll \lambda$ , when the compressibility and density inside the sphere differ by factors of any size from the values outside.

In discussing the scattering of sound waves by a region in the medium, in which the density and compressibility have different values than in the rest of the medium, it is usual to mingle the two effects by writing the wave equation as

$$\Delta^2 \phi + \omega^2 \rho \varkappa \phi = 0 \tag{1}$$

where  $(\omega/2\pi)$  is the frequency,  $\varrho$  the density and  $\varkappa$  the compressibility of the medium. If, within region R,  $\varrho = \varrho_s$  and  $\varkappa = \varkappa_s$ , different from the constant  $\varrho$  and  $\varkappa$  of the rest of the medium, then the difference  $\omega^2 (\varrho_s \varkappa_s - \varrho \varkappa)$  can be taken to the right-hand side of the equation and treated as a scattering term.

However, it is more instructive to set up the equation in dimensionless form by starting from the two basic equations relating fluid velocity with pressure

$$f((\partial \boldsymbol{u}/\partial t) = -(1/\varrho) \operatorname{grad} p; \varkappa (\partial p/\partial t) = -\operatorname{div} \boldsymbol{u}$$

and writing

$$\frac{1}{\varrho \, \omega^2} \, \varDelta^2 \, \rho + \varkappa \, \rho = \begin{cases} 0 \text{ outside } R \\ (\varkappa - \varkappa_s) \, \rho + \frac{1}{\omega^2} \, \operatorname{div} \left[ \left( \frac{1}{\varrho} - \frac{1}{\varrho_s} \right) \operatorname{grad} \rho \right] \text{ inside } R \,. \end{cases}$$

$$(2)$$

Now multiplication by  $\rho \omega^2 = (k^2/\varkappa)$  produces

$$\Delta^{2} p + k^{2} p = \begin{cases} 0 \text{ outside } R \\ -\delta_{k} p + \operatorname{div} [\delta_{p} \operatorname{grad} p] \text{ inside } R, \end{cases}$$
(3)

\*) This note is in memory of pleasant and informative discussions on wave scattering with Professor E. C. G. STUECKELBERG, in Princeton and in Munich, some 35 years ago.

Philipp M. Morse

where

$$\delta_k = \frac{\varkappa_3 - \varkappa}{\varkappa}$$
 and  $\delta_p = \frac{\varrho_s - \varrho}{\varrho_s} \cdot$ 

This equation can now be solved so that the effects of change of compressibility are distinguishable from the effects of change of density.

For example, the integral equation for the scattering of sound from region R, in unbounded space, is

$$p(\mathbf{r}) = A \ e^{i\mathbf{k}_i \cdot \mathbf{r}} + \int_R \left[k^2 \ \delta_k \ p(\mathbf{r}_0) \ g(\mathbf{r} \mid \mathbf{r}_0) + \delta_\varrho \ (\operatorname{grad}_0 \ p) \ . \ (\operatorname{grad}_0 \ g)\right] \ dv_0 \ , \tag{4}$$

where g is the usual Green's function  $[e^{ik|\mathbf{r}-\mathbf{r}_0|}/4 \pi | \mathbf{r}-\mathbf{r}_0|]$ . At large distances from region R the scattered wave becomes A  $(e^{ikr}/r) F_s (\mathbf{k}_i \cdot \mathbf{k}_s)$ 

$$F_{s} = \frac{1}{4\pi} \int_{R} \left[ k^{2} \,\delta_{k} \, p(\boldsymbol{r}_{0}) + k_{p} \, c \, \boldsymbol{k}_{s} \cdot \boldsymbol{u}(\boldsymbol{r}_{0}) \right] \, e^{-i\boldsymbol{k}_{s} \cdot \boldsymbol{r}_{0}} \, dv_{0} \,, \tag{5}$$

where  $\mathbf{k}_s$  is the vector with magnitude k in the direction of scattering. Thus at long wave-lengths the first term in the integral is a spherically symmetric wave produced by the difference in compressibility of the medium in R, from that outside R, whereas the second term represents the dipole radiation caused by the fact that a region of differing density moves at a different rate than does the surrounding medium. In this form of the solution the two effects are clearly distinguished.

This formulation can be extended to include the scattering effect of turbulence, when the fluid velocity within R is U + u, u being the velocity of the sound wave and U the velocity of turbulent motion. The additional term entering on the right-hand side of (3) is then  $-2 \Sigma \left[ \frac{\partial^2}{\partial x_m} \left( \frac{\partial u_m}{\partial x_m} \right) \right]$ , which adds another term

$$- (\varrho/2\pi) \int [\boldsymbol{k}_s \cdot \boldsymbol{U}_{\omega}(\boldsymbol{r}_0)] [\boldsymbol{k}_s \cdot \boldsymbol{u}(\boldsymbol{r}_0)] e^{-i\boldsymbol{k}_s \cdot \boldsymbol{r}_0} dv_0$$
(6)

to the integral of (5), where  $U_w$  is the Fourier transform of U for frequency  $(\omega/2\pi)$ . This represents quadrupole radiation from the turbulent region.

If the differential quantities  $\delta_{\varkappa}$  and  $\delta_{\varrho}$  are small (and U = 0) then at long wavelengths the Born approximation indicates that the monopole and dipole strengths of the scattered wave are

$$S_{s} \simeq \frac{i \ k \ \langle p \rangle}{\varrho \ c} \int_{R} \delta_{\varkappa} \ dv; \ \boldsymbol{D}_{s} \simeq \langle \boldsymbol{u} \rangle \int_{R} \delta_{p} \ dv \ , \tag{7}$$

where  $\langle p \rangle$  and  $\langle u \rangle$  are the mean acoustic pressure and velocity, caused by the incident wave, near region R. If the  $\delta$ 's are not small, but region R is a sphere of radius a, small compared to the wavelength, then separation into spherical harmonic components demonstrates that the scattering strengths are

$$S_{s} \simeq \frac{i \ k \ \langle p \rangle}{\varrho \ c} \int \left( \frac{\varkappa_{s} - \varkappa}{\varkappa} \right) dv; \ \boldsymbol{D}_{s} \simeq \langle \boldsymbol{u} \rangle_{R} \int \left( \frac{3 \ \varrho - 3 \ \varrho_{s}}{2 \ \varrho_{s} + \varrho} \right) dv . \tag{8}$$

In this case, for an incident plane wave in the z direction, of amplitude A, the pressure

54

Vol. 38, 1965

and radial velocity at the surface of the sphere, the scattered pressure at large distances and the scattering and absorption cross sections, at long wavelengths, are:

$$\begin{aligned}
\phi(a,\vartheta) \simeq A \left[ 1 + i \, k \, a \, \frac{3 \, \varrho_s}{2 \, \varrho_s + \varrho} \cos\vartheta \right] & u_r\left(a,\vartheta\right) \simeq \left(A/\varrho \, c\right) \left[ \frac{1}{3} \, i \, k \, a \, \frac{\varkappa_s}{\varkappa} + \frac{3 \, \varrho}{2 \, \varrho_s + \varrho} \cos\vartheta \right] \\
& \phi_s\left(r,\vartheta\right) \to A \left(e^{i \, k \, r}/r\right) \, F_s(\vartheta) \\
& F_s(\vartheta) \simeq \frac{1}{3} \, k^2 \, a^3 \left[ \left(\frac{\varkappa_s - \varkappa}{\varkappa}\right) + \frac{1}{3} \left(\frac{3 \, \varrho_s - 3 \, \varrho}{2 \, \varrho_s + \varrho}\right) \cos\vartheta \right] \\
& Q_s \simeq \frac{4}{\varrho} \, \pi \, a^2 \left(k \, a\right)^4 \left[ \left| \frac{\varkappa_s - \varkappa}{\varkappa} \right|^2 + \frac{1}{3} \left| \frac{3 \, \varrho_s - 3 \, \varrho}{2 \, \varrho_s + \varrho} \right|^2 \right] \\
& Q_a \simeq \frac{4}{3} \, \pi \, a^2 \left(k \, a\right) \, I \, m \left[ \frac{\varkappa_s - \varkappa}{\varkappa} + \frac{3 \, \varrho_s - 3 \, \varrho}{2 \, \varrho_s + \varrho} \right]
\end{aligned}$$

where we have assumed that  $\varkappa_s$  and/or  $\varrho_s$  may be complex quantities, representing energy loss.

For a heavy, incompressible sphere  $(\varkappa_s \ll \varkappa, \varrho_s \gg \varrho)$  the angle-distribution of the scattered intensity has the factor  $(-1 + 3/2 \cos \vartheta)^2$ , with some forward scattering, though with more backward scattering. (This is in contrast to the Born approximation, which would predict no scattering at  $\vartheta = 0$ .) In this case the radial velocity of the surface is zero and the pressure there is  $A (1 + 3/2 i k a \cos \vartheta)$ . For a light, compressible sphere  $(\varkappa_s \gg \varkappa, \varrho_s \ll \varrho)$  the scattered intensity has the factor  $[(\varkappa_s/\varkappa) - 3 \cos \vartheta]^2$ , again differing from the Born approximation. The pressure at r = a is A, the pressure of the incident wave, and the radial velocity at r = a is  $(A/\varrho c) [i k a (\varkappa_s/\varkappa) + 3 \cos \vartheta]$  which can be quite large. The sphere moves back and forth along the z axis with three times the velocity of the surrounding fluid.

Thus separation of the effects of compressibility from the effects of density, by treating the wave equation in its dimensionless form, enables one to see more clearly the separate effects, on the motion and pressure of region R and on the resulting scattered wave, of the two kinds of inhomogeneities.