**Zeitschrift:** Helvetica Physica Acta

**Band:** 38 (1965)

Heft: VI

Artikel: Possibility for a large spin-flip part in the scattering amplitudes of K^-p

charge exchange at 9,5 GeV/e

**Autor:** Fischer, Walter E.

**DOI:** https://doi.org/10.5169/seals-113614

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

## Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Possibility for a Large Spin-Flip Part in the Scattering Amplitudes of K-p Charge Exchange at 9,5 GeV/c

by Walter E. Fischer\*)

CERN-Geneva

(12.6.65)

An experiment on  $K^- p$  charge exchange, carried out recently at CERN¹), shows a momentum transfer distribution in which the cross section bends downwards towards t=0, while the maximum is situated at t=-0.1. The same behaviour appears also in  $\pi^- p$  charge exchange²). This is incompatible with pure exponential behaviour of the elastic scattering at small t. In this letter, which must be considered as an additional remark to the already published paper of the CERN/ETH group¹), we try to explain this somewhat strange behaviour of the momentum transfer distribution by a strong spin-flip part in the scattering amplitude.

Consider now the reaction

$$K^- + p \rightarrow \overline{K}^0 + n$$
.

Convenient variables for the description of this system are the squares of the centre-of-mass total energies in the three channels s, u, and t, with

$$s + u + t = 2 M^2 + 2 m^2$$
.

We denote the nucleon and K mass by M and m, respectively. The differential cross section in the c.m. is given by:

$$\frac{d\sigma}{dt} = \frac{1}{\pi} \left( \frac{M}{4 W k} \right)^2 \sum \mid \overline{u}_2 \ T \ u_1 \mid^2$$

where the sum is taken over final spin states and the average over the initial spin states and

$$T = -A(s,t) + \frac{i}{2} \gamma_{\mu} (q_1 + q_2)_{\mu} B(s,t)$$

k is the c.m. momentum and W the c.m. energy.  $q_1$  and  $q_2$  are the c.m.four-momenta of the  $K^-$  and the  $\overline{K}{}^0$ , respectively.

Omitting the modification due to Coulomb interaction, the barycentric kaonnucleon differential cross section is also given by:

$$\frac{d\sigma}{dt} = \frac{\pi}{k^2} \left| f_1(k\cos\theta) + \cos\theta f_2(k\cos\theta) \right|^2 + \sin^2\theta \left| f_2(k\cos\theta) \right|^2$$

<sup>\*)</sup> Member of the ETH-visiting group.

where the first term of the right-hand side is the non-spin-flip part and the second the spin-flip part of the amplitude.  $\vartheta$  is the scattering angle in the barycentric system.

In terms of helicity state amplitudes

$$f_{\lambda\lambda'} = -\frac{M}{4\pi W} \frac{1}{u} \left(-p_2, \lambda\right) \left(-A + \frac{i}{2} \gamma_{\mu} \left(q_1 + q_2\right)_{\mu} B\right) u \left(p_1, \lambda'\right)$$

one gets the differential cross section

$$\frac{d\sigma}{dt} = \frac{\pi}{k^2} \left[ \left( 1 - \frac{t}{4 M^2} \right) |f_{1/2 1/2}|^2 - \frac{t (t + 4 q^2) s}{4 (4 M^2 - t^2) M^2} |f_{1/2 - 1/2}|^2 \right].$$


Fitting this formula to the experimental cross section and assuming that

$$|f_{1/21/2}|^2 = a e^{\alpha t}$$
 and  $|f_{1/2-1/2}|^2 = b e^{\alpha t}$ 

we get for the three parameters


$$a=229\,rac{\mu b}{
m ster}$$
 ,  $b=114\,rac{\mu b}{
m ster}$  ,  $lpha=7.4\;(G\,e\,V/c)^{-2}$  .

Of course these parameters have a meaning only for momentum transfer |t| < 0.8. For higher values the parametrization no longer fits the experimental distribution, as one can see in Figure 1. These results were published before<sup>1</sup>) in order to explain the maximum in the experimental differential cross section at small momentum transfer.



Experimental momentum transfer distribution and parametrization fit

It is now interesting to see what this parametrisation means in terms of spin-flip and non spin-flip amplitudes. The ratio of spin-flip to non spin-flip terms as a function of the momentum transfer can be seen in Figure 2.



Spin-flip to non-spin-flip ratio for the parametrization and for a  $N \varrho N$  coupling model

We compare now this result with the peripheral model. Assuming that this charge-exchange reaction is dominated by a  $\varrho$ -exchange, we define the  $N \varrho N$ -coupling with the Lagrangian

$$\mathcal{L} = rac{i}{2} \gamma_{N \varrho N} (ar{N} \gamma_{\mu} N) \varrho^{\mu} + rac{\mu_{N \varrho N}}{8 M} (ar{N} \sigma_{\mu \nu} N) (\partial_{\mu} \varrho^{\nu} - \partial_{\nu} \varrho^{\mu})$$

where  $\gamma_{N\varrho N}$  is the  $N\varrho$  N-coupling constant and  $\mu_{N\varrho N}$  the difference of the anomalous magnetic moments of the nucleons. If we believe in a  $\varrho$ -photon analogy, we take

$$\frac{\mu_{N\varrho N}}{\gamma_{N\varrho N}} = 3.7$$

which is just the ratio of the tensor to the vector coupling in the  $N \varrho$  N-coupling. The pseudoscalars  $K^-$  and  $\overline{K}{}^0$  are coupled to the  $\varrho$ -meson by

$$L = \gamma_{K \varrho K} \left( \phi \ \partial_{\mu} \phi^* - \phi^* \ \partial_{\mu} \phi \right) \varrho^{\mu}$$

 $\phi$  and  $\varrho^{\mu}$  are the pseudoscalar- and vector-field operators respectively.  $\gamma_{K\varrho K}$  is the coupling constant of the  $\varrho$ -meson to the K-mesons. With this model we get the following matrix T,

$$T = \gamma_{K\varrho K} \frac{1}{m_{\varrho}^2 - t} \overline{u_2} (-p_2) \left\{ \frac{u - s}{4M} \mu_{N\varrho N} + \frac{i}{2} \gamma_{\mu} (q_1 + q_2)_{\mu} (\gamma_{N\varrho N} + \mu_{N\varrho N}) \right\} u_1(p_1)$$

 $\overline{u_2}$  and  $u_1$  are the Dirac-spinors of the neutron and the proton.  $1/(m_{\varrho}^2 - t)$  of course comes from the  $\varrho$ -meson propagator.

It is now an easy thing to calculate the spin-flip to non spin-flip ratio for this matrix. Since we are only interested in this ratio, we are not bothered by the somewhat unknown  $K \varrho$  K-coupling constant. This ratio is also plotted in Figure 2. One sees that the relatively strong spin-flip term in this coupling is still smaller than demanded by the parametrisation. Therefore it is hopeless to try to reproduce the momentum transfer distribution with a first order perturbation model of the  $\varrho$ -exchange type.

Apart from this, there also arises the difficulty that the  $\varrho$ -meson propagator is too flat if one does not introduce a form factor which is strongly t dependent. One also sees that a change of the ratio  $C_T/C_V$  does not improve the situation. Even if we go over to the model of Gottfried and Jackson³) and take into account the absorption effects in the incoming and outgoing channels, the model is still not able to reproduce the experimental momentum transfer distribution, although the spin-flip to non spin-flip ratio is shifted in the right direction⁴). Although there is a spin-flip contribution in the  $N \varrho N$ -coupling the recoil neutron is not polarized. (Both, the electric and magnetic part of the coupling are real.) In our parametrisation, however, the relative phase of the amplitudes a and b can be such that, due to the large spin-flip term, a polarisation of the recoil neutron is required. This could be checked by an experiment on polarized target⁵).

I am grateful to Dr. D. Amati and Dr. H. Högaasen for helpful comments. I should like to thank also my colleagues of the CERN/ETH group for stimulating discussions. I also wish to express my appreciation to Professor P. Preiswerk for the hospitality extended to me at CERN.

#### References

- 1) P. Astbury, G. Finocchiaro, A. Michelini, C. Verkerk, D. Websdale, C. H. West, W. Beusch, B. Gobbi, M. Pepin, E. Polgar, M. Pouchon, Submitted to Physics Letters.
- <sup>2</sup>) P. Borgeaud, C. Bruneton, P. Falk-Vairant, O. Guisan, J. Kirz, P. Sonderegger, A. V. Stirling, B. Amblard, C. Caverzasio, J. P. Guillaud, M. Yvert, Phys. Rev. Letters 14, 763 (1965).
- 3) K. Gottfrfried and J. D. Jackson, Nuovo Cimento 34, 735 (1964).
- 4) T. H. HÖGAASEN, private communication.
- 5) An experiment of this kind for  $\pi p$ -charge exchange is now prepared at CERN, by the Saclay-Orsay group (see Ref. 2).