

**Zeitschrift:** Helvetica Physica Acta

**Band:** 43 (1970)

**Heft:** 3

**Rubrik:** Zusammenfassungen der letzten eingegangenen Arbeiten = Résumés des derniers articles reçus

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# HELVETICA PHYSICA ACTA

Zusammenfassungen der letzten eingegangenen Arbeiten

Résumés des derniers articles reçus

## Nuclear Magnetic Resonance in $\text{Be}_{22}\text{Re}$ and $\text{Be}_{22}\text{Tc}$

by M. BERNASSON, P. DESCOUTS and G. A. STYLES

Institut der Physique de la Matière Condensée, Université de Genève

(31. I. 70)

*Abstract.* The N. M. R. Knight shifts,  $K$ , and line-widths,  $\Delta H$ , of the  $^9\text{Be}$ ,  $^{185}\text{Re}$ ,  $^{187}\text{Re}$  and  $^{99}\text{Tc}$  resonances have been measured in the intermetallic compounds  $\text{Be}_{22}\text{Re}$  and  $\text{Be}_{22}\text{Tc}$  at three different magnetic field strengths. Both  $K$  and  $\Delta H$  were determined at  $300^\circ\text{K}$ ,  $77^\circ\text{K}$  and  $4.2^\circ\text{K}$  and were found to be independent of temperature. The line-widths of the  $^9\text{Be}$  resonance in both compounds and of the  $^{185}\text{Re}$  and  $^{187}\text{Re}$  resonances in  $\text{Be}_{22}\text{Re}$  exhibit a field-dependence which is shown to be due to second-order quadrupole effects. The values of the Knight shift of  $^9\text{Be}$  ( $K = -0.0027\%$ ), close to that measured in pure Be, together with those of  $^{185}\text{Re}$  ( $K = -0.880\%$ ) and of  $^{99}\text{Tc}$  ( $K = -0.536\%$ ) appear to indicate localisation of the d-electron wavefunctions near to the transition metal ions.

## On the Asymptotic Condition of Scattering Theory

by W. O. AMREIN, PH. A. MARTIN and B. MISRA

Institute of Theoretical Physics, University of Geneva, Geneva, Switzerland

(6. II. 70)

*Abstract.* We propose a new formulation of the asymptotic condition of scattering theory which applies to Coulomb interactions and other long range potentials and which generalizes the usual asymptotic condition. It consists of the requirement that the constants of the free motion be asymptotically stationary also under the real evolution. From this and two supplementary conditions we prove the existence of wave operators and of a scattering operator. It is then shown that the wave operators are strong operator limits as in standard scattering theory except that the free evolution may have to be replaced by a modified propagator. Finally, Dollard's method of proving the asymptotic convergence for the Coulomb interaction is extended to a more general class of potentials.

## The Decay of $^{51}\text{Cr}$

by CL. RIBORDY and O. HUBER

Institut de Physique de l'Université de Fribourg (Suisse)

(16. II. 70)

*Abstract.* The decay of  $^{51}\text{Cr}$  is studied with a  $\text{Ge}(\text{Li})$  detector of  $3.2\text{ cm}^3$  and a double focusing beta spectrometer. Hypothetical transitions (150 keV, 325 keV, 470 keV and 645 keV) in  $^{51}\text{V}$  are excluded. The energy and K-internal conversion coefficient of the single  $\gamma$ -transition in  $^{51}\text{V}$  are measured:  $E_\gamma = (320.032 \pm 0.040)\text{ keV}$  and  $\alpha_K = (1.46 \pm 0.13) \cdot 10^{-3}$ . The  $Q$  value of the decay  $^{51}\text{Cr} \rightarrow ^{51}\text{V}$  is obtained from the end point energy of the internal bremsstrahlung (IB) spectrum:  $Q = (748 \pm 14)\text{ keV}$ . This IB spectrum is measured with a  $\text{Ge}(\text{Li})$  detector and unfolded with a described procedure. Although a pile up rejector was used, this  $\text{Ge}(\text{Li})$  measurement suffers still from a distortion due to the pile up effect. The IB spectrum taken with the magnetic spectrometer (Pile up free) reveals a doubtful structure which remains unclear. The proportion of the electron captures populating the first excited state of  $^{51}\text{V}$  is measured:  $\beta = (10.2 \pm 1\%)$ . A decay scheme is proposed.

### Nucleon Transfer Reactions below the Coulombbarrier

by D. TRAUTMANN and K. ALDER

Institute of Theoretical Physics, Basel, Switzerland

(23. II. 70)

*Abstract.* A DWBA treatment for nucleon transfer reactions is given which is applicable for energies below and in the neighbourhood of the Coulomb barrier. At energies well below the Coulomb barrier the theory is essentially exact and the radial integrals describing the reaction may be evaluated analytically by means of generalized hypergeometric series. Methods for their numerical calculation are given. Semiquantal and semiclassical approximations are considered. From those it is possible to recognize a close similarity between transfer processes and Coulomb excitation. The effects of nuclear interaction, which become important at energies close to the Coulomb barrier, are treated in an approximate manner. Expressions are given for the differential and total cross-section. The deuteron stripping reaction is treated as a special case and the polarisation of the outgoing proton in  $(d, p)$ -reactions is calculated. The connection between this treatment and the diffraction model developed by Dar and Frahn and Sharaf is discussed. An improvement of the model is given. A method similar to the one used in scattering and Coulomb excitation calculations is used to improve the slow convergence of the sum over the orbital angular momenta. Comparisons with actual experiments are discussed.

### Four-Particle-Two-Hole Core-Excitation in Heavy Nuclei

by U. GÖTZ, J. HADERMANN, K. ALDER

Seminar für theoretische Physik der Universität Basel, Basel, Switzerland

(23. II. 70)

*Abstract.* Four-particle-two-hole core-excitation in heavy nuclei with two nucleons outside the closed shells is investigated by including the highest core orbits into the configurations space of a shell-model calculation. The core-excitation is supposed to be effected by an interaction between core nucleons and valence nucleons which is essentially the same as the well-known residual interaction between the valence nucleons. The nucleons in the core are assumed to be excited only in pairs coming from the same core orbit. Therefore, the angular momentum of the core is always even.

For the description of core-excited configurations in second quantization formalism an orthogonal system of four-fermion operators is constructed. Using a phenomenological interaction potential with appropriate spin-spin and tensor parts and restricting to excitations with core-spin 0 energy levels and transition probabilities of the nuclei  $\text{Pb}^{206}$ ,  $\text{Po}^{210}$  and  $\text{Hg}^{206}$  are calculated.