Zeitschrift: Helvetica Physica Acta

Band: 44 (1971)

Heft: 2

Artikel: Mass differences as additional electromagnetic corrections in low

energy elastic and charge exchange N scattering

Autor: Oudes, G.C. / Rasche, G.

DOI: https://doi.org/10.5169/seals-114274

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Mass Differences as Additional Electromagnetic Corrections in Low Energy Elastic and Charge Exchange π N Scattering

by G. C. Oades and G. Rasche

Institut für Theoretische Physik der Universität Zürich

(17. VIII. 70)

Summary. We present a treatment of low energy $\pi^- p \to \pi^- p$ and $\pi^- p \to \pi^0 n$ scattering which includes non-relativistically both mass difference effects and the effect of the long range Coulomb potential. Using this formalism we then show how the corrections to the usual charge independent expressions can be calculated in a first order perturbation treatment.

1. Introduction

In a recent paper [1] we have given a formalism which takes into account the Coulomb corrections to the charge independent nuclear πN interaction. These corrections are important when one wishes to isolate the purely nuclear scattering amplitude for use in dispersion relations or in tests of charge independence. It is usually assumed that these Coulomb effects constitute the bulk of the low energy electromagnetic corrections but, to be consistent, it is also necessary to include the electromagnetic mass differences between the proton and the neutron and between the π^{\pm} and π^{0} in the analysis of the coupled processes $\pi^{-} \phi \to \pi^{-} \phi$ and $\pi^{-} \phi \to \pi^{0} n$.

The usual procedure, when analysing differential cross sections, is to take mass differences into account kinematically when extracting the scattering amplitude; the dynamic effects of these mass differences on the scattering amplitude are not treated. In the present paper we include these latter effects in our non-relativistic treatment of the Coulomb corrections. In so doing we adhere as closely as possible to the notation used in Ref. [1]. In case of any confusion or for a more detailed discussion of the problem the reader is referred to this paper which also quotes some of the earlier literature.

In Section 2 we write down the Schrödinger equation in the simultaneous presence of mass differences and Coulomb effects. In Section 3 we deal with the inner Coulomb and mass difference corrections and in Section 4 we deal with the outer Coulomb and mass difference corrections. In Section 5, for use in experimental analyses, we give perturbation expressions correct to first order in the Coulomb and mass difference effects. Finally in two appendices we give the connection between our corrected S-matrix elements and the differential cross sections and we check that our S-matrix, as constructed from the coupled two channel Schrödinger equation, is unitary and symmetric.

2. The Schrödinger Equation

Since the potential energy operator in the Hamiltonian is non-controversial, we concentrate our attention on the kinetic energy term

$$H_{kin} = \frac{1}{2 m_{ab}^{(1)}} \left(p^{(1)} \right)^2 + \frac{1}{2 m_{ab}^{(2)}} \left(p^{(2)} \right)^2 . \tag{1}$$

Here the superscripts (1) and (2) stand for the pion and nucleon respectively. Because of mass differences the mass factors are operators in isospin space with the form

$$\frac{1}{m_{op}^{(1)}} = -\frac{1}{\mu_{-}} \frac{1}{2} t_{3} (1 - t_{3}) + \frac{1}{\mu_{0}} (1 + t_{3}) (1 - t_{3})$$

$$\frac{1}{m_{op}^{(2)}} = \frac{1}{m_{p}} \frac{1}{2} (1 + \tau_{3}) + \frac{1}{m_{n}} \frac{1}{2} (1 - \tau_{3})$$
(2)

where μ_{-} is the mass of the π^{-} , μ_{0} the mass of the π^{0} , m_{p} the mass of the proton and m_{n} the mass of the neutron. t and $1/2\tau$ are the pion and nucleon isospin operators. The operators in (2) are diagonal in the charge basis of isospin space, $|c\rangle$ (c=-, 0 denoting the π^{-} p and π^{0} n states), and so the inverses of these operators exist, having the form

$$m_{op}^{(1)} = -\mu_{-} \frac{1}{2} t_{3} (1 - t_{3}) + \mu_{0} (1 + t_{3}) (1 - t_{3})$$

$$m_{op}^{(2)} = m_{p} \frac{1}{2} (1 + \tau_{3}) + m_{n} \frac{1}{2} (1 - \tau_{3}) .$$
(3)

The operators $m_{op}^{(1)}$ and $m_{op}^{(2)}$ commute so one can introduce total linear momentum P and relative linear momentum P and transform to the c.m. frame P = 0.

Using the same algebraic manipulations as in the single channel case we obtain

$$H_{kin} = \frac{1}{2 m_{ob}} (p)^2$$
, $P = 0$ (4)

where

$$rac{1}{m_{op}} = rac{1}{m_{op}^{(1)}} + rac{1}{m_{op}^{(2)}} \; \cdot$$

In the charge basis we have in matrix notation

$$\frac{1}{m_{op}} = \begin{pmatrix} \frac{1}{m_p} + \frac{1}{\mu_-} & 0\\ 0 & \frac{1}{m_n} + \frac{1}{\mu_0} \end{pmatrix} = \begin{pmatrix} \frac{1}{m_-} & 0\\ 0 & \frac{1}{m_0} \end{pmatrix}$$
 (5)

where m_- and m_0 are the reduced masses of the $\pi^- p$ and $\pi^0 n$ systems. The inverse of (5) is obviously

$$m_{op} = \begin{pmatrix} m_{-} & 0 \\ 0 & m_{0} \end{pmatrix} \quad . \tag{6}$$

Thus we see that taking into account mass differences in the kinetic energy for the relative motion amounts to replacing the reduced mass by a mass operator which, in the charge basis, has the form given by (6). In addition, one has to introduce a rest mass operator which takes into account the difference in the rest energies of the two channels. The form of this operator is:

$$M_{op} = m_p \frac{1}{2} (1 + \tau_3) + m_n \frac{1}{2} (1 - \tau_3) + \mu_0 (1 + t_3) (1 - t_3) - \mu_{-} \frac{1}{2} t_3 (1 - t_3)$$

or, in the charge basis

$$M_{op} = \begin{pmatrix} m_p + \mu_- & 0 \\ 0 & m_n + \mu_0 \end{pmatrix}$$

Using units with $\hbar = c = 1$ we arrive at the Schrödinger equation

$$\left(-rac{1}{2\;m_{op}}\varDelta+U_{op}+V_{op}+M_{op}
ight)\psi=E\;\psi$$

where U_{op} is the strong, charge independent nuclear potential and V_{op} is the long range Coulomb potential. U_{op} is diagonal in the total isospin basis of isospin space. We assume that no other channels are open so that U_{op} is a real potential. The extension to the inelastic scattering region by the introduction of a complex U_{op} is trivial. The only essential difference being that our 2×2 S-matrix is no longer unitary. V_{op} has the form

$$V_{op} = V_c(r) t_3 1/2 (1 + \tau_3)$$
 ,

 $-V_c(r)$ being the static Coulomb potential in the $\pi^- p$ state. The wave function ψ depends on the relative coordinates in the c.m. frame, on the nucleon spin variable, and on the isospin variables of the pion and of the nucleon.

We now go over to the partial waves corresponding to diagonal absolute value of the total angular momentum, 3rd component of the total angular momentum and parity,

$$\psi_{l\pm}(x,s) = \left[R_{l\pm}^{(-)}(r) \mid -\rangle + R_{l\pm}^{(0)}(r) \mid 0\rangle\right] \frac{1}{r} \Omega_{l\pm,s}(\theta,\phi)$$

where $\Omega_{l\pm,s}$ is defined in Ref. [1]. The radial wave equations for the functions $R_{l\pm}$ can now be written down. For simplicity we only consider the case of l=0, although the results can be immediately generalised, and we use the notation of Ref. [1]

$$R_{o+}^{(c)} \equiv R_c^{TOT}$$
.

The superscript ^{TOT} means that the radial wave functions are exact solutions of the coupled channel system in distinction to solutions of other equations which will be introduced later.

The system of coupled equations determining R_c^{TOT} can be written in matrix form as

$$\left[\frac{1}{2}\begin{bmatrix} \frac{1}{m_{-}} & 0\\ 0 & \frac{1}{m_{0}} \end{bmatrix} \frac{d^{2}}{dr^{2}} + \begin{pmatrix} E - m_{p} - \mu_{-} & 0\\ 0 & E - m_{n} - \mu_{0} \end{pmatrix} - \begin{bmatrix} \frac{1}{3} U_{3} + \frac{2}{3} U_{1} - V_{c} \frac{\sqrt{2}}{3} (U_{3} - U_{1})\\ \frac{\sqrt{2}}{3} (U_{3} - U_{1}) & \frac{2}{3} U_{3} + \frac{1}{3} U_{1} \end{bmatrix} \right] \begin{pmatrix} R_{-}^{TOT} \\ R_{0}^{TOT} \end{pmatrix} = 0.$$
(7)

Here U_3 and U_1 are the s-wave diagonal values of U_{op} in the total isospin basis of isospin space. Defining wave numbers corresponding to the relative linear momenta of the two channels by

$$k_{-}^{2} \equiv 2 m_{-} (E - m_{p} - \mu_{-})$$

 $k_{0}^{2} \equiv 2 m_{0} (E - m_{n} - \mu_{0})$

we can rewrite (7) in the form

We can alternatively transform this equation to the total isospin basis when we obtain the corresponding equation for R_{2T}^{TOT} , the s-wave radial wave functions in the total isospin basis.

Defining

$$m_{-} - m_{0} \equiv \Delta m$$

$$k_{0}^{2} - k_{-}^{2} \equiv \Delta k^{2} \tag{9}$$

the transformed equation becomes

$$\begin{bmatrix}
\frac{d^{2}}{dr^{2}} + k_{-}^{2} + \frac{2}{3} \Delta k^{2} & \frac{\sqrt{2}}{3} \Delta k^{2} \\
\frac{\sqrt{2}}{3} \Delta k^{2} & \frac{d^{2}}{dr^{2}} + k_{-}^{2} + \frac{1}{3} \Delta k^{2}
\end{bmatrix} \\
- \begin{bmatrix}
2 m_{-} U_{3} - \frac{2}{3} m_{-} V_{c} - \frac{4}{3} \Delta m U_{3} & \frac{2\sqrt{2}}{3} m_{-} V_{c} - \frac{2\sqrt{2}}{3} \Delta m U_{1} \\
\frac{2\sqrt{2}}{3} m_{-} V_{c} - \frac{2\sqrt{2}}{3} \Delta m U_{3} & 2 m_{-} U_{1} - \frac{4}{3} m_{-} V_{c} - \frac{2}{3} \Delta m U_{1}
\end{bmatrix} \begin{bmatrix}
R_{3}^{TOT} \\
R_{1}^{TOT}
\end{pmatrix} = 0 .$$
(10)

3. The Inner Corrections

We first consider the functions R_{2T}^{IN} which are solutions of the equation obtained from (10) when mass differences and the Coulomb potential are only included for $r \leqslant r_0$, where

$$r_0 = \max(r_N, r_c) .$$

Here r_N is the range of the nuclear interaction and r_c is the charge radius beyond which the $\pi^- \phi$ Coulomb potential behaves like a point charge potential. Thus we have

$$\begin{bmatrix}
\frac{d^2}{dr^2} + k_-^2 & 0 \\
0 & \frac{d^2}{dr^2} + k_-^2
\end{bmatrix}
\begin{pmatrix}
R_3^{IN} \\
R_1^{IN}
\end{pmatrix} = 0 \quad r \geqslant r_0$$
(11)

while for $r \leqslant r_0$ the functions R_{2T}^{IN} obey (10).

We now introduce solutions R_{2T}^N of the purely nuclear problem i.e.

$$\begin{bmatrix} \frac{d^2}{dr^2} + k_{-}^2 - 2 m_{-} U_3 & 0 \\ 0 & \frac{d^2}{dr^2} + k_{-}^2 - 2 m_{-} U_1 \end{bmatrix} \begin{pmatrix} R_3^N \\ R_1^N \end{pmatrix} = 0$$
 (12)

and choose the two linearly independent solutions which have the behaviour

$$\begin{pmatrix}
R_{3\alpha}^{N} \\
R_{1\alpha}^{N}
\end{pmatrix} = \begin{pmatrix}
\sin(k_{-}r + \delta_{3}) \\
0
\end{pmatrix} \quad r \geqslant r_{N}$$

$$\begin{pmatrix}
R_{3\beta}^{N} \\
R_{1\beta}^{N}
\end{pmatrix} = \begin{pmatrix}
0 \\
\sin(k_{-}r + \delta_{1})
\end{pmatrix} \quad r \geqslant r_{N}$$
(13)

where δ_{2T} are the purely nuclear charge independent s-wave phases. We have arbitrarily defined the charge independent phases to correspond to scattering with the π^{\pm} and proton masses so as to avoid mass corrections in $\pi^{+} \phi$ scattering.

Since (11) and (12) coincide for $r \ge r_0$ we can write the general solution of (11) in terms of four constants α_3 , α_1 , β_3 , β_1 , in the form

$$R_{3}^{IN} = \alpha_{3} R_{3\alpha}^{N} + \beta_{3} R_{1\beta}^{N}, R_{1}^{IN} = \alpha_{1} R_{3\alpha}^{N} + \beta_{1} R_{1\beta}^{N}, \qquad r \geqslant r_{0}$$

$$(14)$$

Performing the same manipulations as in Ref. [1] we arrive at implicit relations between these constants of the form

$$\chi_{3} \equiv \beta_{3} \sin (\delta_{3} - \delta_{1}) =
= -\frac{1}{3 k_{-}} \int_{0}^{r_{0}} dr \, R_{3\alpha}^{N} \left[2 m_{-} V_{c} \left(R_{3}^{IN} - \sqrt{2} R_{1}^{IN} \right) \right.
\left. + 2 \sqrt{2} \, \Delta m \left(\sqrt{2} U_{3} R_{3}^{IN} + U_{1} R_{1}^{IN} \right) \right.
\left. + \sqrt{2} \, \Delta k^{2} \left(\sqrt{2} R_{3}^{IN} + R_{1}^{IN} \right) \right]$$

$$\chi_{1} \equiv \alpha_{1} \sin \left(\delta_{3} - \delta_{1} \right) =$$
(15a)

$$= -\frac{1}{3 k_{-}} \int_{0}^{r_{0}} dr \ R_{1\beta}^{N} \left[2 \sqrt{2} \ m_{-} V_{c} \left(R_{3}^{IN} - \sqrt{2} \ R_{1}^{IN} \right) \right. \\ \left. - 2 \Delta m \left(\sqrt{2} \ U_{3} \ R_{3}^{IN} + U_{1} \ R_{1}^{IN} \right) \right. \\ \left. - \Delta k^{2} \left(\sqrt{2} \ R_{3}^{IN} + R_{1}^{IN} \right) \right] \ . \tag{15b}$$

Thus combining (14) and (15) we have

$$egin{align} R_3^{IN} &= lpha_3 \; R_{3lpha}^N + rac{\chi_3}{\sin{(\delta_3 - \delta_1)}} \, R_{1eta}^N \ & r \!\geqslant\! r_0. \ R_1^{IN} \!=\! rac{\chi_1}{\sin{(\delta_3 - \delta_1)}} \, R_{3lpha}^N + eta_1 \, R_{1eta}^N \ & \end{array}$$

Finally we select two linearly independent solutions $R_{2T,\alpha}^{IN}$ and $R_{2T,\beta}^{IN}$ corresponding to the choices $\alpha_3 = 1$, $\beta_1 = 0$ and $\alpha_3 = 0$, $\beta_1 = 1$ i.e.

$$R_{3\alpha}^{IN} = R_{3\alpha}^{N} + \frac{\chi_{3\alpha}}{\sin(\delta_{3} - \delta_{1})} R_{1\beta}^{N}$$

$$R_{1\alpha}^{IN} = \frac{\chi_{1\alpha}}{\sin(\delta_{3} - \delta_{1})} R_{3\alpha}^{N}$$

$$r \geqslant r_{0}$$

$$R_{3\beta}^{IN} = \frac{\chi_{3\beta}}{\sin(\delta_{3} - \delta_{1})} R_{1\beta}^{N}$$

$$R_{1\beta}^{IN} = \frac{\chi_{1\beta}}{\sin(\delta_{3} - \delta_{1})} R_{3\alpha}^{N} + R_{1\beta}^{N}$$
(16)

4. The Outer Corrections

We now have to take into account the tail of the Coulomb potential and the effect of the mass differences for $r \ge r_0$. It is convenient to work in the charge basis so we transform (16) to obtain

$$R_{-\alpha}^{IN} = \sqrt{\frac{1}{3}} R_{3\alpha}^{N} + \frac{\sqrt{\frac{1}{3}} \chi_{3\alpha} R_{1\beta}^{N} - \sqrt{\frac{2}{3}} \chi_{1\alpha} R_{3\alpha}^{N}}{\sin(\delta_{3} - \delta_{1})}$$

$$R_{-\beta}^{IN} = -\sqrt{\frac{2}{3}} R_{1\beta}^{N} + \frac{\sqrt{\frac{1}{3}} \chi_{3\beta} R_{1\beta}^{N} - \sqrt{\frac{2}{3}} \chi_{1\beta} R_{3\alpha}^{N}}{\sin(\delta_{3} - \delta_{1})}$$

$$r \geqslant r_{0} . \tag{17}$$

$$R_{0\alpha}^{IN} = \sqrt{\frac{2}{3}} R_{3\alpha}^{N} + \frac{\sqrt{\frac{2}{3}} \chi_{3\alpha} R_{1\beta}^{N} + \sqrt{\frac{1}{3}} \chi_{1\alpha} R_{3\alpha}^{N}}{\sin(\delta_{3} - \delta_{1})}$$

$$R_{0\beta}^{IN} = \sqrt{\frac{1}{3}} R_{1\beta}^{N} + \frac{\sqrt{\frac{2}{3}} \chi_{3\beta} R_{1\beta}^{N} + \sqrt{\frac{1}{3}} \chi_{1\beta} R_{3\alpha}^{N}}{\sin(\delta_{3} - \delta_{1})}$$

From (8) we see that R_c^{TOT} and R_c^{IN} satisfy the same coupled equations for $r \leqslant r_0$ and so we can write

$$\begin{pmatrix} R_{-}^{TOT} \\ R_{0}^{TOT} \end{pmatrix} = A_{\alpha} \begin{pmatrix} R_{-\alpha}^{IN} \\ R_{0\alpha}^{IN} \end{pmatrix} + A_{\beta} \begin{pmatrix} R_{-\beta}^{IN} \\ R_{0\beta}^{IN} \end{pmatrix} \qquad r \le r_{0}$$

$$(18)$$

where A_{α} and A_{β} are arbitrary constants. Also from (8) we see that for $r \geqslant r_0$, R_c^{TOT} satisfy the equations

$$\left(\frac{d^2}{dr^2} + k_-^2\right) R_-^{TOT} + 2 m_- V_c R_-^{TOT} = 0$$
(19a)

$$\left(\frac{d^2}{dr^2} + k_0^2\right) R_0^{TOT} = 0 . (19b)$$

We now define two independent solutions of (19a), u_{α} and u_{β} , with asymptotic behaviour

$$u_{\alpha} \sim \sin(k_{-}r + \tau_{\alpha}^{(-)} - \sigma_{0} + \eta \ln 2 k_{-}r)$$

$$u_{\beta} \sim \sin(k_{-}r + \tau_{\beta}^{(-)} - \sigma_{0} + \eta \ln 2 k_{-}r)$$

$$r \to \infty$$
(20)

where the Coulomb parameter η is given by

$$\eta = \frac{e^2 m_-}{k_-}$$
 $(e^2 = \alpha = 137.0388^{-1})$

and the s-wave Coulomb phase is given by

$$\sigma_0 = \arg \Gamma (1 + i \eta) .$$

The phases $\tau_{\alpha}^{(-)}$ and $\tau_{\beta}^{(-)}$ are fixed by imposing the condition

$$\frac{u_{\alpha}'}{u_{\alpha}} \Big|_{r=r_0} = \frac{R_{-\alpha}^{IN'}}{R_{-\alpha}^{IN}} \Big|_{r=r_0}$$

$$\frac{u_{\beta}'}{u_{\beta}} \Big|_{r=r_0} = \frac{R_{-\beta}^{IN'}}{R_{-\beta}^{IN}} \Big|_{r=r_0}$$
(21)

where the prime denotes differentiation with respect to r. For $r \geqslant r_0$, R_-^{TOT} obeys the same equation as u_{α} and u_{β} so we can write

$$R_{-}^{TOT} = B_{\alpha} u_{\alpha} + B_{\beta} u_{\beta} \qquad r \geqslant r_{0}. \tag{22}$$

Matching value and derivative of R_{-}^{TOT} at r_0 as given by (18) and by (22) we obtain

$$B_{\alpha} = X_{\alpha} A_{\alpha} , \quad B_{\beta} = X_{\beta} A_{\beta}$$
 (23)

where the constants X_{α} and X_{β} are given by

$$u_{\alpha}(r_{0}) = X_{\alpha}^{-1} R_{-\alpha}^{IN}(r_{0})$$

$$u_{\beta}(r_{0}) = X_{\beta}^{-1} R_{-\beta}^{IN}(r_{0}) . \tag{24}$$

In the same spirit we define two independent solutions of (19b), v_{α} and v_{β} with asymptotic behaviour

$$v_{\alpha} \sim \sin(k_{0} r + \tau_{\alpha}^{(0)})$$

$$v_{\beta} \sim \sin(k_{0} r + \tau_{\beta}^{(0)}),$$

$$r \to \infty$$
(25)

the phases $\tau_{\alpha}^{(0)}$ and $\tau_{\beta}^{(0)}$ being determined by

$$\frac{v_{\alpha}'}{v_{\alpha}} \Big|_{r=r_0} = \frac{R_{0\alpha}^{IN}}{R_{0\alpha}^{IN}} \Big|_{r=r_0}$$

$$\frac{v_{\beta}'}{v_{\beta}} \Big|_{r=r_0} = \frac{R_{0\beta}^{IN}'}{R_{0\beta}^{IN}} \Big|_{r=r_0}$$
(26)

We also define two constants Y_{α} and Y_{β} by

$$v_{\alpha}(r_0) = Y_{\alpha}^{-1} R_{0\alpha}^{IN}(r_0) v_{\beta}(r_0) = Y_{\beta}^{-1} R_{0\beta}^{IN}(r_0) .$$
 (27)

For $r \ge r_0$, R_0^{TOT} obeys the same equation as v_α and v_β and so we arrive at the analogous result to (22) and (23),

$$R_0^{TOT} = A_{\alpha} Y_{\alpha} v_{\alpha} + A_{\beta} Y_{\beta} v_{\beta} \qquad r \geqslant r_0.$$
 (28)

Thus our general solution of the two channel $\pi^- p$, $\pi^0 n$ problem with Coulomb interaction and mass differences included has the form

$$\begin{pmatrix} R_{-}^{TOT} \\ R_{0}^{TOT} \end{pmatrix} = A_{\alpha} \begin{pmatrix} X_{\alpha} u_{\alpha} \\ Y_{\alpha} v_{\alpha} \end{pmatrix} + A_{\beta} \begin{pmatrix} X_{\beta} u_{\beta} \\ Y_{\beta} v_{\beta} \end{pmatrix} \quad r \geqslant r_{0}$$
(29)

where A_{α} and A_{β} are arbitrary constants.

Using the known asymptotic behaviour of the u's and the v's we can proceed from (29) to obtain the S-matrix elements in the charge basis, taking care to include the correct normalisation factors since the $\pi^- p$ and $\pi^0 n$ systems have different relative velocities. The resulting expressions are:

$$S_{TOT}^{--} = \left[X_{\alpha} Y_{\beta} e^{i \left(\tau_{\alpha}^{(-)} - \tau_{\beta}^{(0)} \right)} - X_{\beta} Y_{\alpha} e^{i \left(\tau_{\beta}^{(-)} - \tau_{\beta}^{(0)} \right)} \right] D_{TOT}^{-1}$$

$$S_{TOT}^{-0} = \left(\frac{k_{-} m_{0}}{k_{0} m_{-}} \right)^{1/2} \left[-2 i X_{\alpha} X_{\beta} \sin \left(\tau_{\alpha}^{(-)} - \tau_{\beta}^{(-)} \right) \right] D_{TOT}^{-1}$$

$$S_{TOT}^{0-} = \left(\frac{k_{0} m_{-}}{k_{-} m_{0}} \right)^{1/2} \left[2 i Y_{\alpha} Y_{\beta} \sin \left(\tau_{\alpha}^{(0)} - \tau_{\beta}^{(0)} \right) \right] D_{TOT}^{-1}$$

$$S_{TOT}^{00} = \left[X_{\alpha} Y_{\beta} e^{-i \left(\tau_{\alpha}^{(-)} - \tau_{\beta}^{(0)} \right)} - X_{\beta} Y_{\alpha} e^{-i \left(\tau_{\beta}^{(-)} - \tau_{\alpha}^{(0)} \right)} \right] D_{TOT}^{-1}$$

$$(30)$$

where

$$D_{TOT} = X_{\alpha} Y_{\beta} e^{-i(\tau_{\alpha}^{(-)} + \tau_{\beta}^{(0)})} - X_{\beta} Y_{\alpha} e^{-i(\tau_{\beta}^{(-)} + \tau_{\alpha}^{(0)})}$$
(31)

In Appendix I we show how the scattering amplitudes are constructed from these S-matrix elements and in Appendix II we check that the values given by (30) and (31) correspond to the elements of a symmetric and unitary matrix.

5. Perturbation Expressions

We have shown in the previous section how the full S-matrix elements can be obtained once the functions R_c^{IN} are known. These R_c^{IN} can be obtained by numerical solution of the coupled equations for $0 \le r \le r_0$ or alternatively they can be approximated by replacing the R_{2T}^{IN} by R_{2T}^{N} in (15a) and (15b) thus obtaining approximate values of the χ 's to substitute into (16). This latter procedure amounts to treating the Coulomb interaction and the mass differences as perturbations. The first order values obtained in this way are

$$\chi_{3\alpha} = -\frac{1}{3 \, k_{-}} \int_{0}^{r_{0}} dr (R_{3\alpha}^{N})^{2} (2 \, m_{-} \, V_{c} + 4 \, \Delta m \, U_{3} + 2 \, \Delta k^{2})$$

$$\chi_{3\beta} = \frac{1}{3 \, k_{-}} \int_{0}^{r_{0}} dr (R_{3\alpha}^{N} \, R_{1\beta}^{N}) (2 \, \sqrt{2} \, m_{-} \, V_{c} - 2 \, \sqrt{2} \, \Delta m \, U_{1} - \sqrt{2} \, \Delta k^{2})$$

$$\chi_{1\alpha} = -\frac{1}{3k_{-}} \int_{0}^{r_{0}} dr (R_{3\alpha}^{N} R_{1\beta}^{N}) (2\sqrt{2} m_{-} V_{c} - 2 | \sqrt{2} \Delta m U_{3} - \sqrt{2} \Delta k^{2})$$

$$\chi_{1\beta} = \frac{1}{3k_{-}} \int_{0}^{r_{0}} dr (R_{1\beta}^{N})^{2} (4 m_{-} V_{c} + 2\Delta m U_{1} + \Delta k^{2}) . \tag{32}$$

The nuclear potentials U_3 and U_1 can be eliminated by using (12) to obtain

$$U_{2T} = \left(\frac{d^2 R_{2T}^N}{dr^2} + k_-^2 R_{2T}^N\right) \cdot (2 m_- R_{2T}^N)^{-1}.$$

Having obtained the R_c^{IN} , either numerically or in the approximation described above, we then have to calculate values for the X's, $\tau^{(-)}$'s, Y's and $\tau^{(0)}$'s. Using (20) and (24) and matching value and derivative at r_0 we obtain

$$\tan \tau_{\alpha}^{(-)} = \frac{F_{0}(-\eta, k_{-}r) R_{-\alpha}^{IN'}(r) - F_{0}(-\eta, k_{-}r)' R_{-\alpha}^{IN}(r)}{G'_{0}(-\eta, k_{-}r) R_{-\alpha}^{IN}(r) - G_{0}(-\eta, k_{-}r) R_{-\alpha}^{IN'}(r)} \bigg|_{r = r_{0}}$$

$$X_{\alpha} = \frac{R_{-\alpha}^{IN}(r)}{\cos \tau_{\alpha}^{(-)} F_{0}(-\eta, k_{-}r) + \sin \tau_{\alpha}^{(-)} G_{0}(-\eta, k_{-}r)} \bigg|_{r = r_{0}}.$$
(33)

Analogous expressions are obtained for $\tau_{\beta}^{(-)}$ and X_{β} by replacing $R_{-\alpha}^{IN}$ by $R_{-\beta}^{IN}$. In a similar way using (25) and (27) we obtain

$$\tan \tau_{\alpha}^{(0)} = \frac{\begin{bmatrix} k_{0} r j_{0}(k_{0} r) \end{bmatrix} R_{0\alpha}^{IN'}(r) - [k_{0} r j_{0}(k_{0} r)]' R_{0\alpha}^{IN}(r)}{[k_{0} r n_{0}(k_{0} r)] R_{0\alpha}^{IN'}(r) - [k_{0} r n_{0}(k_{0} r)]' R_{0\alpha}^{IN}(r)} \Big|_{r = r_{0}}$$

$$Y_{\alpha} = \frac{R_{0\alpha}^{IN}(r)}{\cos \tau_{\alpha}^{(0)} [k_{0} r j_{0}(k_{0} r)] - \sin \tau_{\alpha}^{(0)} [k_{0} r n_{0}(k_{0} r)]} \Big|_{r = r_{0}}.$$
(34)

 $\tau_{\beta}^{(0)}$ and Y_{β} being obtained by replacing $R_{0\alpha}^{IN}$ by $R_{0\beta}^{IN}$. The notation suggests the obvious generalisation to $l \neq 0$.

Having obtained the values for the X's, Y's and τ 's the S-matrix elements can be calculated directly from (30) and (31). Alternatively values can be obtained for the eigenphases τ_3 and τ_1 and the mixing parameter ω as defined in Ref. [1]. We have from Ref. [1]

$$S_{TOT}^{--} + S_{TOT}^{00} = e^{2i\tau_3} + e^{2i\tau_1}$$
(35a)

$$S_{TOT}^{0-} = S_{TOT}^{-0} = \frac{\sqrt{2}}{3} \left(e^{2i\tau_3} - e^{2i\tau_1} \right) + \frac{\tan \omega}{3} \left(e^{2i\delta_3} - e^{2i\delta_1} \right)$$
 (35b)

+ higher order terms.

If we define Δ 's and ε 's by

$$\tau_{\alpha}^{(-)} = \delta_{3} + \Delta_{3}^{(-)} \qquad \tau_{\alpha}^{(0)} = \delta_{3} + \Delta_{3}^{(0)}
\tau_{\beta}^{(-)} = \delta_{1} + \Delta_{1}^{(-)} \qquad \tau_{\beta}^{(0)} = \delta_{1} + \Delta_{1}^{(0)}
X_{\alpha} = \sqrt{\frac{1}{3}} (1 + \varepsilon_{3}^{(-)}) \qquad Y_{\alpha} = \sqrt{\frac{2}{3}} (1 + \varepsilon_{3}^{(0)})
X_{\beta} = -\sqrt{\frac{2}{3}} (1 + \varepsilon_{1}^{(-)}) \qquad Y_{\beta} = \sqrt{\frac{1}{3}} (1 + \varepsilon_{1}^{(0)})$$
(36)

then from (30) and (31) we obtain the first order results

$$S_{TOT}^{--} + S_{TOT}^{00} = e^{2i\delta_3} \left(1 + 2 i \frac{\Delta_3^{(-)} + 2\Delta_3^{(0)}}{3} \right)$$

$$+ e^{2i\delta_1} \left(1 + 2 i \frac{2\Delta_1^{(-)} + \Delta_1^{(0)}}{3} \right)$$

$$S_{TOT}^{-0} = \frac{\sqrt{2}}{3} \left(\left(\frac{k_- m_0}{k_0 m_-} \right)^{1/2} + \frac{2\varepsilon_3^{(-)} - 2\varepsilon_3^{(0)} + \varepsilon_1^{(-)} - \varepsilon_1^{(0)}}{3} + i \frac{\Delta_3^{(-)} + 2\Delta_3^{(0)} + 2\Delta_1^{(-)} + \Delta_1^{(0)}}{3} \right)$$

$$\cdot \left\{ e^{2i\delta_3} \left[1 + i \left(\Delta_3^{(-)} - \Delta_1^{(-)} \right) \right] - e^{2i\delta_1} \left[1 - i \left(\Delta_3^{(-)} - \Delta_1^{(-)} \right) \right] \right\}.$$
(37a)

Since the decomposition of a complex number into the sum of two complex numbers each of modulus 1 is unique, we have on comparing (35a) and (37a),

$$\tau_{3} = \delta_{3} + \frac{\Delta_{3}^{(-)} + 2\Delta_{3}^{(0)}}{3}$$

$$\tau_{1} = \delta_{1} + \frac{2\Delta_{1}^{(-)} + \Delta_{1}^{(0)}}{3}.$$
(38)

Using (38) and comparing (35b) and (37b) we have

$$\tan \omega = \sqrt{2} \left[\left(\frac{k_{-} m_{0}}{k_{0} m_{-}} \right)^{1/2} - 1 \right] + \frac{\sqrt{2}}{3} \left[2 \left(\varepsilon_{3}^{(-)} - \varepsilon_{3}^{(0)} \right) + \left(\varepsilon_{1}^{(-)} - \varepsilon_{1}^{(0)} \right) + \frac{2 \left(\Delta_{3}^{(-)} - \Delta_{3}^{(0)} \right) - \left(\Delta_{1}^{(-)} - \Delta_{1}^{(0)} \right)}{\tan \left(\delta_{3} - \delta_{1} \right)} \right].$$
(39)

APPENDIX I

Differential Cross Section in Terms of S-Matrix Elements

The connection between the S-matrix elements and the differential cross sections proceeds in a similar way to that outlined in Section 3 of Ref. [1]. The asymptotic form of the radial and isotopic part of the wave function for a $\pi^- p$ incoming state of given l value is

$$\left[\sin \left(k_{-} r - \frac{l\pi}{2} - \sigma_{l} + \eta \ln 2 k_{-} r \right) + \frac{1}{2 i} \left(S_{l\pm}^{--} - 1 \right) e^{i(k_{-}r l\pi/2 - \sigma_{l} + \eta \ln 2 k_{-}r)} \mid - \right) \right] + \left[\frac{1}{2 i} \left(\frac{k_{-} m_{0}}{k_{0} m_{-}} \right)^{1/2} S_{l\pm}^{0-} e^{i(k_{0}r - l\pi/2)} \mid 0 \right) \right]$$
(A1.1)

where the $S_{l\pm}^{c'c}$ are the full S-matrix elements i.e. such that

$$S_{0+}^{c'c} = S_{TOT}^{c'c}$$
.

Defining the quantities

$$f_{l\pm}^{--} = \frac{1}{2i} (S_{l\pm}^{--} - 1) e^{-2i\sigma_l}$$

$$f_{l\pm}^{0-} = \frac{1}{2i} S_{l\pm}^{0-} e^{-i\sigma_l}$$

the same treatment as in Ref. [1] shows that the scattering amplitude can be written in the form

$$F^{c'-} = f^{c'-} + i \ \sigma \cdot \hat{n} \ g^{c'-}$$
 $c' = -, 0$

where

$$f^{c'-} = \delta_{c'-} F_{c_p} + \frac{1}{k_-} \sum_{l} \left[(l+1) f_{l\pm}^{c'-} + l f_{l-}^{c'-} \right] P_l (\cos \theta)$$

$$g^{c'-} = \frac{1}{k_-} \sum_{l} \left[f_{l+}^{c'-} - f_{l-}^{c'-} \right] P_l^1 (\cos \theta) .$$

 F_{c_p} is the point charge pure Coulomb scattering amplitude and \hat{n} is the normal to the scattering plane.

In terms of $f^{c'}$ and $g^{c'}$ the differential cross section is given by

$$rac{d\sigma}{d\Omega}_{-
ightarrow c'} = |f^{c'-}|^2 + |g^{c'-}|^2.$$

It should be noted that the usual treatments of $\pi^- p$ charge exchange scattering approximate the expression

$$\frac{1}{2i} \left(\frac{k_{-} m_{0}}{k_{0} m_{-}} \right)^{1/2} S_{l\pm}^{0-}$$

in equation (A1.1) by the charge independent limiting value in neglecting the mass differences and putting $\eta=0$. Making this approximation the mass difference effects are only allowed for in the kinematical factor which comes in, if one goes from the amplitude to cross section. One then can write

$$\frac{d\sigma}{d\Omega_{-\,\,\rightarrow\,\,0}}\,\cong\frac{v_0}{v_-}\,[\mid f_N^{0-}\mid^2\,+\,\mid g_N^{0-}\mid^2]\ .$$

Here v_0 and v_- are the relative c.m. velocities of the π^0 n and π^- p systems; f_N^{0-} and g_N^{0-} are the purely nuclear amplitudes, neglecting all isospin violating effects [2, 3].

APPENDIX II

Symmetry and Unitarity of the Full S-Matrix

As a check on our calculation we show explicitly that the S-matrix we obtain is symmetric and unitary.

a) Symmetry

Since u_{α} and u_{β} are two independent solutions of the same differential equation their Wronskian is independent of r and from (20) we have

$$W[u_{\alpha}, u_{\beta}] = k_{-} \sin(\tau_{\alpha}^{(-)} - \tau_{\beta}^{(-)})$$
.

Using (24) then gives

$$k_{-}\,X_{lpha}\,X_{eta}\sin{(au_{lpha}^{(-)}- au_{eta}^{(-)})}=W[R_{-lpha}^{IN}$$
 , $R_{-eta}^{IN}]_{r=r_{lpha}}$

Similarly

$$k_0 Y_{\alpha} Y_{\beta} \sin(\tau_{\alpha}^{(0)} - \tau_{\beta}^{(0)}) = W[R_{0\alpha}^{IN}, R_{0\beta}^{IN}]_{r=r_0}$$

and so from (30) we have

$$D_{TOT} \cdot (S_{TOT}^{0-} - S_{TOT}^{-0}) =$$

$$2 i \left(\frac{m_{-} m_{0}}{k_{-} k_{0}}\right)^{1/2} \left(\frac{1}{m_{-}} W \left[R_{-\alpha}^{IN}, R_{-\beta}^{IN}\right]_{r=r_{0}} + \frac{1}{m_{0}} W \left[R_{0\alpha}^{IN}, R_{0\beta}^{IN}\right]_{r=r_{0}}\right). \tag{A2.1}$$

However, $R_{c\alpha}^{IN}$ and $R_{c\beta}^{IN}$ are regular solutions of (8) for $r \leqslant r_0$.

Using this system of coupled differential equations one obtains

$$\frac{1}{m_{-}}W\left[R_{-\alpha}^{IN}, R_{-\beta}^{IN}\right] + \frac{1}{m_{0}}W\left[R_{0\alpha}^{IN}, R_{0\beta}^{IN}\right] = 0 \quad , \quad 0 \leqslant r \leqslant r_{0}.$$

Thus the right hand side of (A2.1) vanishes and so

$$S_{TOT}^{0-} = S_{TOT}^{-0}$$
 (A2.2)

b) Unitarity

We have to prove the three relations

$$|S_{TOT}^{--}|^2 + |S_{TOT}^{0-}|^2 = 1$$
, (A2.3)
 $|S_{TOT}^{00}|^2 + |S_{TOT}^{-0}|^2 = 1$, (A2.4)

$$|S_{TOT}^{00}|^2 + |S_{TOT}^{-0}|^2 = 1$$
 , (A2.4)

$$S_{TOT}^{--} S_{TOT}^{0-*} + S_{TOT}^{00*} S_{TOT}^{-0} = 0$$
 (A2.5)

Writing (A2.3) in the form

$$\mid D_{TOT} \cdot S_{TOT}^{0-} \mid^2 = \mid D_{TOT} \mid^2 - \mid D_{TOT} \cdot S_{TOT}^{--} \mid^2$$

we insert the explicit expressions from (30) and (31) to obtain

$$4 \frac{k_0 m_-}{k_- m_0} Y_{\alpha}^2 Y_{\beta}^2 \sin^2(\tau_{\alpha}^{(0)} - \tau_{\beta}^{(0)}) = -4 X_{\alpha} Y_{\beta} X_{\beta} Y_{\alpha} \sin(\tau_{\alpha}^{(-)} - \tau_{\beta}^{(-)}) \sin(\tau_{\alpha}^{(0)} - \tau_{\beta}^{(0)}).$$

Thus we must prove

$$\frac{k_0}{m_0} Y_{\alpha} Y_{\beta} \sin(\tau_{\alpha}^{(0)} - \tau_{\beta}^{(0)}) + \frac{k_{-}}{m_{-}} X_{\alpha} X_{\beta} \sin(\tau_{\alpha}^{(-)} - \tau_{\beta}^{(-)}) = 0$$

but this follows at once from the results of the previous section.

To prove (A2.4) we note from (30) that

$$(D_{TOT} \cdot S_{TOT}^{00})^* = (D_{TOT} \cdot S_{TOT}^{--}) \tag{A2.6}$$

and so (A2.4) follows at once from (A2.3) which we have just proved. Finally we use (A2.2) and (A2.6) to rewrite (A2.5) as

$$(D_{TOT} \cdot S_{TOT}^{--}) \cdot 2 \ Re \ (D_{TOT} \cdot S_{TOT}^{-0}) = 0$$

However, from (30), we see that $D_{TOT} S_{TOT}^{-0}$ is pure imaginary and so this relation is also proved to be true.

REFERENCES

G. C. OADES and G. RASCHE, Helv. phys. Acta, 44, 5 (1971).

For the single channel case see also J. Reignier Bulletin de l'Académie Royale de Belgique (Classe des Sciences) 48, 179 (1962).

- See p. 82, H. A. Bethe and F. de Hoffmann, Mesons and Fields, Vol. II (Row, Peterson + Co., New York 1955).
- See Section 4 of J. Hamilton, Phys. Rev. 110, 1134 (1958).