Zeitschrift: Helvetica Physica Acta

Band: 45 (1972)

Heft: 2

Artikel: Definition of Green's functions for dilute Fermi gases

Autor: Ruelle, David

DOI: https://doi.org/10.5169/seals-114378

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Definition of Green's Functions for Dilute Fermi Gases

by David Ruelle

I.H.E.S., 91, Bures-sur-Yvette, France

(20. XI. 71)

Abstract. The infinite volume limit is shown to exist for Green's functions of a large class of dilute Fermi gases. In particular, this gives meaning to the time evolution of the corresponding infinite systems.

I started struggling to understand statistical mechanics about ten years ago in Zurich, with the kind help of M. Fierz and R. Jost. It is nice to remember these discussions which Fierz insisted to have in German—after a few minutes German was replaced by Schwytzer-Dütsch. Few mathematically solid results were then known about thermodynamic limits, correlation functions, reduced density matrices, etc. Now, 'rigorous results' have become an industry but I hope that M. Fierz will find some pleasure in looking at this note dedicated to him on his sixtieth birthday.

It was shown in a recent note [6] that Ginibre's work on the reduced density matrices of quantum gases [3], [4] can be used to discuss the existence and analyticity of Green's functions. Here we sharpen the results of [6] in the case of Fermi systems. In particular the time correlation functions are seen to be well defined in the limit of an infinite volume for a dilute Fermi gas.

Green's functions are defined by

$$G_{\Lambda}^{\Phi}(\zeta) = Z^{-1} \operatorname{Tr}(A_{1} e^{-(\zeta_{2} - \zeta_{1})H_{\Lambda}} A_{2} \dots e^{-(\zeta_{m} - \zeta_{m-1})H_{\Lambda}} A_{m} e^{-(\beta + \zeta_{1} - \zeta_{m})H_{\Lambda}})$$

where H_{Λ} is the Hamiltonian for a bounded region Λ of \mathbb{R}^{ν} , $Z = \operatorname{Tr} e^{-\beta H_{\Lambda}}$, A_k is either $\int dx \, \phi_k(x) a(x)$ or $\int dx \, \phi_k(x) a^*(x)$, and $\phi_k \in L^2(\mathbb{R}^{\nu})^1$). We have used the notation $\boldsymbol{\zeta} = (\zeta_1, \ldots, \zeta_m), \boldsymbol{\phi} = (\phi_1, \ldots, \phi_m)$ and assumed that $\boldsymbol{\zeta} \in \overline{\mathcal{D}}$,

$$\overline{\mathcal{D}} = \{ \zeta \colon \operatorname{Re} \zeta_1 \leqslant \cdots \leqslant \operatorname{Re} \zeta_m \leqslant \operatorname{Re} \zeta_1 + \beta \}$$

For a system of particles interacting through a suitable pair potential Φ ,²) and for small activity, the operator $e^{-\lambda H_A}$, with $\lambda > 0$, may be defined in terms of Wiener

$$A_{k} = \int dx_{1} \dots dx_{p} dy_{1} \dots dy_{q} \phi_{k}(x_{1}, \dots, x_{p}, y_{1}, \dots, y_{q}) a^{*}(x_{1}) \dots a^{*}(x_{p}) a(y_{1}) \dots a(y_{q})$$

where $\phi_k \in L^2(\mathbb{R}^{v(p+q)})$, see [6].

2) Ginibre's conditions [4] are (A), (B) of the theorem below and

(C)
$$\int_{|x|>c} |\Phi(x)| dx < + \infty \text{ for some } C > 0.$$

We shall replace (C) by the stronger requirement

(C') $\Phi \in L^1(\mathbb{R}^{\nu}) \cap L^2(\mathbb{R}^{\nu})$.

¹⁾ More generally one could take

integrals and is of trace class (see Ginibre [1]). The operators $A_k e^{-\lambda H_A}$ can also be expressed in terms of Wiener integrals and are of trace class.

When λ is complex and $\operatorname{Re} \lambda > 0$, $e^{-\lambda H_A}$ is defined and analytic, therefore G_A^{Φ} is an analytic function of the complex variables $\zeta_k = \beta_k - it_k$ in the domain

$$\mathscr{D} = \{ \boldsymbol{\zeta} : \beta_1 < \cdots < \beta_m < \beta_1 + \beta \}$$

If $\zeta \in \overline{\mathcal{D}}$ and $t_1 = \cdots = t_n$, Ginibre's analysis shows that G_A^{Φ} tends to a limit when $A \to \infty$.³) From this the following result can be deduced (see [6]):

There exists a function G^{Φ} analytic in \mathcal{D} and such that

$$\lim_{\Lambda \to \infty} G_{\Lambda}^{\Phi}(\zeta) = G^{\Phi}(\zeta) \tag{1}$$

uniformly on compacts with respect to $\zeta \in \mathcal{D}$.

The theorem below shows that $G_{\Lambda}^{\Phi}(\zeta)$ has a limit G^{Φ} when $\Lambda \to \infty$ and ζ is in the closure $\overline{\mathcal{D}}$ of \mathcal{D} . This defines infinite volume Green's functions for real times and not just $\zeta \in \mathcal{D}$.

Theorem. Let $\Phi: \mathbb{R}^{\nu} \to \mathbb{R}$ satisfy the following conditions

- (A) Φ is even $(\Phi(x) = \Phi(-x))$ and continuous for $x \neq 0$
- (B) Φ is stable
- (C') $\int |\Phi(x)| dx < +\infty$, $\int |\Phi(x)|^2 dx < +\infty$

For a Fermi system with pair potential Φ at small activity, G^{Φ} extends to a bounded continuous function on $\overline{\mathcal{D}}$ such that

$$\lim_{\Lambda \to \infty} G_{\Lambda}^{\Phi}(\zeta) = G^{\Phi}(\zeta)$$

uniformly on the compacts of $\bar{\mathcal{D}}$.

We decompose the proof in several steps

Lemma 1. Let \mathcal{M}_n be defined by

$$\mathcal{M}_{n} = \left\{ \zeta \in \overline{\mathcal{D}} : \beta_{3} - \beta_{1} \geqslant \frac{\beta}{n}, \beta_{4} - \beta_{2} \geqslant \frac{\beta}{n}, \dots \right.$$

$$\dots, \beta_{m} - \beta_{m-2} \geqslant \frac{\beta}{n}, \beta + \beta_{1} - \beta_{m-1} \geqslant \frac{\beta}{n}, \beta + \beta_{2} - \beta_{m} \geqslant \frac{\beta}{n} \right\}$$

For sufficiently large Λ , G_{Λ}^{\bullet} is bounded on \mathcal{M}_n uniformly with respect to $\boldsymbol{\zeta}$ and Λ ; if ϕ_k is of class C^2 with compact support, the same is true of $\partial G_{\Lambda}^{\boldsymbol{\phi}}/\partial \zeta_k$. Furthermore $G_{\Lambda}^{\boldsymbol{\phi}}$ tends to a limit $G^{\boldsymbol{\phi}}$ uniformly on the compacts of \mathcal{M}_n when $\Lambda \to \infty$.

If $\zeta \in \mathcal{M}_n$, G_A^{Φ} can be expressed in terms of operators $e^{[it_k-(1/4n)\beta]H_A}A_k(\phi_k)e^{-it_kH_A}$, $e^{it_kH_A}A_k(\phi_k)e^{-[it_k+(1/4n)\beta]H_A}$,

The bounded open $\Lambda \subset \mathbb{R}^{\nu}$ form a directed set, when ordered by inclusion. We let $\lim G_{\Lambda}$ when $\Lambda \to \infty$ be by definition the limit of the directed family (G_{Λ}) .

and $e^{-\lambda H_{\Lambda}}$ with $0 < \lambda < \beta$. Using Hölder's inequality) we find an upper bound for $|G_{\Lambda}^{\Phi}|$ in terms of the expressions

$$Z^{-1} \operatorname{Tr}[(e^{-(1/4n)\beta H_A} A_k(\phi_k) A_k(\phi_k)^* e^{-(1/4n)\beta H_A})^n]$$

and

$$Z^{-1} \operatorname{Tr}[(e^{-(1/4n)\beta H_A} A_k(\phi_k)^* A_k(\phi_k) e^{-(1/4n)\beta H_A})^n]$$

These expressions are known to have a limit when $\Lambda \to \infty$. This proves the uniform boundedness of $|G_A^{\Phi}(\zeta)|$.

The derivative $\partial G_{\Lambda}^{\Phi}/\partial \zeta_k$ is obtained by replacing $A_k(\phi_k)$ by $[A_k(\phi_k), H_{\Lambda}]$ in the expression of G_{Λ}^{Φ} . One is thus brought to considering the expression

$$Z^{-1} \operatorname{Tr}[(e^{-(1/4n)\beta H_{\Lambda}}[A_{k}(\phi_{k}), H_{\Lambda}][A_{k}(\phi_{k}), H_{\Lambda}]^{*} e^{-(1/4n)\beta H_{\Lambda}})^{n}]$$
(2)

or a similar one with $[A_k(\phi_k), H_{\Lambda}]$ replaced by $[A_k(\phi_k), H_{\Lambda}]^*$. Since ϕ_k is of class C^2 with compact support, the commutator of $A_k(\phi_k)$ with the kinetic energy part of H_{Λ} is again of the form $A(\phi)$. The potential energy part of H_{Λ} is

$$\sim \int dx \int dy \, \Phi(x-y) a^*(x) a^*(y) a(x) a(y)$$

and its commutator with $A_k(\phi_k)$ is

$$\sim \int dx \int dy \, \phi_{\mathbf{k}}(x) \Phi(x-y) a^*(x) a^*(y) a(y)$$

or

$$\sim \int dx \int dy \, \phi_k(x) \Phi(x-y) a^*(y) a(x) a(y).$$

We insert these expressions in

$$[A_k(\phi_k), H_{\Delta}][A_k(\phi_k), H_{\Delta}]^*$$

and use the anticommutation relations to put the annihilation operators a to the right and the creation operators a^* to the left (Wick ordering). A number of terms are thus obtained which are conveniently described by diagrams; they are integrals of Wick ordered products $a^*(x_1) \dots a(x_r)$ multiplied by continuous functions $\phi(x_i)$, and the pair potential $\Phi(x_i - x_j)$. The pair potential appears as factor 0, 1, or 2 times. If $\Phi(x_i - x_j)$ appears there also appears a factor $\phi(x_i)$ or $\phi(x_j)$; for each variable x_i which does not appear in a factor $\Phi(x_j - x_i)$ there is a factor $\phi(x_i)$. Inserting now in (2) we obtain a sum of terms which are integrals of reduced density matrices multiplied by factors $\phi(x_k)$ and $\Phi(x_k - x_j)$ as described above. Using the condition (C') of the theorem, and the fact that the reduced density matrices are bounded functions, uniformly in Λ [3], be obtain a bound on (2) which is independent of Λ . This gives the desired boundedness of $\partial G_{\Lambda}^{\Phi}/\partial \zeta_k$.

The C^2 functions with compact support are dense in L^2 . Therefore we may assume that ϕ_1,\ldots,ϕ_m are C^2 with compact support in proving that G^{Φ}_{Λ} tends to a limit G^{Φ} uniformly on the compacts of \mathcal{M}_n when $\Lambda \to \infty$. Since the G^{Φ}_{Λ} and their derivatives $\partial G^{\Phi}_{\Lambda}/\partial \zeta_k$ are uniformly bounded, this results from $G^{\Phi}_{\Lambda}(\zeta) \to G^{\Phi}(\zeta)$ for $\zeta \in \mathcal{D}$ (see (1)).

⁴⁾ See [2] Lemma XI, 9-20, p. 1105.

An easy extension of Ginibre's results is actually needed at this point.

Lemma 2. Let ϕ be given and Λ_0 chosen sufficiently large. We assume that $1 \leq l \leq m$, that K is compact in \mathbb{R} and that $\zeta_{l+1}, \ldots, \zeta_m$ are fixed such that

$$0 < \operatorname{Re} \zeta_{l+1} < \cdots < \operatorname{Re} \zeta_m < \frac{\beta}{2}$$

Given $\epsilon > 0$, there exists $\delta > 0$ such that

$$|G_{\mathcal{A}}^{\Phi}(-i\theta_1,\ldots,-i\theta_l,\,\zeta_{l+1},\ldots,\zeta_m)-G_{\mathcal{A}}^{\Phi}(-i\theta_1,\ldots,-i\theta_{l-1},\,\zeta_l,\ldots,\zeta_m)|<\epsilon$$
 (3)

whenever $\Lambda \supset \Lambda_0$, and θ_l , ζ_l satisfy $\theta_l \in K$, $|\zeta_l + i\theta_l| < \delta$, $0 < \operatorname{Re} \zeta_l < \operatorname{Re} \zeta_{l+1}$. Furthermore, $G_{\Lambda}^{\Phi}(-i\theta_1, \ldots, -i\theta_l, \zeta_{l+1}, \ldots, \zeta_m)$ has a limit $G_{(l)}^{\Phi}(-i\theta_1, \ldots, -i\theta_l, \zeta_{l+1}, \ldots, \zeta_m)$ when $\Lambda \to \infty$, uniformly for $(\theta_1, \ldots, \theta_l) \in K^l$.

We have

$$\begin{split} &|G_{A}^{\Phi}(-i\theta_{1},\ldots,-i\theta_{l},\,\zeta_{l+1},\ldots,\zeta_{m})-G_{A}^{\Phi}(-i\theta_{1},\ldots,-i\theta_{l-1},\,\zeta_{l},\,\ldots,\zeta_{m})|\\ &=|Z^{-1}\operatorname{Tr}[A_{1}e^{i(\theta_{2}-\theta_{1})H_{A}}\ldots A_{l-1}\\ &\quad\times (e^{i(\theta_{l}-\theta_{l-1})H_{A}}A_{1}e^{-(\zeta_{l+1}+i\theta_{l})H_{A}}-e^{-(\zeta_{l}+i\theta_{l+1})H_{A}}A_{1}e^{-(\zeta_{l+1}-\zeta_{l})H_{A}})\\ &\quad\times A_{l+1}\ldots e^{-(\zeta_{m}-\zeta_{m-1})H_{A}}A_{m}\,e^{-(\beta-i\theta_{1}-\zeta_{m})H_{A}}]|\\ &\leqslant XY \end{split}$$

where

$$\begin{split} X^2 &= Z^{-1} \operatorname{Tr}(e^{-\beta H_A/2} A_1 e^{i(\theta_2 - \theta_1)H_A} \dots A_{l-1} A_{l-1}^* \dots e^{-i(\theta_2 - \theta_1)H_A} A_1^* e^{-\beta H_A/2}) \\ &\leqslant (\|A_1\| \dots \|A_{l-1}\|)^2 \\ Y^2 &= [F(\zeta_l, \zeta_l) - F(\zeta_l, -i\theta_l) - F(-i\theta_l, \zeta_l) + F(-i\theta_l, -i\theta_l)] \\ F(u, v) &= Z^{-1} \operatorname{Tr}[e^{-(\beta/2 - i\theta_1 - \zeta_m)^* H_A} A_m^* \dots A_{l+1}^* e^{-(\zeta_{l+1} - u)^* H_A} A_1^* \\ &\qquad \qquad \times e^{(-u^* - v)H_A} A_1 e^{-(\zeta_{l+1} - v)H_A} A_{l+1} \dots A_m e^{-(\beta/2 - i\theta_1 - \zeta_m)H_A}] \end{split}$$

and (3) results from Lemma 1.

We prove now the convergence of $G_{\Lambda}^{\Phi}(-i\theta_1,\ldots,-i\theta_l,\,\zeta_{l+1},\ldots,\,\zeta_m)$ when $\Lambda\to\infty$. For l=1 this is again a consequence of Lemma 1. For the general case we use induction on l. By the compactness of K, given $\delta>0$, there exist $\zeta_{l}^{(1)},\ldots,\,\zeta_{l}^{(N)}$ with the following properties

- a) $0 < \operatorname{Re} \zeta^{(j)} < \operatorname{Re} \zeta_{l+1}$,
- b) for each $\theta_l \in K$ there is a j such that $|\zeta_l^{(j)} + i\theta_l| < \delta$.

By the first part of the lemma we have thus

$$|G_{\Lambda}^{\Phi}(-i\theta_1,\ldots,-i\theta_l,\,\zeta_{l+1},\ldots,\zeta_m)-G_{\Lambda}^{\Phi}(-i\theta_1,\ldots,-i\theta_{l-1},\,\zeta_l^{(j)},\,\zeta_l,\ldots,\zeta_m)|<\epsilon.$$

On the other hand the induction assumption implies that, for $(\theta_1, \ldots, \theta_{l-1}) \in K^{l-1}$ and Λ sufficiently large, we have

$$|G_{\Lambda}^{\Phi}(-i\theta_1,\ldots,-i\theta_{l-1},\zeta_l^{(j)},\ldots,\zeta_m)-G_{(l-1)}^{\Phi}(-i\theta_1,\ldots,-i\theta_{l-1},\zeta_l^{(j)},\ldots,\zeta_m)|<\epsilon.$$

The uniform convergence of $G_{\Delta}^{\Phi}(-i\theta_1, \ldots, -i\theta_l, \zeta_{l+1}, \ldots, \zeta_m)$ follows.

Proof of the Theorem

Since the functions G_{Λ}^{Φ} are analytic in \mathcal{D} and bounded continuous in $\overline{\mathcal{D}}$, it is easily seen that uniform convergence on the compacts of the distinguished boundary \mathcal{S} implies uniform convergence on the compacts of $\overline{\mathcal{D}}$. The distinguished boundary is defined by

$$\mathcal{S} = \bigcup_{l=1}^{m} \mathcal{S}_{l}$$

$$\mathcal{S}_{l} = \{ \boldsymbol{\zeta} : \operatorname{Re} \zeta_{1} = \cdots = \operatorname{Re} \zeta_{l}, \operatorname{Re} \zeta_{l+1} = \cdots = \operatorname{Re} \zeta_{m} = \operatorname{Re} \zeta_{1} + \beta \}.$$

Therefore by Lemma 2 we have uniform convergence on the compacts of $\mathscr S$ when $\Lambda \to \infty$, and the theorem is proved.

Remark on time evolution

It is known that for certain quantum lattice systems, time evolution can be described by automorphisms of a 'standard' C*-algebra (see [5]). In this note we prove only the existence of Green's functions (a similar result is known for a large class of lattice systems [7]). It is probable that time evolution cannot be described here by automorphisms of the C*-algebra of canonical anticommutation relations, and it is unclear if it is given by automorphisms of its weak closure. For further considerations on this problem see [1].

REFERENCES

- [1] D. A. Dubin and G. L. Sewell, J. Math. Phys. 11, 2990-2998 (1970).
- [2] N. DUNFORD and J. SCHWARTZ, Linear Operators (Interscience, New York 1963).
- [3] J. GINIBRE, J. Math. Phys. 6, 238-251 (1965); 6, 252-262 (1965); 6, 1432-1446 (1965).
- [4] J. Ginibre, in: Statistical Mechanics, p. 148 (proceedings of the I.U.P.A.P. meeting, Copenhagen 1966) (Ed. T. Bak; Benjamin, New York 1967).
- [5] D. W. Robinson, Commun. math. Phys. 7, 337-348 (1968).
- [6] D. RUELLE, J. Math. Phys. 12, 901-903 (1971).
- [7] M. B. Ruskai, Commun. math. Phys. 20, 193-204 (1971).