Zeitschrift:	Helvetica Physica Acta
Band:	45 (1972)
Heft:	5
Artikel:	Coherence resonances in the alignment-orientation coupling process induced by an electric field
Autor:	Faist, A.
DOI:	https://doi.org/10.5169/seals-114412

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Birkhäuser Verlag Basel

# Coherence Resonances in the Alignment-Orientation Coupling Process Induced by an Electric Field

# by A. Faist

#### Laboratoire d'optique physique, EPF, Lausanne

#### (6. III. 72)

Summary. The alignment-orientation coupling process is studied in the case of atomic excited states in the presence of an electric field. The basic properties are obtained by applying a static electric field. The phenomenon is more complex in the case of an oscillating electric field and new resonances of the coherence type appear. The general properties of these new resonances are studied with the aid of computational methods.

We consider an atomic excited state of angular momentum J = 1 (I = 0) subject, in the presence of an electric field E, to a Stark effect giving rise to an energy separation [1].

 $\Delta W = -m^2 \beta_s = -m^2 \cdot a \cdot E^2$ 

These sublevels are optically pumped from a unique ground state (m = 0) and one observes the light re-emitted by fluorescence.

# A. Static Field E

Let us first consider the case of a static electric field. If  $\vec{E} = E \cdot [\vec{z}]$ , Oz being chosen as the axis of quantification, then the Hamiltonian is for the excited state:

$$\mathscr{H} = \begin{pmatrix} -\beta_s & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -\beta_s \end{pmatrix} \qquad \begin{array}{c} m = 1 \\ 0 \\ 0 \\ -1 \end{array}$$

As this matrix is diagonal and time independent, the stationary solution of the matrix density equation [2]

$$\dot{
ho}_{mm'}=\Gamma
ho_{mm'}^{\mathfrak{o}}-\Gamma
ho_{mm'}-i[\mathscr{H},
ho]_{mm'}$$

is given by  $(\hbar = 1)$ 

$$\rho_{mm'}^{\text{stat}} = \rho_{mm'}^{0} \frac{\Gamma}{\Gamma + i(W_m - W_{m'})}$$

where  $\Gamma$  is  $1/\tau$ ,  $\tau$  being the lifetime of the excited sublevel 1m.

The vector of polarization of the pumping light is chosen so as to give rise to a coherent excitation of the three sublevels



Geometric disposition of the polarizations of the pumping and fluorescence lights.

This gives the initial condition

$$\rho_{mm'}^{o} = \frac{1}{2} \begin{pmatrix} \frac{1}{2} & -\frac{1}{\sqrt{2}} & -\frac{1}{2} \\ -\frac{1}{\sqrt{2}} & 1 & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & 1 & \frac{1}{\sqrt{2}} \\ -\frac{1}{2} & \frac{1}{\sqrt{2}} & \frac{1}{2} \end{pmatrix}.$$

The stationary solution can now be written down

$$\rho_{\textit{mm'}}^{\text{stat}} = \frac{1}{2} \begin{pmatrix} \frac{1}{2} & -\frac{1}{\sqrt{2}} \frac{\Gamma}{\Gamma - i\beta_s} & -\frac{1}{2} \\ -\frac{1}{\sqrt{2}} \frac{\Gamma}{\Gamma + i\beta_s} & 1 & \frac{1}{\sqrt{2}} \frac{\Gamma}{\Gamma + i\beta_s} \\ -\frac{1}{2} & \frac{1}{\sqrt{2}} \frac{\Gamma}{\Gamma - i\beta_s} & \frac{1}{2} \end{pmatrix}.$$

The intensity of the re-emitted light can be calculated using the relation

$$I_{\mathbf{fl}} = \sum_{m} \sum_{m'} \langle 0 | \vec{e}_{\lambda} \vec{D} | m \rangle \langle m | \rho | m' \rangle \langle m' | \vec{e}_{\lambda} \vec{D} | 0 \rangle.$$

We shall first consider linearly polarized light with the vector of polarization parallel (+) or perpendicular (-) to the polarization of the pumping light:

$$\vec{e}_{\lambda} = rac{1}{\sqrt{2}} ([\vec{x}] \pm [\vec{z}])$$

Vol. 45, 1972 Coherence Resonances in Alignment-Orientation Coupling Process

$$\begin{split} I_{\rm fl}^{\rm lin} &\propto \frac{1}{2} (1 \pm {\rm Re}\,\rho_{10}) = \frac{1}{2} \left( 1 \pm \frac{\Gamma^2}{\Gamma^2 + \beta_s^2} \right) \\ &= \frac{1}{2} \left( 1 \pm \frac{\Gamma^2}{\Gamma^2 + a^2 E^4} \right). \end{split}$$

These are absorption curves given on Figure 2:



Figure 2

Linearly polarized re-emitted light.

As  $E^2$  is always positive, these curves have no left part. The shape is similar to the right-hand part of a Hanle curve with

 $\Gamma = aE_1^2$ 

where  $\Gamma$  is the inverse of the lifetime and a the Stark constant of the considered level. This particular aspect of the phenomenon had already been observed previously

[3].

The linearly polarized light with  $\vec{e}_{\lambda} = [\vec{x}]$  or  $\vec{e}_{\lambda} = [\vec{z}]$  has a constant intensity  $I = \frac{1}{2}$ . Let us now consider circularly polarized light re-emitted in the Oy direction [4]. We find:





Again  $I_{fl}^{circ\pm}$  does not depend on the sign of E.

One should notice that the 'creation' of orientation is maximum for  $E_1 = \sqrt{\Gamma/a}$  and vanishes for higher values of E[5].

## **B.** Oscillating Electric Field

Let us now consider the action of an oscillating electric field  $\vec{E} = [\vec{z}] E \cos \omega t$ . The (time-dependent) Stark separation is now

$$\Delta W = -m^2 a E^2 \cos^2 \omega t$$
$$= -m^2 \frac{a}{2} E^2 (1 + \cos 2\omega t)$$
$$= -\beta (1 + \cos 2\omega t)$$

Owing to the fact that the Stark effect is quadratic, the Hamiltonian is now the sum of a time-independent part  $(-\beta)$  and a time-dependent part  $(-\beta \cos 2\omega t)$ .

As before the only field strength dependent terms are the  $\rho_{m,m\pm 1}$  terms which are solution of the equation

$$\dot{\rho} = \Gamma \rho^{0} - \Gamma \rho + i(\beta + \beta \cos 2\omega t) \rho$$

where

$$\rho = \rho_{1,0} = \rho_{0,1}^{\times} = \rho_{-1,0} = \rho_{0,-1}^{\times}.$$

This is a typical coherence–resonance equation with the solution [6]

$$\frac{\rho}{\rho_0} = A_0 + \sum_{p=1}^{\infty} \left( A_p \, e^{i \, 2 \, p \, \omega t} + B_p \, e^{-i \, 2 \, p \, \omega t} \right)$$

with

$$A_{0} = \sum_{n=-\infty}^{+\infty} \frac{\Gamma}{\Gamma - i(\beta + 2n\omega)} \,\mathfrak{I}_{n}^{2} \left(\frac{\beta}{2\omega}\right)$$
$$A_{p} = \sum_{n=-\infty}^{+\infty} \frac{\Gamma}{\Gamma - i(\beta + 2n\omega)} \,\mathfrak{I}_{n} \left(\frac{\beta}{2\omega}\right) \mathfrak{I}_{n+p} \left(\frac{\beta}{2\omega}\right)$$
$$B_{p} = \sum_{n=-\infty}^{+\infty} \frac{\Gamma}{\Gamma - i(\beta + 2n\omega)} \,\mathfrak{I}_{n} \left(\frac{\beta}{2\omega}\right) \mathfrak{I}_{n-p} \left(\frac{\beta}{2\omega}\right).$$

In the case of ordinary coherence-resonances, the intensity of the oscillating magnetic field appears only in the argument of the Bessel functions. Here the intensity of the oscillating electric field appears not only in the Bessel functions but also in the expression  $\Gamma/\Gamma - i \ (\beta + 2n\omega)$ . Therefore it is possible to pass a resonance by varying only the intensity of the oscillating field. For the same reason, the different resonances corresponding to different values of n are not separated for ordinary values of the electric field.

When  $\beta$  is positive, only the terms corresponding to  $n \leq 0$  are resonant.

Vol. 45, 1972

Let us again calculate the intensity of the re-emitted light:

Circularly polarized light

$$\begin{aligned} I_{f1}^{\text{circ}\pm} &\propto \frac{1}{2} (1 \pm \text{Im}\,\rho) \\ &= \frac{1}{2} \left\{ 1 \pm \left( I_I + \sum_{p=1}^{\infty} I_c^p + \sum_{p=1}^{\infty} I_s^p \right) \right\} \end{aligned}$$

where

$$I_{I} = \sum_{n=-\infty}^{+\infty} \frac{x + 2ny}{1 + (x + 2ny)^{2}} \,\mathfrak{I}_{n}^{2} \left(\frac{x}{2y}\right)$$

$$I_{c}^{p} = \sum_{n=-\infty}^{+\infty} \frac{x + 2ny}{1 + (x + 2ny)^{2}} \left\{\mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n+p} \left(\frac{x}{2y}\right) + \mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n-p} \left(\frac{x}{2y}\right)\right\} \cos 2p\omega t$$

$$I_{s}^{p} = \sum_{n=-\infty}^{+\infty} \frac{1}{1 + (x + 2ny)^{2}} \left\{\mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n+p} \left(\frac{x}{2y}\right) - \mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n-p} \left(\frac{x}{2y}\right)\right\} \sin 2p\omega t$$

with  $x = \beta/\Gamma$  and  $y = \omega/\Gamma$ .



Figure 4

Intensity of the static part of the circularly polarized fluorescence light.



760

Intensity of the first harmonic (p = 1) of the circularly polarized fluorescence light (in phase term).

The re-emitted light consists of a static part  $I_I$  and a time-dependent part which can be separated into  $I_c^p$  (in phase term) and  $I_s^p$  (in quadrature term). With a lock-in type detection, it is possible to detect  $I_c^p$  or  $I_s^p$  corresponding to a selected harmonic of order p. The intensities  $I_I$ ,  $I_c^p$  and  $I_s^p$  have been computed for different values of x, y and p and some of them are given on Figures 4, 5 and 6. For y = 1, the re-emitted light  $I_I$  (Fig. 4) presents a sharp peak for x = 0, 85. The position and shape of this peak depends on the value of y. It is interesting to notice that the  $I_c^1$  term (p = 1, Fig. 5) presents at a frequency inferior to  $\Gamma(y = \omega/\Gamma = 0, 2)$  a sharp peak for a Stark separation equal to half of the natural linewidth ( $x = \beta/\Gamma = 0, 5$ ). The width is of the order of  $\Gamma$ .

Magnetic coherence resonances with the same properties could be obtained by varying the intensity of the oscillating field proportionally to the intensity of the static field.

Linearly polarized light

$$\begin{split} I_{fI}^{\text{lin}} &\propto \frac{1}{2} (1 \pm \text{Re}\,\rho) \\ &= \frac{1}{2} \left\{ 1 \pm \left( I_I + \sum_{p=1}^{\infty} \Gamma_c^p + \sum_{p=1}^{\infty} I_s^p \right) \right\} \end{split}$$





Intensity of the first harmonic (p = 1) of the circularly polarized fluorescence light (in quadrature term).

where

$$I_{I} = \sum_{n=-\infty}^{+\infty} \frac{1}{1 + (x + 2ny)^{2}} \,\mathfrak{I}_{n}^{2} \left(\frac{x}{2y}\right)$$

$$I_{c}^{p} = \sum_{n=-\infty}^{+\infty} \frac{1}{1 + (x + 2ny)^{2}} \left\{\mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n+p} \left(\frac{x}{2y}\right) + \mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n-p} \left(\frac{x}{2y}\right)\right\} \cos 2p\omega t$$

$$I_{s}^{p} = \sum_{n=-\infty}^{+\infty} \frac{x + 2ny}{1 + (x + 2ny)^{2}} \left\{\mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n+p} \left(\frac{x}{2y}\right) - \mathfrak{I}_{n} \left(\frac{x}{2y}\right) \mathfrak{I}_{n-p} \left(\frac{x}{2y}\right)\right\} \sin 2p\omega t$$

with  $x = \beta/\Gamma$  and  $y = \omega/\Gamma$ 

The intensities of  $I_I$ ,  $I_c^p$  and  $I_s^p$  have also been computed for different values of x, y and p (Figs. 7, 8 and 9). They do not seem to present very interesting properties.

## Interpretation in terms of the 'dressed atom' theory

In this theory [7] we consider the energy levels of the atoms 'dressed' with N photons. Because of the properties of the second order Stark effect, only states with  $\Delta N = 2$  have to be taken into account [8].

Coherence resonances of the kind described above occur at  $\beta = 0, 2 \omega, 4 \omega, ...$ when the pumping light puts the atoms in a superposition state  $m = 0, \pm 1$ . These resonances correspond to

$$n = 0, -1, -2, \ldots$$



Vol. 45, 1972





Intensity of the first harmonic (p = 1) of the linearly polarized fluorescence light (in quadrature term).

and we see again that if the Stark constant 'a' is positive, it is not possible to observe the resonances corresponding to n > 0.



# Conclusion

It has been shown that resonances of the coherence type can be observed in the alignment-orientation coupling process. The principal properties of these new resonances have been described. They could be applied to the measurement of lifetimes, and to the development of an electrometer. If this theory can be applied to states having a sufficiently long lifetime, weak electric fields could be detected and measured.

# Acknowledgments

I am indebted to Dr. E. Geneux and Prof. A. Quattropani for many fruitful discussions and to O. D'Auriol for the computation work.

#### REFERENCES

- [1] L. LANDAU and E. LIFCHITZ, Mir (1966).
- [2] C. COHEN-TANNOUDJI, Ann. Phys. 7, 423 (1962).
- [3] E. GENEUX, private communication (1971).
- [4] L. LOMBARDI, Thesis (Grenoble 1969).
- [5] C. COHEN-TANNOUDJI and J. DUPONT-ROC, Opt. Com. 1, No. 4, 184 (1969).
- [6] C. J. FAVRE and E. GENEUX, Phys. Lett. 8, No. 3, 190 (1964).
- [7] C. COHEN-TANNOUDJI and S. HAROCHE, C.r. Acad. Sci. Paris 262, 37 (1966).

[8] A. FAIST, Thesis (Geneva 1970).