Zeitschrift: Helvetica Physica Acta

Band: 45 (1972)

Heft: 7

Artikel: On the spectral properties of some one-particle Schrödinger

Hamiltonians

Autor: Thomas, Lawrence E.

DOI: https://doi.org/10.5169/seals-114426

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On the Spectral Properties of Some One-Particle Schrödinger Hamiltonians

by Lawrence E. Thomas

Forschungsinstitut für Mathematik, ETH, Zürich

(6. VII. 72)

Abstract. We consider a class of relatively compact perturbations $\{V\}$ of $H_0 = p_1^2 + p_2^2 + p_3^2$ acting in momentum space, $L^2(\mathbb{R}^3, d^2p)$. Resolvent matrix elements $[\phi(1/H_0 + V - z)\phi]$ are shown to be meromorphic in a neighborhood of the positive real axis, ϕ belonging to a dense set. Absolute continuity of the continuous spectrum follows.

1. Introduction

In this article we discuss spectral properties of some one-particle Schrödinger Hamiltonians. We consider a class of perturbations $\{V\}$ of $H_0 = p_1^2 + p_2^2 + p_3^2$ acting in momentum space, $L^2(\mathbb{R}^3, d^3p)$, for which the following spectral properties of $H = H_0 + V$ are shown;

- i) the absolutely continuous part of the spectrum of H and the spectrum of H_0 coincide.
- ii) the eigenvalues of H are isolated from one another except perhaps at the origin, where they may accumulate,
- iii) H has no singular continuous part.

Each of these spectral properties is probably desirable in a mathematically rigorous scattering theory. This is particularly true in the time-dependent perturbation scheme, in which one wishes to establish the existence and completeness of wave operators (defined in some canonical way), effecting a unitary transformation between H_0 and the absolutely continuous part of H[1]. Property i) is in fact a necessary condition for the existence of such operators. Properties ii) and iii) bear on the boundary value behavior of resolvent matrix elements and hence on the analytic properties of the S-matrix itself.

The perturbations considered are relatively compact, from which it follows that the essential spectra of H_0 and H coincide. Of course, there exist compact perturbations of H_0 transforming the continuous spectrum of H_0 into a discrete spectrum for H. There also exist second-order ordinary differential operators with singular continuous spectrum [2]. But by imposing additional analytic conditions on V we can rule out these pathologies and attain the above spectral properties.

We prove the above spectral properties for the class of perturbations $\{V\}$ by exhibiting a dense set of vectors \mathcal{D} for which the resolvent matrix elements $[\psi(1/H-z)\phi]$ ϕ , $\psi \in \mathcal{D}$ are meromorphic in z as z crosses the positive real axis (the essential spectrum

of H minus the origin) and travels into the second sheet. Aguilar and Combes [3] have given a proof that meromorphy of the resolvent matrix elements from a dense set implies the spectral properties. We do not repeat that argument but only show the meromorphy.

The method described here accommodates perturbations which are not necessarily short range [4], repulsive [5], spherically symmetric [6], or dilatation analytic [3]. In addition, the method is applicable to a wider class of problems, for example the description of spectral properties of multiparticle Hamiltonians and the discussion of positive bound states and resonances. These applications will be reported elsewhere.

Section 2 introduces the important notion of bounded contour distortion and discusses the resolvent meromorphy for a restricted class of perturbations (which includes some short-range potentials). Section 3 extends the results on meromorphy to perturbations (including some long-range potentials) which are limiting cases of perturbations in Section 2. Section 4 summarizes basic applications of the theory.

2. Second Sheet Continuation of Resolvent Matrix Elements

We will be working throughout in three-dimensional momentum space \mathbb{R}^3 , and three-dimensional complex space \mathbb{C}^3 . Let $\mathscr{H}=L^2(\mathbb{R}^3,d^3p)$ and let $\mathscr{D}=\{\phi\in\mathscr{H}|\phi\text{ is entire in }\mathbb{C}^3\}$. \mathscr{D} is dense in \mathscr{H} . We set $H_0=p^2=p_1^2+p_2^2+p_3^2$ and $H=H_0+V$ where V is the convolution by a function $v(\vec{p})$ with properties described below. A point in \mathbb{C}^3 (as well as in $\mathbb{R}^3\subset\mathbb{C}^3$) will be designated by \vec{p} . The complex valued function $p_1^2+p_2^2+p_3^2$ on \mathbb{C}^3 is simply written p^2 . We denote $|p_1^2|+p_2^2|+|p_3^2|$ defined on \mathbb{C}^3 by $|\vec{p}|^2$.

The convolution function $v(\vec{p})$ is assumed in this section to have the following properties:

- i) v(p) is an analytic function on an open set χ of \mathbb{C}^3 containing \mathbb{R}^3 ,
- ii) for any $\vec{p} \in \chi$ there exists a real $M(\vec{p}) \ge 0$ such that

$$\int\limits_{\mathbb{R}^3\,\cap\,\{\vec{k}||\vec{k}|>M(p)\}} v(\overrightarrow{p-k})v^*(\overrightarrow{p-k})d^3\,k<\infty.$$

Example 1. $v(\vec{p}) = \cos \alpha p/p^2 + m^2$, α a real number. For $\alpha = 0$, V is just the Yukawa potential. For $\alpha \neq 0$, $v(\vec{p})$ satisfies the above conditions but is not dilatation analytic.

Example 2. $v(\dot{p}) = \sin p^2/p^2 + m^2$. This function is cited as an example which satisfies conditions i), but not ii). Hence it will not satisfy the hypotheses of the theorem below.

Let U be a simply connected open set of the complex plane $\mathbb C.$

Definition 1: Bounded contour distortion. Let $\sigma(z, \vec{r}) : U \times \mathbb{R}^3 \to \mathbb{C}^3$ be a continuous function, and let $\sum(z)$ be the range of σ for fixed z. $\sum(z)$ is a bounded contour distortion if for fixed z

- i) σ maps \mathbb{R}^3 to $\Sigma(z)$ homeomorphically, $\Sigma(z)$ is piecewise smooth, and the (complex valued) Jacobian $\partial \sigma/\partial r = \partial(\vec{p})/\partial(\vec{r})$ is bounded and bounded away from zero almost everywhere,
- ii) there is an M(z) > 0 such that if $|\vec{r}| > M(z)$, $\sigma(z, \vec{r}) = \vec{r}$.

Theorem 1. Let $\sum(z)$ be a bounded contour distortion defined in an open set U_s intersecting the quadrant $\mathbb{C}_{++} = \{z | \text{re } z > 0, \text{ im } z > 0\}$ such that

i) for some open set $N \subset U_s \cap \mathbb{C}_{++}$, $\sum (z) = \mathbb{R}^3$, $z \in N$;

ii) for z fixed in U_s , $p^2 - z \neq 0$ for each $\vec{p} \in \Sigma(z)$ and for each $\vec{q} \in \Sigma(z)$, $v(\vec{p} - \vec{q})$ is analytic in \vec{p} for \vec{p} in a \mathbb{C}^3 neighborhood containing $\Sigma(z)$.

Then the resolvent matrix elements $[\psi(1/H-z)\phi]\phi$, $\psi \in \mathcal{D}$ may be meromorphically continued throughout U_s .

We begin the proof of this theorem by defining a family of (separable) Hilbert spaces \mathcal{H}_z , $z \in U_s$. Let \mathcal{H}_z be the space of square integrable functions defined on $\sum (z)$, with inner product

$$(\phi, \psi)_{\mathcal{H}_z} = \int\limits_{\Sigma(z)} \psi^*(\vec{p}) \phi(p) |d^3p| = \int\limits_{\mathbb{R}^3} \phi^*(\sigma(z, \vec{r})) \phi(\sigma(z, r)) \left| \frac{\partial \sigma}{\partial r} d^3r \right|.$$

 \mathcal{H}_z is just \mathcal{H} for z in $N \subset V_s \cap \mathbb{C}_{++}$.

In each \mathcal{H}_z we define the integral operator $K_z(\delta):\mathcal{H}_z\to\mathcal{H}_z$, depending on the complex variable δ , as

$$(K_z(\delta)\phi)(\vec{p}) = \int\limits_{\Sigma(z)} \frac{v(\vec{p}-\vec{q})}{q^2-z-\delta} \phi(\vec{q})d^3q.$$

(The reader should note that no absolute value signs appear around the differential form $d^3q = (\partial \sigma/\partial r)d^3r$. It is in general complex valued). It is clear that for z in the neighborhood N and $|\delta|$ sufficiently small, $K_z(\delta)$ is just $V(1/H_0 - z - \delta)$.

Lemma 1. For sufficiently small $\eta(z) > 0$, $K_z(\delta)$ is compact analytic, $|\delta| < \eta(z)$.

Proof: Choose $\eta(z) = \frac{1}{2} \min |q^2 - z|$. Then $q^2 - z - \delta \neq 0$, $\vec{q} \in \sum(z)$, and $K_z(\delta)$ is Hilbert–Schmidt since

$$\int\limits_{\Sigma(z)\times\Sigma(z)}\left|\frac{v(\dot{p}-\dot{q})}{q^2-z-\delta}\right|^2|d^3p\,d^3q|<\infty$$

by the definition of $\sum(z)$, assumptions ii) of the theorem and i) on v. K clearly depends analytically on δ .

We next introduce a linear mapping $A_{wz}^c: \mathcal{H}_z \to \mathcal{H}_w$, $z, w \in U_s$. Let c be a smooth curve running from z to w in U_s . Let $\mathcal{D}(A_{wz}^c) = \{\phi \in \mathcal{H}_z | \exists \ a \ \mathbb{C}^3 \text{ neighborhood } W \text{ containing } \bigcup_{x \in c} \sum(x) \text{ and } \phi \text{ is analytic in } W\}$. Then we define $A_{wz}^c \phi = \phi|_{\Sigma(w)}$. Hence A_{wz}^c is analytic continuation of ϕ from $\sum(z)$ to $\sum(w)$. Note that $A_{wz}^{c-1} = A_{zw}^c$ and that this inverse is defined on the range of A_{wz}^c . If x is a point on the curve c, we have $A_{zw}^c = A_{zx}^c A_{xw}^c$, for elements $\phi \in \mathcal{D}(A_{zw}^c)$.

Let z be a point in U_s and let θ_z be the connected part of $\{z' \in U_s | |z' - z| < \eta(z)\}$, $\eta(z)$ the same as in Lemma 1.

Lemma 2. For $z + \delta \in \theta_z$ and any path c from z to $z + \delta$ lying in θ_z , $K_z(\delta)\phi = A_{z,z+\delta}^c K_{z+\delta}(0) A_{z+\delta,z}^c \phi$, $\phi \in \mathcal{D}(A_{z+\delta,z}^c)$.

Proof: We have

$$(K_z(\delta)\phi)|_{\Sigma(z)} = \int\limits_{\Sigma(z)} \frac{v(\vec{p}-\vec{q})}{q^2-z-\delta} \,\phi(\vec{q})d^3q.$$

By condition ii) of the theorem, we may choose a complex $z_1 \in c$, $z_1 - z \neq 0$ such that $v(\vec{p} - \vec{q})\phi(\vec{q})$ is nonsingular for \vec{p} , \vec{q} ranging independently over a \mathbb{C}^3 neighborhood containing $\bigcup_{x \in I_1} \sum (x)$, where I_1 is the interval on c from z to z_1 . $[v(\vec{p} - \vec{q})/q^2 - z - \delta]\phi(\vec{q})d^3q$ is an analytic closed differential form in \vec{q} on this neighborhood. Using the complex form of Stokes' theorem [7], we may replace the integration path $\sum (z)$ of the above integral by $\sum (z_1)$ to get

$$(K_{z}(\delta)\phi)|_{\Sigma(z)} = \int_{\Sigma(z_{1})} \frac{v(\vec{p} - \vec{q})}{q^{2} - z - \delta} \phi(\vec{q}) d^{3} q = A_{z,z_{1}}^{c} K_{z_{1}}(\delta - z_{1} + z) A_{z_{1},z}^{c} (\phi|_{\Sigma(z)}).$$

(Note that because $\sum(x)$ is a bounded contour distortion, $\sum(z_1) - \sum(z)$ is compact. $\sum(z_1) - \sum(z)$ may be regarded as the boundary of a four-dimensional region in the domain of analyticity of the differential form. This allows application of Stokes' theorem.) We next choose $z_2 \in c$ such that $v(\vec{p} - \vec{q})\phi(\vec{q})$ is nonsingular, \vec{p} , \vec{q} ranging independently over a neighborhood of $\bigcup_{x \in I_2} \sum(x)$, I_2 the portion of c from z_1 to z_2 . It follows in a similar manner that

$$(K_{z_1}(\delta-z_1+z)\phi)|_{\Sigma(z_1)}=A^c_{z_1,z_2}K_{z_2}(\delta-z_2+z)A^c_{z_2,z}(\phi|_{\Sigma(z_1)}).$$

Combining this equation with the previous one, we get

$$(K_{\boldsymbol{z}}(\delta)\phi)|_{\boldsymbol{\Sigma}(\boldsymbol{z})} = A_{\boldsymbol{z},\boldsymbol{z}_2}^{\,\boldsymbol{c}} K_{\boldsymbol{z}_2}(\delta - \boldsymbol{z}_2 + \boldsymbol{z}) A_{\boldsymbol{z}_2,\boldsymbol{z}}^{\,\boldsymbol{c}}(\phi|_{\boldsymbol{\Sigma}(\boldsymbol{z})}).$$

By repeated application of this process, a finite set z_1, z_2, \ldots, z_k can be obtained such that $z_k = z + \delta$, and

$$K_{z}(\delta)\phi = A_{z,z+\delta}^{c} K_{z+\delta}(0) A_{z+\delta,z}^{c} \phi.$$

Only a finite number of z_j 's are required since otherwise one could conclude the existence of a point $z_s \in c$ such that $v(\vec{p} - \vec{q})$ would be singular for \vec{p} , \vec{q} ranging over $\sum (z_s)$.

Lemma 3. Let $\phi \in \mathcal{D}$ and let ψ be a solution to the integral equation

$$\Psi + K_z(\delta)\psi = \phi|_{\Sigma(z)}, \quad z + \delta \in \theta_z.$$

Then $\psi \in \mathcal{D}(A_{z+\delta,z}^c)$ and $\psi|_{\Sigma(z+\delta)} = A_{z+\delta,z}^c \psi$ satisfies

$$\psi|_{\Sigma(z+\delta)} + K_{z+\delta}(0)(\psi|_{\Sigma(z+\delta)}) = \phi|_{\Sigma(z+\delta)}$$

where c is any path in θ_z from z to $z + \delta$.

Proof: The proof of this lemma closely resembles that of Lemma 2. Condition ii) of the theorem and the entirety of $\phi \in \mathcal{D}$ imply the existence of a $z_1 \in c$, $z_1 - z \neq 0$, such

that

$$\psi(\vec{p}) = -\int\limits_{\Sigma(z)} \frac{v(\vec{p}-\vec{q})}{q^2-z-\delta} \, \psi(\vec{q}) d^3 \, q + \phi(\vec{p})$$

is analytic in a \mathbb{C}^3 neighborhood of $\bigcup_{x \in I_1} \sum (x)$, I_1 the interval of c from z to z_1 . Using Lemma 2 we may then write

$$\begin{split} \psi|_{\Sigma^{(z_1)}} &= A^c_{z_1,z} \, K_z(\delta) \psi + \phi|_{\Sigma^{(z_1)}} \\ &= K_{z_1} (\delta - z_1 + z) (\psi_{\Sigma^{(z_1)}}) + \phi|_{\Sigma^{(z_1)}}. \end{split}$$

We next choose a $z_2 \in c$ such that $\psi(\not p)$ is analytic in a neighborhood of $\bigcup_{x \in I_2} \sum (x)$, I_2 the interval of c from z_1 to z_2 . Again by repeated application of this process we can obtain a finite set $z_1, z_2, \ldots, z_k, z_k = z + \delta$ and

$$\psi|_{\Sigma^{(z+\delta)}} + K_{z+\delta}(0)(\psi|_{\Sigma^{(z+\delta)}}) = \phi|_{\Sigma^{(z+\delta)}}.$$

Only a finite number of z_j 's are required since otherwise there would be a $z_s \in c$ such that $\psi(\vec{p})$ would be singular on $\sum (z_s)$, and yet nonsingular on $\sum (z_s - \rho)$, $z_s - \rho \in c$, $\rho \neq 0$. But this is impossible since ψ has the representation

$$\psi(\vec{p}) = -\int\limits_{\Sigma(z_s-\rho)} \frac{v(\vec{p}-\vec{q})}{q^2-z-\delta} \, \psi|_{\Sigma(z_s-\rho)}(\vec{q}) d^3 \, q + \phi(\vec{p})$$

which for $|\rho|$ sufficiently small surely is analytic in a \mathbb{C}^3 neighborhood of $\sum (z_s)$. Since ψ is analytic in a neighborhood of $\sum (x)$ for any $x \in \theta_z$, the analytic continuation of ψ to $\psi|_{\sum (z+\delta)}$ is path independent.

We are now able to prove Theorem 1. We show that the meromorphic continuation of $[\psi(1/H-z)\phi]$ ϕ , $\psi \in \mathcal{D}$ throughout U_s is given by

$$\mathcal{M}(z) = \int_{\Sigma(z)} (\psi^*|_{\Sigma(z)})(\vec{p}) \frac{1}{\vec{p}^2 - z} (1 + K_z(0))^{-1} (\phi|_{\Sigma(z)})(\vec{p}) d^3 p.$$

 $(\psi^*|_{\Sigma^{(z)}})$ is the analytic continuation of ψ^* from \mathbb{R}^3 to $\Sigma^{(z)}$.) First note that if $z \in N \subset U_s \cap \mathbb{C}_{++}$, the integral expression on the right-hand side is

$$\mathcal{M}(z) = \int_{\mathbb{R}^3} \psi^*(p) \frac{1}{H_0 - z} \left(1 + V \frac{1}{H_0 - z} \right)^{-1} \phi(\vec{p}) d^3 p = [\psi(1/H - z)\phi]_{\mathcal{H}},$$

i.e., $\mathcal{M}(z)$ is the resolvent matrix element for z in N. It remains only to check the meromorphy of $\mathcal{M}(z')$, $z' \in \theta_z$, $z \in U_s$. By Lemma 3 the integrand of $\mathcal{M}(z')$ may be analytically continued from $\sum (z')$ to $\sum (z)$. Applying Stokes' theorem as in Lemma 2, we obtain

$$\begin{split} \mathcal{M}(z') &= \int\limits_{\Sigma(z')} (\psi^*|_{\Sigma(z')}) (\vec{p}) \frac{1}{p^2 - z'} (1 + K_{z'}(0))^{-1} \phi|_{\Sigma(z')} (\vec{p}) d^3 p \\ &= \int\limits_{\Sigma(z)} (\psi^*|_{\Sigma(z)}) (\vec{p}) \frac{1}{p^2 - z'} (A_{zz'} (1 + K_{z'}(0))^{-1} \phi|_{\Sigma(z')}) (\vec{p}) d^3 p \\ &= \int\limits_{\Sigma(z)} (\psi^*|_{\Sigma(z)}) (\vec{p}) \frac{1}{p^2 - z'} (1 + K_{z}(z' - z))^{-1} \phi|_{\Sigma(z)} (\vec{p}) d^3 p. \end{split}$$

Lawrence E. Thomas H. P. A.

But from Lemma 1, $K_z(z'-z)$ is compact and analytic in z'. Hence the latter expression is meromorphic in θ_z [8]. This completes the proof.

Example 3. $v(\vec{p}) = \cos \alpha p/p^2 + m^2$, α a real number. We discuss the meromorphy domains of the resolvent in two cases by constructing bounded contour distortions. In both cases z starts from a neighborhood $N \subset \mathbb{C}_{++}$, crosses over the positive real axis and travels into the second sheet.

Case 1. The matrix elements of the resolvent $[\psi(1/H-z)\phi]$ ϕ , $\psi \in \mathcal{D}$ will be meromorphic in the second sheet for im $\sqrt{z} > -\frac{1}{2}m$, $z \neq 0$. In the complex plane \mathbb{C} , let $S_z(t)$ $0 \leq t < \infty$ be a simple smooth curve depending continuously on z which originates at the origin, avoids the points $\pm \sqrt{z}$, and lies in the strip $|\text{im } x| < \frac{1}{2}m$. In addition, let the locus of $S_z(t)$ be the positive real axis for all but a finite part of the curve, and for a neighborhood $N \subset \mathbb{C}_{++}$, let the locus be the entire positive real axis, $z \in N$. $S_z(t)$ is parameterized in such a way that $S_z(t) = t$ for t sufficiently large. Then the mapping $\sigma(z,\vec{r})$ for $\Sigma(z)$ is given by $\sigma(z,\vec{r}) = S_z(r)(\vec{r}/r)$. One can verify that $\Sigma(z)$ satisfies the conditions of Theorem 1. In particular, for \vec{p} , $\vec{q} \in \Sigma(z)$, $p^2 - z \neq 0$, and $v(\vec{p} - \vec{q})$ is analytic for \vec{p} in a neighborhood of $\Sigma(z)$, $\vec{q} \in \Sigma(z)$.

Case 2. The matrix elements of the resolvent $[\psi(1/H-z)\phi] \phi, \psi \in \mathcal{D}$ will be meromorphic in the second sheet region $\arg z > -\pi/2$. Let x(z) be a point in $\mathbb C$ depending continuously on z which lies on the positive real axis for z in a neighborhood $N \subset \mathbb C_{++}$, and otherwise lies on the vertical line $\operatorname{Re} x = \operatorname{Re} \sqrt{z}$, $-\operatorname{Re} \sqrt{z} < \operatorname{im} x(z) < \operatorname{im} \sqrt{z}$. Let $S_z(t)$ $0 \le t < \infty$ be the piecewise smooth curve with the locus of points consisting of the three straight line segments, [0, x(z)], $[x(z), 2\operatorname{Re} \sqrt{z}]$, $[2\operatorname{Re} \sqrt{z}, +\infty]$ in $\mathbb C$. Again assume $S_z(t) = t$ for t sufficiently large. (Note that $S_z(t)$ is so constructed that for any two points $x_1, x_2 \in S_z(t)$, $|\operatorname{re}(x_1 - x_2)| > |\operatorname{im}(x_1 - x_2)|$.) Then the mapping $\sigma(z, \tilde{r})$ for $\Sigma(z)$ is $\sigma(z, \tilde{r}) = S_z(r)(\tilde{r}/r)$. Again one can verify that $\Sigma(z)$ satisfies the conditions of Theorem 1.

3. Continuation of Resolvent Matrix Elements for Long-Range Potentials

The results in the previous section concerning the meromorphy of resolvent matrix elements may be extended to a larger class of perturbations. This class consists of potentials which are limits, in a sense defined below, of potentials considered in Theorem 1. The class includes certain long-range potentials.

Let V_n , $n=1, 2, \ldots$ be a sequence of potentials with corresponding convolution functions $v_n(\vec{p})$ and assume the v_n satisfy the conditions given in Section 2. Let V be a potential with convolution function $v(\vec{p})$.

Theorem 2. Suppose

- i) there is a bounded contour distortion $\sum(z)$, independent of n, defined throughout an open neighborhood U_s satisfying the conditions of Theorem 1 for each $v_n(\vec{p})$;
- ii) the V_n converge to V in the sense that the integral operators $K_{nz}:\mathcal{H}_z\to\mathcal{H}_z$, $K_z:\mathcal{H}_z\to\mathcal{H}_z$,

$$(K_{nz}(\delta)\phi)(\vec{p}) = \int_{\Sigma(z)} \frac{v_n(\vec{p}-\vec{q})}{q^2-z-\delta} \phi(q)d^3q,$$

Vol. 45, 1972 On the Spectral Properties of Some One-Particle Schrödinger Hamiltonians 1063

$$(K_z(\delta)\phi)(\vec{p}) = \int\limits_{\Sigma(z)} rac{v(\vec{p}-\vec{q})}{q^2-z-\delta} \, \phi(q) d^3 \, q$$

satisfy

$$\lim_{n\to\infty} K_{nz}(0) \to \mathrm{K}_z(0)$$

in norm, uniformly in z.

Then $[\psi(1/H_0 + V - z)\phi] \phi$, $\psi \in \mathcal{D}$ can be meromorphically continued throughout U_s .

Proof: Again let θ_z be the connected neighborhood of z defined above Lemma 2 in Section 2. $K_z(0)$ is compact since it is the limit in norm of compact operators. $K_z(\delta)$ is compact analytic in δ , $z + \delta \in \theta_z$ since it can be written as the composition of a bounded analytic (multiplication) operator and a compact operator,

$$K_{\mathbf{z}}(\delta)\phi(p) = \int_{\Sigma(\mathbf{z})} \frac{v(\vec{p}-\vec{q})}{(q^2-z)} \frac{1}{[1-(\delta/q^2-z)]} \phi(\vec{q})d^3q.$$

Note that $K_{nz}(\delta)$ converges uniformly to $K_z(\delta)$, $z + \delta \in \theta$. Now set

$$\mathscr{M}(z) = \int\limits_{\Sigma(z)} \psi^*|_{\Sigma(z)}(\vec{p}) \frac{1}{\vec{p}^2 - z} (1 + K_z(0))^{-1} \phi(\vec{p}) d^3 p, \quad \phi, \psi \in \mathscr{D}.$$

For z in N, $\mathcal{M}(z)$ is just equal to $[\psi(1/H-z)\phi]$. $\mathcal{M}(z')$ is meromorphic about the point $z,z'\in\theta_z$ because

$$\mathcal{M}(z') = \lim_{n \to \infty} \int_{\Sigma(z')} \psi^* |_{\Sigma(z')}(\vec{p}) \frac{1}{p^2 - z'} (1 + K_{nz}(0))^{-1} \phi |_{\Sigma(z')}(\vec{p}) d^3 p$$

$$= \lim_{n \to \infty} \int_{\Sigma(z)} \psi^* |_{\Sigma(z)}(\vec{p}) \frac{1}{p^2 - z'} (1 + K_{nz}(z' - z))^{-1} \phi |_{\Sigma(z)}(\vec{p}) d^3 p$$

$$= \int_{\Sigma(z)} \psi^* |_{\Sigma(z)}(\vec{p}) \frac{1}{p^2 - z'} (1 + K_z(z' - z))^{-1} \phi |_{\Sigma(z)}(\vec{p}) d^3 p.$$

The latter expression is meromorphic in z' since $K_z(z'-z)$ is compact analytic. This proves the theorem.

Example 4. $v = \frac{\cos \alpha p}{p^2}$, $v_n = \frac{\cos \alpha p}{p^2 + (1/n)}$, α a nonnegative real number. In configu-

ration space, $V\alpha(r)$ is

Lawrence E. Thomas H. P. A.

Case 2 of example 3 in the previous section provides a contour distortion $\sum(z)$ in the region $R_2 = \{z \in \mathbb{C} | \arg z > -(\pi/2)\}$ for which all v_n satisfy the conditions of Theorem 1. (Note that in Case 2, the contour distortion did not depend on m.) To establish the meromorphy of the matrix elements $[\psi(1/H_0 + V - z)\phi] \phi, \psi \in \mathcal{D}$ in R_2 , one must show the uniform convergence $K_{nz} \to K_z$ for z in any compact neighborhood $B \subseteq R_2$. We show only the boundedness of K_z in B, by writing K_z as the sum of two operators, $K_z = K_z^1 + K_z^2$,

$$\begin{split} K_{z} \phi(p) &= \int\limits_{\Sigma(z)} \frac{\cos \alpha (p-q)}{(p-q)^{2}} \frac{1}{q^{2}-z} \phi(\vec{q}) d^{3} q \\ &= \int\limits_{\Sigma(z) \ \cap \ (\vec{q}||\vec{p}-\vec{q}| \leqslant M)} \frac{\cos \alpha (p-q)}{(p-q)^{2}} \frac{1}{q^{2}-z} \phi(q) d^{3} q \\ &+ \int\limits_{\Sigma(z) \ \cap \ \{\vec{q}||\vec{p}-\vec{q}| \geqslant M\}} \frac{\cos \alpha (p-q)}{(p-q)^{2}} \frac{1}{q^{2}-z} \phi(q) d^{3} q, \end{split}$$

where M is an arbitrary positive constant. The first term is bounded since the integration has kernel satisfying the Holmgren criteria for boundedness of the operation. (Namely, if

$$T\psi = \int K(x,y)\psi(y)d\mu(y), \ s_1 = \sup_{x} \int |K(x,y)|d\mu(y), \ s_2 = \sup_{y} \int |K(x,y)|d\mu(x),$$

then $|T| \leqslant (s_1 s_2)^{1/2}$ [9].) The second operation is bounded since it is Hilbert–Schmidt. The uniform convergence $K_{nz} \to K_z$ may be similarly demonstrated by breaking up the path of integration for the operator $(K_z - K_{nz})$ into the two parts again and showing the uniform convergence of the $K_z^1 - K_{nz}^1$ and $K_z^2 - K_{nz}^2$ separately.

4. Concluding Remarks

In this section we make some remarks concerning conditions for V in configuration space in order that the convolution function v for V in momentum space permit applications of Theorem 1 or 2, for z in a neighborhood of the positive real axis. If V is multiplication by an L^2 -function of compact support, then V is convolution by an entire function in momentum space. Theorem 1 may be applied in this case to show that the resolvent matrix elements of \mathscr{D} are meromorphic on an (infinitely sheeted, in general) Riemann surface $\{z \mid -\infty < \arg z < \infty, z \neq 0\}$. If V is multiplication by an L^2 -function w such that $\int w \, e^{m|r|} d^3 r < \infty$ for some m > 0, V will be convolution by a function u analytic in the region im $|\vec{p}| < m$. Theorem 1 will give meromorphy of the resolvent matrix elements in the region $\{z \in \mathbb{C} | \lim \sqrt{z}| < m/2, z \neq 0\}$. This latter result is that of Dolph, McLeod and Thoe [4]. Theorem 2 and the example following it show resolvent meromorphy in a neighborhood of the positive real axis for V_{α} multiplication by

$$w_{\alpha} = \begin{pmatrix} 0 & r < \alpha \\ \\ \frac{1}{r} & r > \alpha \end{pmatrix}$$

in configuration space. One can show as well resolvent meromorphy in a neighborhood of the positive real axis for $V = V_1 + V_2$ where V_1 is multiplication by a function

$$w^1 \in L^2(\mathbb{R}^3), \quad \int w^1 e^{m|r|} d^3r < \infty,$$

and V_2 is multiplication by

$$w^{2}(\vec{r}) = \sum_{i=1}^{N} a_{i} w_{\alpha^{i}}(\overrightarrow{r-r_{0}^{i}}),$$

 a_i real, $w_{\alpha i}$ defined above. Considerably more general conditions on V in configuration space can be given, so that the convolution function v has appropriate analytic properties in momentum space for application of Theorem 2. The proof of the sufficiency of these conditions, however, requires a rather detailed examination of the Fourier transform of the potential and so we do not describe the conditions here.

Acknowledgments

I am indebted to Dr. W. G. Faris, Battelle Advanced Studies Center, Geneva, for many helpful discussions. The work was completed while the author was a guest at the Forschungsinstitut für Mathematik, ETH, Zürich.

REFERENCES

- [1] T. Kato, Perturbation Theory for Linear Operators, chap. X (Springer-Verlag, New York Inc., 1966).
- [2] T. Kato, Perturbation Theory for Linear Operators (Springer-Verlag, New York Inc., 1966), pp. 244, 518, 523.
- [3] J. AGUILAR and J. M. COMBES, Comm. Math. Phys. 22, 269 (1971).
- [4] C. L. Dolph, J. B. McLeod and D. Thoe, J. math. Analysis Applic. 16, 311 (1966).
- [5] R. LAVINE, Comm. Math. Phys. 20, 301 (1971).
- [6] J. WEIDMANN, Math. Z. 98, 268 (1967).
- [7] V. S. VLADIMIROV, Methods of the Theory of Functions of Many Complex Variables (MIT Press, Cambridge 1966), p. 198.
- [8] N. DUNFORD and J. T. Schwartz, *Linear Operators*, part I (Interscience Publishers, Inc., New York 1958), p. 592.
- [9] K. O. FRIEDRICHS, Perturbation of Spectra in Hilbert Space (American Mathematical Society, Providence, 1965), p. 107.