

<b>Zeitschrift:</b>	Helvetica Physica Acta
<b>Band:</b>	49 (1976)
<b>Heft:</b>	6
<b>Rubrik:</b>	Zusammenfassungen der letzten eingegangenen Arbeiten = Résumés des derniers articles reçus

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**HELVETICA PHYSICA ACTA**  
**Zusammenfassungen der letzten eingegangenen Arbeiten**  
**Résumés des derniers articles reçus**

---

**Distinguished Self-adjoint Extension for Dirac Operator with Potential Dominated by Multicenter Coulomb Potentials**

by G. NENCIU

Institut of Atomic Physics, Bucharest, Romania

(20. V. 1976)

*Abstract.* The existence and the uniqueness of the distinguished self-adjoint extension of the Dirac operator describing an electron in the field of a finite number of point charges with  $Z < 137$  is proved.

**Bremsstrahlung and Čerenkov Radiation of High Energy Particles in an Excited Medium**

by S. P. ANDREIEV<sup>1)</sup>

Département de Physique Théorique, Université de Genève, CH-1211 GENEVE 4 – Switzerland

(21. V. 1976)

*Abstract.* Bremsstrahlung and Čerenkov radiation of a high energy particle in a homogeneous medium excited by a resonant electromagnetic field are investigated. It is shown that the field greatly influences the radiation by the particle moving in the medium. The field leads to a suppression of Čerenkov radiation. In some cases the presence of the field leads to an increase of Bremsstrahlung. The intensities of both Bremsstrahlung and Čerenkov radiation are shown to depend non-linearly on the intensity of the field.

**Bestimmung der spektroskopischen Faktoren mehrerer Energieniveaux der Isotope Ni<sup>59, 61, 63, 65</sup> mit (*d, p*) – Reaktionen bei 2,8 MeV Deuteronenenergie**

von P. STAUB, E. BAUMGARTNER, J. X. SALADIN,<sup>1)</sup> H. SCHÄR und D. TRAUTMANN

Institut für Physik der Universität Basel

(24. V. 1976)

*Abstract.* The differential cross section of Ni<sup>58</sup>(*d, p*)Ni<sup>59</sup>, Ni<sup>60</sup>(*d, p*)Ni<sup>61</sup>, Ni<sup>62</sup>(*d, p*)Ni<sup>63</sup> and Ni<sup>64</sup>(*d, p*)Ni<sup>65</sup> to different excitation states of the final nuclei have been measured at a deuteron energy of 2.8 MeV at angles between 50 and 160 degrees. The experimental differential cross sections were compared with theoretical curves. The spectroscopic factors for 76 different states of the product nuclides were determined.

**Untersuchungen zur Bildung von Tracks in Kristallen**

von A. SIGRIST und R. BALZER

Laboratorium für Kernphysik, ETH, 8093 Zürich

(28. V. 1976)

*Abstract.* The minimal energy loss  $(dE/d\xi)_k$  [MeV · cm<sup>2</sup> · mg<sup>-1</sup>] for the formation of a track in an insulator has been determined for mica, tourmaline, quartz glass, quartz crystal, lithium niobate and beryl. The investigations show that the  $(dE/d\xi)_k$ -value of a track detector can be roughly estimated from the thermal conductivity of the given material.

**Modèle Semiphénoménologique de l'Interaction Nucléon – Nucléon**

par A. HOURIET et X. BAGNOUD

Institut de Physique Théorique, Université de Fribourg

(2. VI. 1976)

*Abstract.* A nucleon with isobars is used to elaborate a model of the nucleon–nucleon interaction at low energy ( $E_{CM} < 160$  MeV). Three free parameters only are introduced among which  $f_r^2$ , the pion-nucleon renormalized coupling constant. The model establishes a very good coordination for deuton and  $p-p$  scattering-polarization measurements ( ${}^1K_0$ ,  ${}^1D_2$ ,  ${}^1G_4$  phase shifts), and permits the determination of  $f_r^2$  for every independent experimental value. For 21 such values, we obtain the mean value  $\langle f_r^2 \rangle = 0.0785$  with  $\Delta f_r^2 = 0.0024$  (3%).

**Bestimmung spektroskopischer Faktoren für Energieniveaux der Isotope  ${}^{41, 43, 45, 49}\text{Ca}$  mit  $(d,p)$  – Reaktionen bei 2,5 MeV Deuteronenenergie**

von H. SCHÄR, D. TRAUTMANN und E. BAUMGARTNER

(Institut für Physik der Universität Basel)

(2. VI. 1976)

*Abstract.* Differential  $(d, p)$ -cross sections on  ${}^{40}\text{Ca}$ ,  ${}^{42}\text{Ca}$ ,  ${}^{44}\text{Ca}$ ,  ${}^{48}\text{Ca}$  at 2.5 MeV incident deuteron energy are measured in 5 degree intervals between 40 and 160 degrees. The angular distributions are compared with distorted-wave Born approximation calculations, and the spectroscopic factors are extracted for a number of transitions.

**Spin Corrections to the Two-Body Eikonal Amplitude<sup>1</sup>)**

by M. QUIROS<sup>2</sup>)

University of Geneva

(22. VI. 1976)

*Abstract.* Using the one-parameter eikonal representation and  $\alpha$ -space techniques developed in an earlier paper, the first-order spin corrections are obtained and its high-energy behaviour proved to be non-negligible. In the limit where the range of interaction goes to infinity, bound-states, in the electron positron annihilation region, appear as poles in the  $s$ -channel and Regge trajectories in the  $t$ -channel, as in the usual eikonal model. Spin corrections are associated with double poles.

The Cooperative Institute for Research in Environmental Sciences (Cires) offers one year Visiting Fellowships to scientists with research interests in the areas of -

ATMOSPHERIC SCIENCE

SOLID EARTH SCIENCE

ATMOSPHERIC AND TERRESTRIAL E.M. WAVE PROPAGATION

PALEO AND ROCK MAGNETISM

Recipients of these awards are completely free to pursue their own research programs. Selection is made on the likelihood of an active exchange of ideas between the Visiting Fellow and the scientists in Cires and other local research groups. Stipend is scaled to research experience, and the program is open to scientists of all countries. Applications for the 1977-78 academic year awards should be sent to:

The Director  
Visiting Fellows Program  
Cires

University of Colorado  
Boulder, Colorado 80302 U.S.A.

before February 1, 1977 and should include a curriculum vitae, publications list, the names of three professional references and a brief outline of the intended research. Cires is jointly sponsored by the University of Colorado and the Environmental Research Laboratories, National Oceanic and Atmospheric Administration.

