

Zeitschrift: Helvetica Physica Acta
Band: 56 (1983)
Heft: 1-3

Artikel: Localization and interaction in one dimension
Autor: Apel, W. / Rice, T.M.
DOI: <https://doi.org/10.5169/seals-115357>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LOCALIZATION AND INTERACTION IN ONE DIMENSION

W. Apel and T.M. Rice

Theoretische Physik

ETH Hönggerberg
CH 8093 Zürich

Abstract:

A new approach is given to the problem of an interacting impure one-dimensional electronic system. First the inelastic mean free path $l_{in}(T)$, which results from collisions with other electrons is calculated. The elastic mean free path $l_{el}(T)$ due to impurity scattering is obtained from the resistivity of an infinitesimal short sample. Following Thouless /1/ we argue that the impurity scattering is classical when $l_{in}(T) \ll l_{el}(T)$. From the temperature dependence of the mean free path's we conclude that for $2g_2 - g_1 < -\frac{3}{5} 2\pi v_f$ and $g_1 > 0$ (excluding the Peierls transition) this condition is satisfied at sufficiently low temperature and as a result the resistance is proportional to the length of the sample. The value of the resistivity vanishes in the limit $T \rightarrow 0$ due to the complete screening of the impurity potential. In the other regime, $l_{in}(T) \gg l_{el}(T)$ and the impurity scattering must be treated quantum mechanically. The resistance diverges exponentially with a positive power (depending on the interaction) of the length. The results of the present approach agree with those obtained previously using a scaling theory /2/ and lead to a simpler interpretation. A more detailed account of this approach will appear elsewhere /3/.

References:

1. D.J. Thouless: Solid State Commun. 34, 683 (1980)
2. W. Apel, T.M. Rice: Phys. Rev. B26, 7063 (1982)
3. W. Apel, T.M. Rice: J. Phys. C in press