

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 56 (1983)  
**Heft:** 6

**Artikel:** On finite volume corrections to the equation of state of a free Bose gas  
**Autor:** Berg, M. van den  
**DOI:** <https://doi.org/10.5169/seals-115441>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# On finite volume corrections to the equation of state of a free Bose gas

By M. van den Berg, Dublin Institute for Advanced Studies,  
10 Burlington Road, Dublin 4, Ireland

(6. VI. 1983; rev. 16. VII. 1983)

*Abstract.* We calculate and discuss the asymptotic behaviour of the finite volume correction term to the equation of state of a free Bose gas in the bulk limit.

## 1. Introduction

In studying a phase-transition one is faced with an apparent dilemma: the phase-transition manifests itself in a mathematically clean fashion as a singularity in a thermodynamic function only in the thermodynamic limit, in which the volume as well as the number of particles is infinite. On the other hand, any practical detection of a phase-transition makes use of a sample consisting of a finite number of molecules in a finite volume. The standard reply to such an objection to the use of the thermodynamic limit is that this procedure yields the first term in an asymptotic expansion of the thermodynamic functions. There is no doubt that this is true in general; nevertheless it might prove troublesome to demonstrate this in any particular case. The calculation of the correction term would become important in a critical examination of experimental data near a critical point.

For example: is the observed singularity in the specific heat (the so called  $\lambda$ -singularity) [1–3] in liquid helium due to a weak external gravitation field?

In this paper we will examine the finite volume corrections to the equation of state for a free Bose gas. So we will neglect interaction between the particles. However, we have shown elsewhere [4] that the behaviour of the free gas pressure controls the phase-transition in the mean-field model. We expect this also to be true in the interacting gas.

In studying the finite volume correction to the equation of state for a free Bose gas previous workers [5, 6] have studied the grand canonical pressure at fixed chemical potential; this approach runs into difficulties near the critical density. It is necessary to study it at fixed mean density. We now formulate the problem and state the results.

Consider a free boson gas in a  $d$ -dimensional convex region  $B$  in euclidean space with volume  $V(B)$  and surface area  $S(B)$ . For the single particle hamiltonian  $H(B)$  we take  $-\Delta/2$  with Dirichlet boundary conditions on the boundary  $\partial B$  of  $B$ . The equation of state is given in the implicit form: The grand canonical

pressure  $p_B(\rho)$  is given by

$$p_B(\rho) = \frac{1}{V(B)} \sum_{n=1}^{\infty} \frac{(z(B; \rho))^n}{n} \text{trace}(e^{n \Delta/2}), \quad (1)$$

where  $z(B; \rho)$  is the unique positive solution of

$$\rho = \frac{1}{V(B)} \sum_{n=1}^{\infty} (z(B; \rho))^n \text{trace}(e^{n \Delta/2}); \quad (2)$$

$\rho$  is the mean particle density in the grand canonical ensemble. In the thermodynamic limit in which we keep  $\rho$  fixed and in which we take for  $B$  a sequence  $B_l$  ( $B_1 \subset B_2 \subset B_3 \dots$ ) such that  $S(B_l)/V(B_l) \rightarrow 0$  one can prove [9] that

$$\lim_{l \rightarrow \infty} p_{B_l}(\rho) = p(\rho) = \sum_{n=1}^{\infty} \frac{(\zeta(\rho))^n}{n(2\pi n)^{d/2}}, \quad (3)$$

where

$$\zeta(\rho) = \begin{cases} \zeta, & \rho < \rho_c \\ 1, & \rho \geq \rho_c \end{cases} \quad (4)$$

and  $\zeta$  is the unique solution in  $[0, 1]$  of

$$\rho = \sum_{n=1}^{\infty} \frac{\zeta^n}{(2\pi n)^{d/2}}. \quad (5)$$

We will only consider cases where the critical density  $\rho_c$  is finite ( $d = 3, 4, \dots$ ):

$$\rho_c = \sum_{n=1}^{\infty} \frac{1}{(2\pi n)^{d/2}}, \quad d = 3, 4, \dots \quad (6)$$

Clearly the right hand side of (3) is the first term of an asymptotic expansion of  $p_{B_l}(\rho)$ . In order to find the second term we have to solve equation (2) for  $z(B_l; \rho)$  and substitute the value into (1). In previous papers [9–11] we have shown that there exist different subsequences  $B_l$  which lead to different asymptotic behaviour of  $z(B_l; \rho)$  (for  $\rho \geq \rho_c$ ). That is the reason that the condensate has different structures for different subsequences (that was overlooked by previous workers [7, 8]). We pick out one particular subsequence. Let  $B_l$  be the dilation of a convex region  $B_1$  with unit volume:

$$B_l = \left\{ x \in R^d : \frac{x}{l} \in B_1 \right\}, \quad (7)$$

so that in particular

$$S_l = S(B_l) = l^{d-1} S(B_1), \quad (8)$$

$$V_l = V(B_l) = l^d, \quad (9)$$

$$E_k^l = E_k(B_l) = l^{-2} E_k(B_1), \quad (10)$$

where  $E_1(B_1) < E_2(B_1) \leq E_3(B_1) \leq \dots$  are the eigenvalues of  $-\Delta/2$  with Dirichlet boundary conditions on  $\partial B_1$ . We will also assume that the curvature at each point of  $\partial B_1$  is bounded from above by  $1/R_1$  ( $R_1 > 0$ ). It is convenient to introduce a

scaled fugacity:

$$\zeta_l(\rho) = e^{-E_l} z(B_l; \rho), \quad (11)$$

so that (1) and (2) can be rewritten as follows:

$$p_l(\rho) = p_{B_l}(\rho) = \frac{1}{l^d} \sum_{n=1}^{\infty} \frac{(\zeta_l(\rho))^n}{n} \sum_{k=1}^{\infty} \exp[-n(E_k^l - E_1^l)], \quad (12)$$

and

$$\rho = \frac{1}{l^d} \sum_{n=1}^{\infty} (\zeta_l(\rho))^n \sum_{k=1}^{\infty} \exp[-n(E_k^l - E_1^l)], \quad (13)$$

Our main result is contained in the following

**Theorem 1.** For  $l \rightarrow \infty$ :

$$p_l(\rho) \sim \begin{cases} p(\rho) + \frac{S_1(2\pi)^{1/2}}{4l} \left[ \rho \frac{\sum_{n=1}^{\infty} (\zeta(\rho))^n \cdot n^{(1-d)/2}}{\sum_{n=1}^{\infty} (\zeta(\rho))^n \cdot n^{(2-d)/2}} - \sum_{n=1}^{\infty} (\zeta(\rho))^n \cdot (2\pi n)^{-(d+1)/2} \right], & \rho < \rho_c, \quad (14) \\ p(\rho) - \frac{\pi S_1}{2l} \sum_{n=1}^{\infty} \frac{1}{(2\pi n)^{(d+1)/2}}, & \rho \geq \rho_c, \quad (15) \end{cases}$$

and

**Theorem 2.** The occupation density of the ground state  $\rho_l(1)$  is asymptotically given by ( $l \rightarrow \infty$ )

$$\rho_l(1) \equiv \frac{1}{l^d} \cdot \frac{\zeta_l(\rho)}{1 - \zeta_l(\rho)} \sim \begin{cases} \frac{1}{l^d} \cdot \frac{\zeta(\rho)}{1 - \zeta(\rho)}, & \rho < \rho_c \quad (16) \\ \frac{S_1 \log l}{4\pi l}, & \rho = \rho_c, \quad d = 3 \quad (17) \\ \frac{S_1}{4l} \sum_{n=1}^{\infty} (2\pi n)^{(1-d)/2}, & \rho = \rho_c, \quad d \geq 4 \quad (18) \\ \rho - \rho_c, & \rho > \rho_c, \quad d \geq 3. \quad (19) \end{cases}$$

It is clear that Theorem 2 contains the asymptotic behaviour of  $\zeta_l(\rho)$ . The asymptotic behaviour of  $\zeta_l(\rho)$  for  $\rho > \rho_c$  was proved in [12] in the case where  $B_1$  is a star-shaped region with unit volume. Before we prove these theorems we would like to mention that one can extract from [9] a bound on  $p_B(\rho)$  which holds for all

possible convex regions  $B$ :

$$|p_B(\rho) - p(\rho)| \leq \begin{cases} c_3(\rho) \frac{S(B)}{V(B)} \log \frac{V(B)}{S(B)}, \\ c_4(\rho) \cdot \frac{S(B)}{V(B)}, \end{cases} \quad (20)$$

for positive functions  $c_3(\rho), c_4(\rho) \dots$  (which are bounded for finite  $\rho$ ).

## 2. The asymptotic behaviour of the ground state density and the pressure

In order to prove the Theorems 1 and 2 we need some sharp estimates on

$$Z(t) \equiv \text{trace} (e^{t \Delta/2}) = \sum_{k=1}^{\infty} e^{-t E_k^B}, \quad t > 0. \quad (22)$$

These will be given in the following lemmas.

**Lemma 1.** *For any region  $B$  with a regular boundary*

$$Z(t) \leq \frac{V(B)}{(2\pi t)^{d/2}}, \quad t > 0. \quad (23)$$

*For the proof we refer to [13].*

**Lemma 2.** *For convex regions  $B$*

$$\left| Z(t) - \frac{V(B)}{(2\pi t)^{d/2}} \right| \leq \frac{e^{d/2} \cdot S(B)}{2 \cdot (2\pi t)^{(d-1)/2}}, \quad t > 0. \quad (24)$$

*For the proof we refer to [14] or [15].*

**Lemma 3.** *For convex regions  $B$  with a boundary  $\partial B$  such that at each point of  $\partial B$  the curvature is bounded from above by  $1/R(B)$  ( $R(B) > 0$ ) one has*

$$\left| Z(t) - \frac{V(B)}{(2\pi t)^{d/2}} + \frac{S(B)}{4 \cdot (2\pi t)^{(d-1)/2}} \right| \leq \frac{t \cdot S(B)}{2 \cdot (2\pi t)^{d/2} \cdot R(B)} \left\{ (d-1) \log \left( 1 + \frac{2R^2(B)}{t} \right) + \pi^{1/2} \cdot d(d^{3/2} + \frac{1}{2}) \right\}. \quad (25)$$

Lemma 3 was proved in [15].

**Lemma 4.** *For  $B_1$  convex,  $R(B_1) > 0$  and  $B_l$  the dilation of  $B_1$  we have ( $l \rightarrow \infty$ )*

$$\frac{1}{l^d} \sum_{k=1}^{\infty} (e^{E_k^l} - 1)^{-1} \sim \begin{cases} \rho_c - \frac{S_1 \log l}{4\pi l} + O\left(\frac{1}{l}\right) \end{cases} \quad (26)$$

$$\begin{cases} \rho_c - \frac{S_1}{4l} \sum_{n=1}^{\infty} (2\pi n)^{(1-d)/2} + O\left(\frac{\log^2 l}{l^2}\right) \end{cases} \quad (27)$$

*Proof.* Let  $[l^2]$  be the greatest integer equal or less than  $l^2$ . By Lemma 3, (6),

(8), (9) and (10) we get

$$\begin{aligned}
 \frac{1}{l^d} \sum_{k=1}^{\infty} (e^{E_k^l} - 1)^{-1} &\geq \frac{1}{l^d} \sum_{n=1}^{[l^2]} \sum_{k=1}^{\infty} e^{-nE_k^l} \\
 &\geq \rho_c - \sum_{n=[l^2]+1}^{\infty} (2\pi n)^{-d/2} - \sum_{n=1}^{[l^2]} \frac{S_1}{4l \cdot (2\pi n)^{(d-1)/2}} \\
 &\quad - \sum_{n=1}^{[l^2]} \frac{S_1}{l^2 R_1} \cdot n^{1-d/2} \left( 1 + \log \left( 1 + \frac{2l^2 R_1^2}{n} \right) \right) \\
 &\geq \begin{cases} \rho_c - \frac{S_1 \log l}{4\pi l} - O\left(\frac{1}{l}\right), & d=3 \\ \rho_c - \frac{S_1}{4l} \sum_{n=1}^{\infty} (2\pi n)^{(1-d)/2} - O\left(\frac{\log^2 l}{l^2}\right), & d \geq 4. \end{cases}
 \end{aligned} \tag{28}$$

We obtain an upperbound by Lemmas 1, 3 and (6), (8), (9) and (10):

$$\begin{aligned}
 \frac{1}{l^d} \sum_{k=1}^{\infty} (e^{E_k^l} - 1)^{-1} &\leq \frac{1}{l^d} \sum_{n=1}^{[l^2]} \sum_{k=1}^{\infty} e^{-nE_k^l} + \sum_{n=[l^2]+1}^{\infty} \frac{1}{(2\pi n)^{d/2}} \\
 &\leq \rho_c - \frac{S_1}{4l} \sum_{n=1}^{[l^2]} (2\pi n)^{(1-d)/2} + \sum_{n=1}^{[l^2]} \frac{S_1}{l^2 R_1} n^{1-d/2} \left( 1 + \log \left( 1 + \frac{2l^2 R_1^2}{n} \right) \right) \\
 &\leq \begin{cases} \rho_c - \frac{S_1 \log l}{4\pi l} + O\left(\frac{1}{l}\right), & d=3 \\ \rho_c - \frac{S_1}{4l} \sum_{n=1}^{\infty} (2\pi n)^{(1-d)/2} + O\left(\frac{\log^2 l}{l^2}\right), & d=4. \end{cases}
 \end{aligned} \tag{29}$$

*Proof of Theorem 2.* For  $\rho \geq \rho_c$  we use inequality (12) of [9]:

$$e^{-E_1^l} \leq \zeta_l(\rho) \leq 1, \quad \rho \geq \rho_c. \tag{30}$$

Thus

$$\begin{aligned}
 &l^{-d} \sum_{k=2}^{\infty} \zeta_l(\rho) (e^{E_k^l - E_1^l} - \zeta_l(\rho))^{-1} - l^{-d} \sum_{k=1}^{\infty} (e^{E_k^l} - 1)^{-1} \\
 &\leq l^{-d} (e^{E_1^l} - 1)^{-1} + l^{-d} \sum_{k=2}^{\infty} \{ (e^{E_k^l - E_1^l} - 1)^{-1} - (e^{E_k^l} - 1)^{-1} \} \\
 &\leq l^{-d} (E_1^l)^{-1} + l^{-d} \sum_{k=2}^{\infty} e^{E_k^l} (e^{E_1^l} - 1) (e^{E_k^l} - e^{E_1^l})^{-1} \cdot (e^{E_k^l} - 1)^{-1} \\
 &\leq l^{-d+2} (E_1^l)^{-1} + l^{-d} \sup_{m \geq 2} \frac{e^{E_m^l} - 1}{e^{E_m^l} - e^{E_1^l}} \cdot \sum_{k=2}^{\infty} \frac{e^{E_k^l} (e^{E_1^l} - 1)}{(e^{E_k^l} - 1)^2} \\
 &\leq l^{-d+2} (E_1^l)^{-1} + l^{-d} \frac{(e^{E_1^l} - 1)(e^{E_2^l} - 1)}{(e^{E_2^l} - e^{E_1^l})} \cdot \sum_{n=1}^{\infty} n e^{-nE_1^l/2} \cdot \sum_{k=1}^{\infty} e^{-nE_k^l/2} \\
 &\leq l^{-d+2} (E_1^l)^{-1} + e^{E_2^l/l^2} \cdot \frac{E_1^1 \cdot E_2^1}{E_2^1 - E_1^1} \cdot \frac{1}{l^2 \pi^{d/2}} \sum_{n=1}^{\infty} n^{1-d/2} e^{-nE_1^l/(2l^2)} \\
 &\leq \begin{cases} (lE_1^l)^{-1} + e^{E_2^l/l^2} \cdot \frac{E_2^1 \cdot (E_1^l)^{1/2}}{E_2^1 - E_1^1} \cdot \frac{1}{l}, & d=3 \\ l^{-d+2} (E_1^l)^{-1} - e^{E_2^l/l^2} \frac{E_2^1 \cdot E_1^1}{E_2^1 - E_1^1} \cdot \frac{1}{l^2} \log(1 - e^{-E_1^l/(2l^2)}), & d \geq 4. \end{cases}
 \end{aligned} \tag{31}$$

We have used the inequality  $e^x - 1 \leq xe^x$  and Lemma 1. The combination of (31), Lemma 4 and (13) proves (17), (18) and (19) of Theorem 2. Line (16) follows simply from the convergence of  $\zeta_l(\rho) \rightarrow \zeta(\rho)$  (see [9] or [12] for the proof).

Without proof we state a result (sharper than (16)) for  $\rho < \rho_c$ :

$$\zeta_l(\rho) \sim \zeta(\rho) + \zeta(\rho) \frac{S_1 \cdot (2\pi)^{1/2}}{4l} \left( \sum_{n=1}^{\infty} (\zeta(\rho))^n \cdot n^{(1-d)/2} \right) \cdot \left( \sum_{n=1}^{\infty} (\zeta(\rho))^n \cdot n^{(2-d)/2} \right)^{-1}. \quad (32)$$

We see that there are two essential features in the proof of Theorem 2: the scaling of the eigenvalues (relation (10)) and the non-degeneracy of the ground state. From this it follows already that the occupation density of the second level  $\rho_l(2)$  becomes small:

$$\rho_l(2) \equiv \frac{1}{l^d} \cdot \frac{\zeta_l(\rho)}{e^{E_2^l - E_1^l} - \zeta_l(\rho)} \leq \frac{1}{l^d (E_2^l - E_1^l)} = O(l^{2-d}). \quad (33)$$

This is of course not true for general subsequences  $B_l$  (see [9]).

*Proof of Theorem 1.* We start with an estimate:

$$\begin{aligned} & \left| \rho_l(\rho) - \frac{1}{l^d} \sum_{n=1}^{\infty} \frac{(\zeta_l(\rho))^n}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} \right| \\ & \leq \frac{1}{l^d} \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{(\zeta_l(\rho))^n}{n} e^{-n(E_k^l - E_1^l)} (1 - e^{-nE_1^l}) \\ & \leq \frac{E_1^l}{l^d} \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} (\zeta_l(\rho))^n e^{-n(E_k^l - E_1^l)} \leq \frac{E_1^l \rho}{l^2}. \end{aligned} \quad (34)$$

Consider  $\rho \geq \rho_c$ : We use (30) and Lemma 1 to obtain:

$$\begin{aligned} & \frac{1}{l^d} \sum_{n=1}^{\infty} \frac{(\zeta_l(\rho))^n}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} \\ & \leq \frac{1}{l^d} \sum_{n=1}^{[l^2]} \frac{1}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} + \sum_{n=[l^2]+1}^{\infty} \frac{1}{n \cdot (2\pi n)^{d/2}}, \end{aligned} \quad (35)$$

and

$$\begin{aligned} & \frac{1}{l^d} \sum_{n=1}^{\infty} \frac{(\zeta_l(\rho))^n}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} \geq \frac{1}{l^d} \sum_{n=1}^{\infty} \frac{e^{-nE_1^l}}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} \\ & \geq \frac{1}{l^d} \sum_{n=1}^{[l^2]} \frac{1}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} - \frac{E_1^l}{l^2} \sum_{n=1}^{\infty} (2\pi n)^{-d/2}. \end{aligned} \quad (36)$$

Furthermore we obtain (as in Lemma 4):

$$\frac{1}{l^d} \sum_{n=1}^{[l^2]} \frac{1}{n} \sum_{k=1}^{\infty} e^{-nE_k^l} \sim \sum_{n=1}^{\infty} \frac{1}{n \cdot (2\pi n)^{d/2}} - \frac{\pi}{2l} \sum_{n=1}^{\infty} (2\pi n)^{-(d+1)/2} \cdot S_1. \quad (37)$$

The combination of (34)–(37) proves Theorem 1 for  $\rho \geq \rho_c$ . One proves Theorem 1 for  $\rho < \rho_c$  using (32), (34) and the Lemmas 2 and 3.

Estimate (34) illustrates that Theorem 1 states an asymptotic expansion of  $p_l(\rho)$  for large  $l$  at fixed mean density  $\rho$ . Hence, for  $l$  fixed one can always find a large mean density  $\rho$ , for which the expansion is a bad approximation to  $p_l(\rho)$ .

## Acknowledgement

I would like to thank Professor J. T. Lewis for helpful discussions and suggestions.

## REFERENCES

- [1] M. J. BUCKINGHAM and W. M. FAIRBANK, *The nature of the  $\lambda$ -transition in liquid helium*. Progress in Low Temperature Physics. Ed. Gorter, C. J., North-Holland 3, 80–112 (1961).
- [2] M. VAN DEN BERG and J. T. LEWIS, *On the free boson gas in a weak external potential*. Commun. Math. Phys. 81, 475–494 (1981).
- [3] M. VAN DEN BERG and J. T. LEWIS, *On the  $\lambda$ -singularity in the specific heat of a free boson gas in a gravitational field*. Internal Report 177, (1981). Institute for Theoretical Physics, University of Groningen, The Netherlands.
- [4] M. VAN DEN BERG, J. T. LEWIS and PH. DE SMEDT, *The imperfect boson gas* (in preparation).
- [5] M. N. BARBER, *Critical phenomena of finite thickness II*. Aust. J. Phys. 26, 483–500 (1973).
- [6] M. N. BARBER and M. E. FISHER, *Critical phenomena in systems of finite thickness III*. Phys. Rev. A8, 1124–1135 (1973).
- [7] S. GREENSPOON and R. K. PATHRIA, *Bose–Einstein condensation in finite noninteracting systems*. Phys. Rev. A9, 2103–2110 (1974).
- [8] C. S. ZASADA and R. K. PATHRIA, *Low temperature behaviour of bose systems confined to restricted geometries*. Phys. Rev. A14, 1269–1280 (1976).
- [9] M. VAN DEN BERG, *On condensation in the free boson gas and the spectrum of the laplacian*. Journ. Stat. Phys. 31, 623–637 (1983).
- [10] M. VAN DEN BERG and J. T. LEWIS, *On generalized condensation in the free boson gas*. Physica 110A, 550–564 (1982).
- [11] M. VAN DEN BERG, *On boson condensation into an infinite number of low-lying levels*. Journ. Math. Phys. 23, 1159–1161 (1982).
- [12] J. T. LEWIS and J. V. PULÈ, *The equilibrium states of the free boson gas*. Commun. Math. Phys. 36, 1–18 (1974).
- [13] D. B. RAY, *Spectra of second-order differential operators*. Trans. Am. Math. Soc. 77, 299–321 (1954).
- [14] N. ANGELESCU and G. NENCIU, *On the independence of the thermodynamic limit on the boundary conditions in quantum statistical mechanics*. Commun. Math. Phys. 29, 15–30 (1973).
- [15] M. VAN DEN BERG, *A uniform bound on trace  $(e^{t\Delta})$  for convex regions in  $R^n$  with smooth boundaries*. Accepted for publication in Commun. Math. Phys.