

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 58 (1985)  
**Heft:** 6

**Artikel:** Spectral concentration for Hamiltonian systems  
**Autor:** Hinton, D.B. / Shaw, J.K.  
**DOI:** <https://doi.org/10.5169/seals-115630>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 17.09.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Spectral concentration for Hamiltonian systems

By D. B. Hinton

Department of Mathematics, University of Tennessee, Knoxville, TN 37996-1300

J. K. Shaw

Department of Mathematics, Virginia Tech, Blacksburg, VA 24061-4097

(7. III. 1985; rev. 12. VI. 1985)

*Abstract.* We consider a sequence of operators  $T_n$  arising from Hamiltonian systems and require that  $T_n \rightarrow T$ , where  $T$  corresponds to a fixed Hamiltonian system. Letting  $\tau_n(\lambda)$  and  $\tau(\lambda)$  denote the respective spectral functions, we give sufficient conditions that  $\tau_n(\lambda_2) - \tau_n(\lambda_1) \rightarrow \tau(\lambda_2) - \tau(\lambda_1)$  at continuity points  $\lambda_1$  and  $\lambda_2$  of  $\tau(\lambda)$ .

## 1. Introduction

‘Spectral concentration’ is a term arising from perturbation theory when the spectrum  $\sigma(T_0)$  of an operator  $T_0$  acting in a Hilbert space  $H$  is compared to the spectrum  $\sigma(T_\varepsilon)$  of a perturbed operator  $T_\varepsilon$ , where  $\varepsilon > 0$  is a small parameter and  $T_\varepsilon \rightarrow T_0$  (in a suitable sense) as  $\varepsilon \rightarrow 0$ . In the classical cases of physical interest  $\sigma(T_\varepsilon)$  is qualitatively different from, but quantitatively near to,  $\sigma(T_0)$ .

The expression itself originated with work of Titchmarsh ([22], [23], [24]); also see [20, p. 60]) on the mathematical formulation of the Stark effect in hydrogen. Here one considers the Sturm–Liouville eigenvalue problem

$$-y'' + \left( \frac{l(l+1)}{x^2} - \frac{c}{x} - \varepsilon x \right) y = \lambda y, \quad 0 < x < \infty, \quad (1.1)$$

where  $l$  is a positive integer,  $c$  is a constant,  $\varepsilon > 0$  and  $\lambda$  is the spectral parameter. In [23] the more general equation with  $\varepsilon x$  replaced by  $\varepsilon \sigma(x)$  is actually treated, where  $\sigma(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , and in its forerunner [22] the simpler equation  $-y'' + \varepsilon xy = \lambda y$ ,  $0 \leq x < \infty$ , is studied. For  $\varepsilon = 0$  (1.1) is the Hamiltonian in one dimension for a hydrogen atom ([7], [8]) and the perturbed case  $\varepsilon > 0$  is the corresponding Hamiltonian in a constant, weak electric field of strength  $\varepsilon$ . The unperturbed equation has continuous spectrum in  $[0, \infty)$  and a sequence of isolated negative eigenvalues clustering at  $\lambda = 0$  ([7], [15]). But for any positive  $\varepsilon$  the spectrum of (1.1) continuously fills the entire  $\lambda$ -axis, with no imbedded eigenvalues ([23], [15]). This apparent discrepancy throughout the negative energy spectrum is most easily explained in terms of the spectral function  $\rho_\varepsilon(\lambda)$  for (1.1) (we give precise definitions of all relevant terms from spectral theory in §2 below). For  $\lambda < 0$  the unperturbed spectral function  $\rho_0(\lambda)$  ( $\varepsilon = 0$ ) is a nondecreasing step function with jumps at the eigenvalues. But while  $\rho_\varepsilon(\lambda)$  is continuously differentiable ([15]) on  $-\infty < \lambda < \infty$ , the ‘spectral density’ function  $\rho'_\varepsilon(\lambda)$  has sharp peaks near the unperturbed eigenvalues and is small elsewhere for  $\lambda < 0$ .

Thus the main strength of the negative spectrum is concentrated near the unperturbed eigenvalues; see [22], [23], [24], [20, §12.5 and Notes] and also [19] for another example and further discussion.

Titchmarsh develops in [22] and [23] a theory which attributes the sharp peaks in  $\rho'_\varepsilon(\lambda)$  to poles of the perturbed  $m_\varepsilon(\lambda)$  function (defined in §3 below) lying near the eigenvalues but just below the negative real axis in the complex  $\lambda$ -plane. These ‘perturbed poles’, also called ‘resonances’ ([7]) and ‘pseudo-eigenvalues’ ([19]), transmit their strength to the spectral function through the Titchmarsh–Kodaira formula ([3])

$$\rho_\varepsilon(\lambda_2) - \rho_\varepsilon(\lambda_1) = \lim_{\delta \rightarrow 0^+} \pi^{-1} \int_{\lambda_1}^{\lambda_2} \operatorname{Im} m_\varepsilon(\mu + i\delta) d\mu, \quad (1.2)$$

which is valid at points of continuity  $\lambda_1$  and  $\lambda_2$  of  $\rho_\varepsilon(\lambda)$ . The resonances have exponentially small (as  $\varepsilon \rightarrow 0$ ) imaginary parts in the case of (1.1) ([7], [8]) and are given approximately by Titchmarsh’s classical real eigenvalue perturbation formula in [22]. Less well-behaved perturbing potentials may produce resonances which do not satisfy such relations ([7], [6]).

Another feature of the concentration phenomenon, also introduced by Titchmarsh ([24]), seeks to measure the extent of concentration of the spectrum on small intervals about unperturbed eigenvalues. Suppose that  $T_0$  and  $T_\varepsilon$  are the operators, having spectra  $\sigma(T_0)$  and  $\sigma(T_\varepsilon)$ , respectively, where  $\sigma(T_0)$  is discrete and  $\sigma(T_\varepsilon)$  is continuous. Let  $f \in H = L^2[0, \infty)$  and let  $g_\varepsilon(\lambda)$  be the transform of  $f$ ; see [3, chapter 7], [24] and Remark 5 in §6 below. Then one has the Parseval formula

$$\int_0^\infty |f(x)|^2 dx = \int_{-\infty}^\infty |g_\varepsilon(\lambda)|^2 d\rho_\varepsilon(\lambda),$$

and the theory asserts that there are open intervals  $\Delta_n = \Delta_n(\varepsilon)$ , each  $\Delta_n$  containing the eigenvalue  $\lambda_n$  and whose diameters  $|\Delta_n(\varepsilon)| \rightarrow 0$ , as  $\varepsilon \rightarrow 0$ , such that if  $\Delta = (\bigcup_{n=1}^\infty \Delta_n)$  then

$$\int_0^\infty |f(x)|^2 dx = \int_\Delta |g_\varepsilon(\lambda)|^2 d\rho_\varepsilon(\lambda) + o(1), \quad \varepsilon \rightarrow 0. \quad (1.3)$$

It is then a question of how large the intervals  $\Delta_n$  must be in order that (1.3) should hold. One says, for example, that the spectrum is concentrated to order  $p$  if the widths can be taken to be  $2\varepsilon^p$ ; see [4], [19], [24] and the Notes, pp. 64–68, of [20] where there are numerous references.

The present paper takes a somewhat different approach to the subject. Although a great deal is known about convergence of the spectral family  $E_\varepsilon(\lambda)$  of  $T_\varepsilon$  to the unperturbed family  $E_0(\lambda)$  of  $T_0$ , when  $T_\varepsilon \rightarrow T_0$  in some appropriate way (see [25, p. 286]), there does not appear to be a proof that the spectral jumps  $[\rho_\varepsilon(\lambda)]_{\lambda_1}^{\lambda_2} = \rho_\varepsilon(\lambda_2) - \rho_\varepsilon(\lambda_1)$  converge to the corresponding unperturbed expression  $[\rho_0(\lambda)]_{\lambda_1}^{\lambda_2}$ , as  $\varepsilon \rightarrow 0$ , even for the scalar equation (1.1). The proof of such a convergence theorem, in a systems context which includes (1.1), is the main purpose of this paper.

The convergence of  $[\rho_\varepsilon(\lambda)]_{\lambda_1}^{\lambda_2}$  to  $[\rho_0(\lambda)]_{\lambda_1}^{\lambda_2}$  is a qualitative measure of concentration. It implies, for example, that in the typical case where  $\rho_\varepsilon(\lambda)$  is differentiable

in a neighborhood of an isolated unperturbed eigenvalue  $\lambda_0$ ,  $\rho'_\varepsilon(\lambda)$  must necessarily be small except for a sharp peak very near  $\lambda_0$ .

It will be convenient to take as our basic equation the Hamiltonian system of [5] (and also [1], [10], [11]), which includes scalar equations as one case, on an interval  $[a, b]$ ,  $-\infty < a < b \leq \infty$ . Our principal tool is a generalization (§4) of a deep result of Atkinson in [2] which links  $\rho(\lambda)$  to  $m(\lambda)$  in an alternative way to (1.2). Our main convergence result (Theorem 1) is stated in §2, following the introduction of some notation and terminology, and is proved in §5.

A concluding section (§6) contains examples, a discussion of two singular endpoint problems and a connection between the convergence theorem and a type of convergence of spectral families.

## 2. Hamiltonian systems

This section contains background material on Hamiltonian systems and their spectral theory, included here for completeness.

We consider the  $2k \times 1$  first order system ([5], [1])

$$Jy' = [\lambda A(x) + B(x)]y, \quad a \leq x < b \leq \infty, \quad \lambda \text{ complex}, \quad (2.1)$$

where  $J$ ,  $A$  and  $B$  are  $2k \times 2k$  complex matrices,  $A^* = A$  and  $B^* = B$  (conjugate transpose),  $A \geq 0$  (nonnegative definite) and  $J = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ ,  $I$  being the  $k \times k$  identity. We regard (2.1) as regular at  $x = a$ , singular at  $x = b$  and take the coefficients to be locally integrable so that the usual existence and uniqueness properties hold. Let  $A$  be decomposable as

$$A(x) = \begin{pmatrix} A_1(x) & 0 \\ 0 & 0 \end{pmatrix}, \quad (2.2)$$

where  $A_1(x)$  is  $r \times r$  and invertible,  $1 \leq r \leq 2k$ ,  $A = A_1$  in case  $r = 2k$ , and finally let us assume the definiteness hypothesis ([1, p. 253])

$$\int_{\alpha}^{\beta} y^*(x) A(x) y(x) dx > 0, \quad a < \alpha < \beta < b, \quad (2.3)$$

for all solutions  $y$  of (2.1) which do not vanish identically.

Scalar equations  $-(pu')' + qu = \lambda wu$  are special cases of (2.1), as can be seen from the identity

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u \\ pu' \end{pmatrix}' = \left[ \lambda \begin{pmatrix} w & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} -q & 0 \\ 0 & p^{-1} \end{pmatrix} \right] \begin{pmatrix} u \\ pu' \end{pmatrix}. \quad (2.4)$$

The Dirac systems studied in [18] comprise another special case.

Associated with (2.1) is the linear space ([1])  $L_A^2[a, b)$  of equivalence classes of measurable functions  $f(x)$  defined on  $[a, b)$  such that  $\int_a^b f^*(x) A(x) f(x) dx < \infty$ , and it is natural to introduce the inner product  $(f, g)_A = \int_a^b f^* A g$  for  $f, g \in L_A^2[a, b)$ . There is induced the seminorm  $\|f\|_A = (f, f)_A^{1/2}$ , which is a norm in the event that  $A$  is invertible. Let  $P_r$  be the  $2k \times 2k$  matrix  $\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}$ , where  $I_r$  is the identity matrix

which has the size of  $A_1(x)$  in (2.2), and for  $g \in L_A^2[a, b]$  define  $g_1 = P_r g$ . Letting  $P_r L_A^2[a, b] = \{P_r g \mid g \in L_A^2[a, b]\}$ , we have  $\|f\|_A^2 = \int_a^b f_1^* A_1 f_1$  for all  $f \in P_r L_A^2[a, b]$ . Since  $A_1$  is invertible then  $\|\cdot\|_A$  restricted to  $P_r L_A^2[a, b]$  is a norm; hence  $P_r L_A^2[a, b]$  is a Hilbert space under the inner product  $(\cdot, \cdot)_A$ .

The limit point-limit circle theory of (2.1) is well-known; see [21], [10] and [13]. We know that the number of linearly independent solutions of (2.1) lying in  $L_A^2[a, b]$  is independent of  $\lambda$  in each halfplane  $\text{Im}(\lambda) > 0$  and  $\text{Im}(\lambda) < 0$ , and can range from  $k$  to  $2k$ . If there are exactly  $k$  linearly independent solutions for  $\text{Im}(\lambda) \neq 0$  we say that (2.1) is of limit-point type at  $x = b$ . We take as a basic hypothesis in this paper that

$$\text{system (2.1) is of limit point type at } x = b. \quad (2.5)$$

This hypothesis implies, for example, that certain boundary value problems associated with (2.1) have unique solutions. To discuss one particular type of boundary problem it will be convenient to introduce the  $2k \times 2k$  matrices  $E_\alpha$  given by

$$E_\alpha = \begin{pmatrix} \alpha_1^* & -\alpha_2^* \\ \alpha_2^* & \alpha_1^* \end{pmatrix}$$

where the  $\alpha_j$  are  $k \times k$  and satisfy  $\text{rank}[\alpha_1, \alpha_2] = k$ ,  $\alpha_1 \alpha_2^* = \alpha_2 \alpha_1^*$  and  $\alpha_1 \alpha_1^* + \alpha_2 \alpha_2^* = I$ . A typical boundary condition placed on a vector  $y$  at  $x = a$  is

$$[\alpha_1, \alpha_2]y(a) = v,$$

where  $v$  is a  $k \times 1$  constant vector. In [10] and [12] the present authors proved under assumption (2.5) that the problem  $Jy' = (\lambda A + B)y + Af$ , where  $f \in L_A^2[a, b]$ ,  $\text{Im}(\lambda) \neq 0$  and  $[\alpha_1, \alpha_2]y(a) = v$ , has a unique solution  $y \in L_A^2[a, b]$ . This result has operator-theoretic consequences which we now proceed to describe (see [11], [12]). Define an operator  $T$  with domain  $D(T) \subset L_A^2[a, b]$  by saying  $y \in D(T)$  provided (i)  $y \in L_A^2[a, b]$  (ii),  $y$  is locally absolutely continuous, (iii)  $[\alpha_1, \alpha_2]y(a) = 0$ , (iv)  $\begin{pmatrix} A_1^{-1} & 0 \\ 0 & 0 \end{pmatrix}(Jy' - By) \in P_r L_A^2[a, b]$ , (v)  $\begin{pmatrix} 0 & 0 \\ 0 & I_{2k-r} \end{pmatrix}(Jy' - By) = 0$ , and then defining  $T: D(T) \rightarrow P_r L_A^2[a, b]$  by the equation

$$(Ty)(x) = \begin{pmatrix} A_1^{-1}(x) & 0 \\ 0 & 0 \end{pmatrix}(Jy'(x) - B(x)y(x)). \quad (2.6)$$

Then the existence and uniqueness result cited above may be rephrased by saying that  $(T - \lambda P_r): D(T) \rightarrow P_r L_A^2[a, b]$  is a one-to-one and onto map whenever  $\text{Im}(\lambda) \neq 0$ . It also happens ([10], [11], [12]) that  $T - \lambda P_r$  has a bounded inverse for  $\text{Im}(\lambda) \neq 0$ , and in fact

$$\|(T - \lambda P_r)^{-1}f\|_A \leq |\text{Im}(\lambda)|^{-1} \|f\|_A. \quad (2.7)$$

The *resolvent set* (see [11], [12])  $\rho(T)$  of  $T$  is defined as the set of all complex  $\lambda$  such that  $(T - \lambda P_r)^{-1}: P_r L_A^2[a, b] \rightarrow L_A^2[a, b]$  exists and is bounded; thus  $\rho(T)$  contains all nonreal numbers by (2.7). The *spectrum*  $\sigma(T)$  of  $T$  is the complement of  $\rho(T)$  in the set of complex numbers. The set of isolated points of  $\sigma(T)$  is called the *point spectrum*, and is denoted by  $P(T)$ . The set  $\sigma(T) - P(T)$  is called the *essential spectrum* and the subset  $PC(T) \subset E(T)$  consisting of imbedded eigenvalues is termed the *point-continuous spectrum*; see [11]. We call  $C(T) =$

$E(T) - PC(T)$  the *continuous spectrum* of  $T$ . We cannot appeal to the standard definitions of the above terms because  $T$  is not in the strict sense a Hilbert space operator.

The spectral function  $\tau(\lambda)$  of the operator  $T$  is a  $2k \times 2k$  matrix function defined on  $-\infty < \lambda < \infty$  which is right-continuous, nondecreasing (in the nonnegative definite sense) and normalized by  $\tau(0) = 0$ ; see [11] and [12]. It is constant on that part of the resolvent intersecting the real axis, so that its points of increase comprise  $\sigma(T)$ , there are jump discontinuities at both isolated and imbedded eigenvalues, and  $\tau(\lambda)$  is continuous on  $C(T)$ . There is an analogue of the Titchmarsh–Kodaira formula (1.2); we give this in §3. In the scalar case where (2.1) reduces to (2.4)  $\tau_{22}(\lambda)$  is the usual scalar spectral function.

The convergence theorem we prove will be for spectral functions  $\tau_n(\lambda)$ ,  $n = 1, 2, 3, \dots$ , corresponding to operators  $T_n$  given by (2.6). By means of the imbedding (2.4) our result will include spectral functions for scalar equations. Thus let there be given sequences  $\{A_n(x)\}_{n=1}^\infty$ ,  $\{B_n(x)\}_{n=1}^\infty$  and  $\{E_{n,\alpha}\}_{n=1}^\infty$  such that the systems

$$Jy' = [\lambda A_n(x) + B_n(x)]y, \quad a \leq x < b \leq \infty, \quad (2.1)_n$$

satisfy the same hypotheses (2.2)–(2.6) as system (2.1). Let  $T_n$  and  $\tau_n$  denote the corresponding operators and spectral functions. As regards the convergence of  $T_n$  to  $T$ , we assume specifically that  $A_n \rightarrow A$  and  $B_n \rightarrow B$  in the  $L^1$ -norm on compact subsets of  $[a, b)$ ; i.e.,

$$\lim_{n \rightarrow \infty} \int_c^d \|A_n(x) - A(x)\| dx = \lim_{n \rightarrow \infty} \int_c^d \|B_n(x) - B(x)\| dx = 0, \quad (2.8)$$

where  $a \leq c < d < b$  and where  $\|\cdot\|$  denotes the matrix operator norm  $\|A(x)\| = \sup \{\|A(x)v\| \mid \|v\| = 1\}$ . Let  $E_{n,\alpha} \rightarrow E_\alpha$ . Then our principal result may be stated as follows.

**Theorem 1.** Let  $\lambda_1$  and  $\lambda_2$  be points of continuity of the spectral function  $\tau(\lambda)$ . Then

$$\lim_{n \rightarrow \infty} (\tau_n(\lambda_2) - \tau_n(\lambda_1)) = \tau(\lambda_2) - \tau(\lambda_1). \quad (2.9)$$

This is proved in §5. The method of proof will cover the case of a continuous perturbation parameter  $\varepsilon$ ; i.e.,  $A_\varepsilon \rightarrow A$  and  $B_\varepsilon \rightarrow B$ , as  $\varepsilon \rightarrow 0$ , in the sense of (2.8). Hence we include operators of the type (1.1) restricted to  $0 < a \leq x < \infty$ . Two singular endpoint problems will be discussed in §6.

### 3. Preliminaries

Titchmarsh–Weyl functions for Hamiltonian systems may be defined in terms of the initial value matrices  $E_\alpha$  of §2. Following [10] and [12], let  $Y(x, \lambda)$  be the unique fundamental matrix solution of (2.1) which satisfies for all  $\lambda$  the initial condition  $Y(a, \lambda) = E_\alpha$ , and let us partition  $Y(x, \lambda)$  into  $k \times k$  blocks by writing

$$Y(x, \lambda) = \begin{pmatrix} \Theta(x, \lambda) & \Phi(x, \lambda) \\ \hat{\Theta}(x, \lambda) & \hat{\Phi}(x, \lambda) \end{pmatrix}. \quad (3.1)$$



Then under hypothesis (2.5) the limit

$$M(\lambda) = -\lim_{x \rightarrow b} \Phi^{-1}(x, \lambda) \Theta(x, \lambda), \quad \operatorname{Im}(\lambda) \neq 0, \quad (3.2)$$

exists and defines a  $k \times k$  matrix analytic function of  $\lambda$  in the two half  $\lambda$ -planes. We suppress in our notation the dependence of  $M(\lambda)$  and  $Y(x, \lambda)$  on the initial value matrix  $E_\alpha$ . For the scalar case arising via (2.4),  $M(\lambda)$  reduces to the usual scalar  $m(\lambda)$  function as given, for example, in [3].

Thus there is associated with the operator (2.6) the uniquely defined Titchmarsh–Weyl coefficient (3.2). The  $M(\lambda)$  function carries complete information on the spectrum  $\sigma(T)$  of  $T$ . To quote only part of the main result of [11] (see also [12]) we can say that  $\lambda \in \rho(T)$  if and only if  $M$  is analytic at  $\lambda$ ,  $\lambda \in P(T)$  if and only if  $M$  has a simple pole at  $\lambda$  and  $\lambda \in C(T)$  if and only if  $M$  is not analytic at  $\lambda$  and we have  $\lim_{\nu \rightarrow 0} \nu M(\lambda + i\nu) = 0$ ; for the characterization of  $PC(T)$ , see [12].

There is a Titchmarsh–Kodaira formula analogous to (1.2), but for this we need to mention the ‘characteristic function’  $F(\lambda)$ , introduced by F. V. Atkinson in [1], for the operator  $T$ . This is a  $2k \times 2k$  matrix function which may be shown ([10], [11]) to equal

$$F(\lambda) = \begin{pmatrix} 0 & (\frac{1}{2})I \\ (\frac{1}{2})I & M(\lambda) \end{pmatrix}. \quad (3.3)$$

In (3.3) we have reversed the sign of  $F(\lambda)$  from the way it appears in [10] and [11]. In [11] we prove the following relations between  $F(\lambda)$  and  $\tau(\lambda)$ :

$$\lim_{\delta \rightarrow 0+} \pi^{-1} \int_{\lambda_1}^{\lambda_2} \operatorname{Im} F(\mu + i\delta) d\mu = \tau(\lambda_2) - \tau(\lambda_1), \quad (3.4)$$

( $\operatorname{Im} F = (F - F^*)/2i$ ) at points  $\lambda_1$  and  $\lambda_2$  of continuity of  $\tau$ ;

$$F(\lambda) = \int_{-\infty}^{\infty} \left( \frac{1}{\mu - \lambda} - \frac{\mu}{1 + \mu^2} \right) d\tau(\mu) + K_1 \lambda + K_2, \quad \operatorname{Im}(\lambda) \neq 0, \quad (3.5)$$

for constant Hermitian matrices  $K_1 \geq 0$  and  $K_2$ . Recall in (3.5) that  $d\tau(\mu) \geq 0$ .

Let  $Y_n(x, \lambda)$  be the fundamental matrix solution of  $(2.1)_n$  which satisfies initial values  $Y_n(a, \lambda) = E_{n,\alpha}$ , for all  $\lambda$ , where

$$E_{n,\alpha} = \begin{pmatrix} \alpha_{1,n}^* & -\alpha_{2,n}^* \\ \alpha_{2,n}^* & \alpha_{1,n}^* \end{pmatrix} \rightarrow E_\alpha, \quad n \rightarrow \infty,$$

and where the  $\alpha_{i,n}$  satisfy the same hypotheses as the  $\alpha_i$  in  $E_\alpha$ . Let us denote by  $M_n(\lambda)$  and  $F_n(\lambda)$  the corresponding Titchmarsh–Weyl coefficient and characteristic function for the operator  $T_n$  arising via (2.6) from equation  $(2.1)_n$  and initial values  $E_{n,\alpha}$ . We are going to establish that  $M_n(\lambda) \rightarrow M(\lambda)$ ,  $n \rightarrow \infty$ , uniformly on compact subsets of  $\operatorname{Im}(\lambda) \neq 0$  (so that the same is true for  $F_n \rightarrow F$  by (3.3)) and the first step in doing so is the following.

**Lemma 3.1.** *Let  $[c, d]$  be a compact subinterval of  $[a, b)$  and let  $K$  be a compact subset of the  $\lambda$ -plane. Then  $Y_n(x, \lambda) \rightarrow Y(x, \lambda)$ ,  $n \rightarrow \infty$ , uniformly on  $[c, d] \times K$ .*

*Proof.* We have for  $a \leq x \leq d$  and  $\lambda \in K$  that

$$\begin{aligned} Y(x, \lambda) - Y_n(x, \lambda) &= (E_\alpha - E_{n,\alpha}) + J^{-1} \int_a^x [(\lambda A + B)Y - (\lambda A_n + B_n)Y_n] dt \\ &= (E_\alpha - E_{n,\alpha}) + J^{-1} \int_a^x (\lambda A_n + B_n)(Y - Y_n) dt + J^{-1} \\ &\quad \times \int_a^x [\lambda(A - A_n) + (B - B_n)]Y dt. \end{aligned}$$

It follows that

$$\begin{aligned} \|Y(x, \lambda) - Y_n(x, \lambda)\| &\leq \|E_\alpha - E_{n,\alpha}\| + \int_a^d (|\lambda| \|A - A_n\| + \|B - B_n\|) \|Y\| dt \\ &\quad + \int_a^d (|\lambda| \|A_n\| + \|B_n\|) \|Y - Y_n\| dt \\ &= \varepsilon_n + \int_a^d r_n(t) \|Y - Y_n\| dt \end{aligned}$$

where  $\varepsilon_n$  stands for the first two terms on the right of the inequality and  $r_n(t)$  is the first part of the integrand in the third. From the Gronwall inequality ([9, p. 24])

$$\|Y - Y_n\| \leq \varepsilon_n \exp \int_a^d r_n(t) dt,$$

and this expression clearly tends to 0 uniformly over  $K$  because of (2.8) and the fact that  $\|Y(x, \lambda)\|$  is uniformly bounded on  $[c, d] \times K$ . This completes the proof.

**Lemma 3.2.** *Let  $K^+$  be a compact subset of  $\text{Im } \lambda > 0$ . Then  $M_n(\lambda) \rightarrow M(\lambda)$  uniformly on  $K^+$  (and therefore  $F_n(\lambda) \rightarrow F(\lambda)$  uniformly on  $K^+$ ).*

*Proof.* We will represent  $M_n(\lambda)$  and  $M(\lambda)$  by the matrix circle method of [13]. For  $a < d < b$  and  $\text{Im } \lambda > 0$  let

$$\begin{pmatrix} \mathcal{A} & \mathcal{B}^* \\ \mathcal{B} & \mathcal{D} \end{pmatrix}(d, \lambda) = -iY^*(d, \lambda)JY(d, \lambda) \quad (3.6)$$

and then let

$$\begin{aligned} C_d(\lambda) &= -\mathcal{D}^{-1}(d, \lambda)\mathcal{B}(d, \lambda), \\ R_d^1(\lambda) &= \mathcal{D}^{-1/2}(d, \lambda), \\ R_d^2(\lambda) &= [\mathcal{B}^*\mathcal{D}^{-1}\mathcal{B} - \mathcal{A}]^{1/2}(d, \lambda); \end{aligned} \quad (3.7)$$

see [13] for proofs that these expressions exist. Let  $\mathcal{A}_n, \mathcal{B}_n$ , etc, be the corresponding expressions generated by the fundamental matrix  $Y_n(d, \lambda)$ . Then there are matrices  $V_d(\lambda)$  and  $V_{n,d}(\lambda)$  such that  $V_d^*V_d \leq I$ ,  $V_{n,d}^*V_{n,d} \leq I$  (see [13]) and

$$\begin{aligned} M &= C_d + R_d^1 V_d R_d^2, \\ M_n &= C_{n,d} + R_{n,d}^1 V_{n,d} R_{n,d}^2. \end{aligned}$$



Therefore

$$\begin{aligned} \|M(\lambda) - M_n(\lambda)\| &\leq \|C_d(\lambda) - C_{n,d}(\lambda)\| + \|R_d^1(\lambda)\| \|R_d^2(\lambda)\| \\ &\quad + \|R_{n,d}^1(\lambda)\| \|R_{n,d}^2(\lambda)\|. \end{aligned} \quad (3.8)$$

We know from [13] that  $R_d^1(\lambda)$  and  $R_d^2(\lambda)$  are continuous in  $d$  and  $\lambda$  and decreases to 0 as  $d$  increases to  $b$ ; hence  $R_d^1(\lambda) \rightarrow 0$  and  $R_d^2(\lambda) \rightarrow 0$  uniformly on  $K^+$  as  $d \rightarrow b$ . Thus given  $\varepsilon > 0$  we may choose  $d$  sufficiently near  $b$  that  $\|R_d^i(\lambda)\| < \varepsilon$ . Having fixed  $d$  we then note the formulas (3.6) and (3.7) and invoke Lemma 3.1 to conclude  $\|C_d(\lambda) - C_{n,d}(\lambda)\| < \varepsilon$  and  $\|R_d^i(\lambda) - R_{n,d}^i(\lambda)\| < \varepsilon$  for  $n$  sufficiently large. Then from (3.8) we get  $\|M(\lambda) - M_n(\lambda)\| < \varepsilon + \varepsilon^2 + 4\varepsilon^2$ , when  $n$  is large and  $\lambda \in K^+$ , and from there the conclusion of the lemma.

#### 4. Atkinson inequality

Here we give a matrix version of Atkinson's extension in [2] of the Titchmarsh-Kodaira formula (1.2). In our case we shall be extending (3.4).

Let  $\delta > 0$ ,  $\lambda_1 < \lambda_2$  and define

$$\begin{aligned} G(\lambda_1, \lambda_2) &= \operatorname{Im} \int_{\lambda_1 + i\delta}^{\lambda_2 + i\delta} F(\lambda) d\lambda - [\pi\tau(\mu)]_{\lambda_1}^{\lambda_2} \\ &\quad - [\delta \operatorname{Re} F(\mu + i\delta)]_{\lambda_1}^{\lambda_2}, \end{aligned} \quad (4.1)$$

where  $\operatorname{Re} F = (F + F^*)/2$ , and define

$$H(\lambda_1, \lambda_2) = (\pi\delta/2)[\operatorname{Im} F(\lambda_1 + i\delta) + \operatorname{Im} F(\lambda_2 + i\delta)]. \quad (4.2)$$

We choose for path of integration in (4.1) the horizontal line joining  $\lambda_1 + i\delta$  to  $\lambda_2 + i\delta$ , so that  $d\lambda$  is real and the symbol  $\operatorname{Im}$  can be moved under the integral sign and attached to  $F(\lambda)$ .

**Lemma 4.1.** *The functions  $G$  and  $H$  are Hermitian,  $H > 0$  and*

$$-H \leq G \leq H \text{ (nonnegative definite sense)}. \quad (4.3)$$

*Proof.* The Hermitian properties follow from that of  $\tau$  ([11]) and the definitions of  $\operatorname{Re} F$  and  $\operatorname{Im} F$ . Positive definiteness of  $H$  is a consequence of the Pick-Neumanlinna property  $\operatorname{Im} F(\lambda) > 0$  for  $\operatorname{Im}(\lambda) > 0$ ; see [10] and [11].

Emulating Atkinson's proof, we take (3.5) as the starting point for the proof of (4.3). For the purpose of the proof, in fact, it is best to regard  $\tau(\mu)$  as given and (3.5) as the definition of  $F(\lambda)$ , for this opens the way for some preliminary reductions. Note first that the constant  $K_2$  does not appear in either (4.1) or (4.2) due to cancellation. Hence there is no loss in generality in setting  $K_2 = 0$ . Furthermore,  $K_1$  does not appear in (4.1), because its effect is cancelled between the first and third terms of (4.1), but it contributes a nonnegative definite term to (4.2). Therefore (4.3) is true for general  $K_1 \geq 0$  if it is true for  $K_1 = 0$ . This being the case, we take  $K_1 = K_2 = 0$  and prove (4.3) for the special form

$$F(\lambda) = \int_{-\infty}^{\infty} \left( \frac{1}{\mu - \lambda} - \frac{\mu}{1 + \mu^2} \right) d\tau(\mu). \quad (4.4)$$

A final preliminary reduction is accomplished by replacing  $\tau(\mu)$  by a simplified

function  $\tau(\mu; R)$  which is nondecreasing and agrees with  $\tau(\mu)$  for  $-R \leq \mu \leq R$  but is constant outside  $[-R, R]$ . Let  $F(\lambda; R)$  denote the corresponding characteristic function given by (4.4). Then we have  $\tau(\mu; R) \rightarrow \tau(\mu)$  and  $F(\lambda; R) \rightarrow F(\lambda)$  as  $R \rightarrow \infty$ . We may assume that  $R$  is large enough that the term involving  $\tau(\mu)$  in (4.1) is unaffected by the change. If we can prove (4.3) for compactly supported  $d\tau(\mu)$ , then we obtain the general result by letting  $R \rightarrow \infty$ . Hence we proceed with the proof of (4.3) for the special case (4.4), and in which  $\tau(\mu)$  is constant outside some interval  $[-R, R]$ .

Introducing

$$f(\lambda, \mu) = \delta\{(\mu - \lambda)^2 + \delta^2\}^{-1},$$

we have  $\int_{-\infty}^{\infty} f(\lambda, \mu) d\lambda = \pi$  and

$$\int_{\lambda_1 + i\delta}^{\lambda_2 + i\delta} \text{Im } F(\lambda) d\lambda = \int_{-\infty}^{\infty} d\tau(\mu) \int_{\lambda_1}^{\lambda_2} f(\lambda, \mu) d\lambda,$$

since the integral on the right is finite in this restricted case of compactly supported  $d\tau(\mu)$ . Let

$$R(\gamma) = \int_{\gamma}^{\infty} d\tau(\mu) \int_{-\infty}^{\gamma} f(\lambda, \mu) d\lambda - \int_{-\infty}^{\gamma} d\tau(\mu) \int_{\gamma}^{\infty} f(\lambda, \mu) d\lambda,$$

so that a calculation leads to

$$\int_{\lambda_1 + i\delta}^{\lambda_2 + i\delta} \text{Im } F(\lambda) d\lambda = [\pi\tau(\lambda) + R(\lambda)]_{\lambda_1}^{\lambda_2}.$$

Now by (4.1)  $G(\lambda_1, \lambda_2) = [R(\lambda) - \delta \text{Re } F(\lambda + i\delta)]_{\lambda_1}^{\lambda_2}$ , so if we let  $S(\lambda) = R(\lambda) - \delta \text{Re } F(\lambda + i\delta)$ , then what we must prove is

$$-H(\lambda_1, \lambda_2) \leq S(\lambda_2) - S(\lambda_1) \leq H(\lambda_1, \lambda_2). \quad (4.5)$$

The identical argument in Atkinson's proof gives

$$S(\lambda) = \int_{\lambda}^{\infty} d\tau(\mu) h\left(\frac{\mu - \lambda}{\delta}\right) - \int_{-\infty}^{\lambda} d\tau(\mu) h\left(\frac{\lambda - \mu}{\delta}\right), \quad (4.6)$$

where  $h(\sigma) = \int_{\sigma}^{\infty} (1 + t^2)^{-1} dt - \sigma(1 + \sigma^2)^{-1}$ . However,  $0 \leq h(\sigma) \leq (\pi/2)(1 + \sigma^2)^{-1}$  and  $d\tau(\mu)$  is nonnegative. Thus (4.6) implies

$$S(\lambda) \leq (\pi/2) \int_{-\infty}^{\infty} d\tau(\mu) (1 + (\mu - \lambda)^2 \delta^{-2})^{-1}. \quad (4.7)$$

But in (4.4) we can compute  $\text{Im } F(\lambda + i\delta)$  by dropping the term  $\mu/(1 + \mu^2)$ , which is real, and the result is  $\text{Im } F(\lambda + i\delta) = \delta \int_{-\infty}^{\infty} d\tau(\mu) / ((\mu - \lambda)^2 + \delta^2)$ . Therefore (4.7) becomes

$$S(\lambda) \leq (\pi\delta/2) \text{Im } F(\lambda + i\delta). \quad (4.8)$$

In (4.6) we could have used the second term to bound  $S(\lambda)$  below, and the result would then read

$$S(\lambda) \geq -(\pi\delta/2) \text{Im } F(\lambda + i\delta). \quad (4.9)$$

Combining (4.8) and the reversed negative of (4.9) we get

$$S(\lambda_2) - S(\lambda_1) \leq (\pi\delta/2)[\operatorname{Im} F(\lambda_2 + i\delta) + \operatorname{Im} F(\lambda_1 + i\delta)].$$

The other half of (4.3) follows from switching the roles of (4.8) and (4.9) as used above. This completes the proof of the lemma.

Since the norm of a Hermitian matrix equals the maximum of the moduli of its eigenvalues, (4.3) gives the useful inequality

$$\|G\| \leq \|H\|. \quad (4.10)$$

## 5. Proof of main result

Assume the hypotheses of Theorem 1. Then by (4.1) we have for any  $\delta > 0$

$$\begin{aligned} \pi[\tau(\mu) - \tau_n(\mu)]_{\lambda_1}^{\lambda_2} &= \operatorname{Im} \int_{\lambda_1 + i\delta}^{\lambda_2 + i\delta} [F(\lambda) - F_n(\lambda)] d\lambda \\ &\quad - \delta[\operatorname{Re} F(\mu + i\delta) - \operatorname{Re} F_n(\mu + i\delta)]_{\lambda_1}^{\lambda_2} \\ &\quad - G(\lambda_1, \lambda_2) + G_n(\lambda_1, \lambda_2), \end{aligned} \quad (5.1)$$

where  $G_n$  is the function of (4.1) formed from  $\tau_n$  and  $F_n$ ; we will denote by  $H_n$  the corresponding function of (4.2). Each of the first two terms on the right of (5.1) can be made small by making  $n \rightarrow \infty$ , once  $\delta$  has been fixed, by the uniform convergence of  $F_n$  to  $F$  on the straight line path of integration, in the first instance, and at its endpoints in the second. We will thus pick  $\delta$  so that the expression  $G_n(\lambda_1, \lambda_2) - G(\lambda_1, \lambda_2)$  is also small for large  $n$ .

Since  $\lambda_1$  and  $\lambda_2$  are in either the continuous spectrum or resolvent set, we have  $\delta F(\lambda_j + i\delta) \rightarrow 0$ , as  $\delta \rightarrow 0$ , by the result in [11] quoted below (3.2). Consequently,  $H(\lambda_1, \lambda_2)$  can be made small by making  $\delta$  suitably small, and the same must hold for  $G(\lambda_1, \lambda_2)$  by (4.10). Turning then to the remaining term  $G_n(\lambda_1, \lambda_2)$  in (5.1), one has

$$\|G_n(\lambda_1, \lambda_2)\| \leq \|H_n(\lambda_1, \lambda_2)\| \leq \|H_n(\lambda_1, \lambda_2) - H(\lambda_1, \lambda_2)\| + \|H(\lambda_1, \lambda_2)\| \quad (5.2)$$

by the triangle inequality. As noted above, the term  $\|H(\lambda_1, \lambda_2)\|$  on the right of (5.2) has been made small by choosing  $\delta$  near 0. For this fixed  $\delta$  we drive the term adjacent to it to 0 by making  $n \rightarrow \infty$ . This completes the proof of the theorem.

## 6. Examples, extensions and applications

We close the paper with some examples illustrating Theorem 1, a discussion of two singular endpoint problems and some remarks on convergence of spectral functions.

(1) Consider the Sturm–Liouville equation  $-y'' + r(x)y - \varepsilon n(x)y = \lambda \omega(x)y$  on  $a \leq x < \infty$ , where the coefficients are differentiable,  $n(x) > 0$  and  $\omega(x) > 0$ . Writing the equation as a system, we may apply the methods of [17] and [15] to give conditions where the unperturbed ( $\varepsilon = 0$ ) spectrum is purely discrete and the perturbed spectrum ( $\varepsilon > 0$ ) is purely continuously differentiable. Briefly, we need

$r(x)$  to dominate  $\omega(x)$  for discreteness and  $n(x)$  to dominate both  $r(x)$  and  $\omega(x)$  for continuous perturbed spectrum; see [17] and [15] for technicalities. For example, when  $\omega(x) = x^\alpha$ ,  $r(x) = x^k$  and  $n(x) = x^m$  the desired situation prevails when  $-2 < \alpha < k < m < (\alpha + 1)$ .

(2) Similarly, there are criteria given discrete spectrum and continuous spectrum for the Dirac system

$$y' = \begin{pmatrix} 0 & \lambda(\alpha(x) + \varepsilon\hat{\alpha}(x)) - p(x) \\ -\lambda(\alpha(x) + \varepsilon\hat{\alpha}(x)) - p(x) & 0 \end{pmatrix} y, \quad a \leq x < \infty,$$

in the unperturbed and perturbed cases, respectively; see [17] and [16] for details. Requirements on the coefficients are similar to Example (1) above;  $p(x)$  must dominate  $\alpha(x)$  and  $\hat{\alpha}(x)$  must dominate  $p(x)$ . If  $\alpha(x) = x^\gamma$ ,  $p(x) = x^k$  and  $\hat{\alpha}(x) = x^\delta$ ,  $1 \leq x < \infty$ , then  $(-\frac{1}{2}) \leq \gamma < k < \delta$  will be sufficient.

(3) Our results apply to perturbations of the type  $-y'' + (\ln x - x^\varepsilon)y = \lambda y$ ,  $1 \leq x < \infty$ ,  $2 \geq \varepsilon \geq 0$ . As in Example (1), [15] shows there to be purely continuous spectrum over  $(-\infty, \infty)$  when  $\varepsilon > 0$ . For  $\varepsilon = 0$  the spectrum is discrete because  $\ln x \rightarrow \infty$  as  $x \rightarrow \infty$ ; see [3].

(4) Suppose now that (2.1) and (2.1)<sub>n</sub> are singular and of limit point type at both ends of  $(a, b)$ . Selfadjoint operators  $T$  and  $T_n$  are defined as in (2.6) except that no boundary condition of the type (iii), above (2.6), is necessary. The characteristic function for  $T$  is given by (see [14])

$$F(\lambda) = \begin{pmatrix} (M_a - M_b)^{-1} & (\frac{1}{2})(M_a - M_b)^{-1}(M_a + M_b) \\ (\frac{1}{2})(M_a + M_b)(M_a - M_b)^{-1} & M_b(M_a - M_b)^{-1}M_a \end{pmatrix}, \quad (6.1)$$

where  $M_a(\lambda)$  and  $M_b(\lambda)$  are the Titchmarsh–Weyl functions for  $T$  at  $x = a$  and  $x = b$ , respectively. The spectral function  $\tau(\lambda)$  is linked to  $F(\lambda)$  by precisely the same formulas (3.4) and (3.5), and the spectrum  $\sigma(T)$  may be characterized by the regular, pole and singular structure of  $F$  just as we described below (3.2) for the one endpoint problem. All this was done in [14]. If we assume that (2.8) holds, then it is true that  $F_n(\lambda) \rightarrow F(\lambda)$  on compact subsets of  $\text{Im}(\lambda) > 0$  by (6.1). So the proof of Theorem 1 would be valid for (6.1). The conclusion is therefore valid for two singular endpoint problems, including (1.1).

(5) We apply our results now to spectral families of scalar (limit point) operators  $Ly = -y'' + q(x)y$ ,  $0 \leq x < \infty$ , where with  $\omega(x) = p(x) = 1$  in (2.4),  $Ly$  appears as the first component in (2.6). The domain of  $L$  is defined in the usual way by assigning a boundary condition  $\cos \alpha y(0) + \sin \alpha y'(0) = 0$ . Letting  $\varphi(x, \lambda)$  be the solution of  $-y'' + q(x)y = \lambda y$  with  $\varphi(0, \lambda) = -\sin \alpha$  and  $\varphi'(0, \lambda) = \cos \alpha$ , the “transform” with respect to  $L$  of a function  $f \in L^2[0, \infty)$  is defined as ([3])

$$g_0(\lambda) = \int_0^\infty f(t)\varphi(t, \lambda) dt.$$

If  $\rho(\lambda)$  is the spectral function for  $L$ , then the eigenfunction expansion

$$f(t) = \int_{-\infty}^\infty g_0(\lambda)\varphi(t, \lambda) d\rho(\lambda)$$

converges in the mean to  $f \in L^2[0, \infty)$ ; see [3]. The spectral family ([25]) of  $L$

consists of the projection operators on  $L^2[0, \infty)$  given by

$$[S(\mu)f](t) = \int_{-\infty}^{\mu} g_0(\lambda) \varphi(t, \lambda) d\rho(\lambda).$$

Now let  $\{q(x)\}$  be a sequence of potentials such that  $q_n(x) \rightarrow q(x)$  uniformly on compact subsets of  $[0, \infty)$ . Then it follows from [25, pp. 283–286] that for the corresponding operators  $L_n$  (with same boundary condition at  $x = 0$  as  $L$ ),  $L_n \rightarrow L$  in the strong resolvent sense, and therefore that  $S_n(\mu) \rightarrow S(\mu)$  (i.e.,  $S_n(\mu)f \rightarrow S(\mu)f$  in  $L^2$  norm for all  $f \in L^2[0, \infty)$ ) where  $S_n(\mu)$  is the spectral family for  $L_n$ .

We now use Theorem 1 to prove pointwise convergence of the projection operators  $[S_n(\mu_2) - S_n(\mu_1)]f$  to  $[S(\mu_2) - S(\mu_1)]f$ , where  $\rho$  is continuous at  $\mu_1$  and  $\mu_2$ , for functions  $f$  of compact support. For if  $f$  is supported on  $[c, d]$ , then

$$\begin{aligned} [S_n(\mu_2) - S_n(\mu_1)]f(x) &= \int_{\mu_1}^{\mu_2} \left[ \int_c^d f(t) \varphi_n(t, \lambda) dt \right] \varphi_n(x, \lambda) d\rho_n(\lambda) \\ &= \int_c^d f(t) \left[ \int_{\mu_1}^{\mu_2} \varphi_n(t, \lambda) \varphi_n(x, \lambda) d\rho_n(\lambda) \right] dt. \end{aligned} \quad (6.2)$$

By an extension of the proof of the Helly–Bray Theorem in [26, p. 31],  $\int_{\mu_1}^{\mu_2} \varphi_n(t, \lambda) \varphi_n(x, \lambda) d\rho_n(\lambda) \rightarrow \int_{\mu_1}^{\mu_2} \varphi(t, \lambda) \varphi(x, \lambda) d\rho(\lambda)$ ,  $n \rightarrow \infty$ , uniformly for  $c \leq t \leq d$ , in view of Theorem 1. Hence (6.2) approaches  $[S(\mu_2) - S(\mu_1)]f(x)$ , as  $n \rightarrow \infty$ , and this is the desired result.

The above proof shows that  $(S_n(\mu)f)(x) \rightarrow (S(\mu)f)(x)$ , pointwise for functions  $f$  of compact support, if the spectra of  $L$  and  $L_n$  are bounded below by a single number, say  $d\rho_n(\lambda) = d\rho(\lambda) = 0$  on  $-\infty < \lambda < \lambda_0$ . For the integrals from  $\mu_1$  to  $\mu_2$  in (6.2) could then be replaced by the integrals from  $-\infty$  to  $\mu$ . To prove the analogous result for spectra unbounded from below would require uniform smallness of the integrals  $\int_{-\infty}^{\lambda_0} \varphi_n(t, \lambda) \varphi_n(x, \lambda) d\rho_n(\lambda)$  for large negative  $\lambda_0$  and  $n = 1, 2, 3, \dots$ . While we do know that each  $\rho_n(\lambda)$  is exponentially small as  $\lambda \rightarrow -\infty$  ([2, p. 18]) and that the terms  $\varphi_n(t, \lambda) \varphi_n(x, \lambda)$  are exponentially bounded as  $\lambda \rightarrow -\infty$ , it is not clear that the resulting bounds are independent of  $n$ .

## REFERENCES

- [1] F. V. ATKINSON, *Discrete and Continuous Boundary Problems*, Academic Press, New York, 1964.
- [2] F. V. ATKINSON, *On the asymptotic behavior of the Titchmarsh–Weyl  $m$ -coefficient and the spectral function for scalar second-order differential expressions*, in *Ordinary and Partial Differential Equations*, pp. 1–27, W. N. Everitt and B. D. Sleeman, editors, Springer-Verlag Lecture Notes in Mathematics, vol. 964, Berlin, 1982.
- [3] E. A. CODDINGTON and N. LEVINSON, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
- [4] C. C. CONLEY and P. A. REJTO, *Spectral concentration II—general theory*, in *Perturbation Theory and its Applications in Quantum Mechanics*, pp. 129–144, C. H. Wilcox, Editor, John Wiley and Sons, Inc., New York, 1966.
- [5] W. A. COPPEL, *Disconjugacy*, Springer-Verlag Lecture Notes in Mathematics, vol. 220, Berlin, 1971.
- [6] L. C. DETWILER and J. R. KLAUDER, *Supersingular quantum perturbations*, *Phys. Rev.* 11 (1975), 1436–1441.
- [7] E. HARRELL and B. SIMON, *The mathematical theory of resonances whose widths are exponentially small*, *Duke Math. J.* 47 (1980), 845–902.



- [8] E. HARRELL, N. CORNGOLD and B. SIMON, *The mathematical theory of resonances whose widths are exponentially small, II*, J. Math. Anal. Appl. 99 (1984), 447–457.
- [9] P. HARTMAN, *Ordinary Differential Equations*, 2nd. ed., Birkhäuser, Boston, 1982.
- [10] D. B. HINTON and J. K. SHAW, *On Titchmarsh–Weyl  $m$ -functions for linear Hamiltonian systems*, J. Diff. Eqs. 40 (1981), 316–342.
- [11] D. B. HINTON and J. K. SHAW, *On the spectrum of a singular Hamiltonian system*, Quaes. Math. 5 (1982), 29–81.
- [12] D. B. HINTON and J. K. SHAW, *Titchmarsh–Weyl theory for Hamiltonian systems*, in *Spectral Theory of Differential Operators*, pp. 219–231, I. W. Knowles and R. T. Lewis, editors, North-Holland, New York, 1981.
- [13] D. B. HINTON and J. K. SHAW, *On boundary value problems for Hamiltonian systems with two singular points*, SIAM J. Math. Anal. 15 (1984), 272–286.
- [14] D. B. HINTON and J. K. SHAW, *On the spectrum of a singular Hamiltonian system, II*, submitted.
- [15] D. B. HINTON and J. K. SHAW, *Absolutely continuous spectra of second order differential operators with short and long range potentials*, to appear in SIAM J. Math. Anal.
- [16] D. B. HINTON and J. K. SHAW, *Absolutely continuous spectra of Dirac systems with long range, short range and oscillating potentials*, Quart. J. Math. Oxford (2), 36 (1985), 183–213.
- [17] D. B. HINTON and J. K. SHAW, *Dirac systems with discrete spectra*, submitted.
- [18] B. M. LEVITAN and I. S. SARGSIAN, *Introduction to Spectral Theory: Self-adjoint Ordinary Differential Operators*, Translations of Mathematical Monographs 39, American Mathematical Society, Providence, 1975.
- [19] J. B. MCLEOD, *Spectral concentration I – the one-dimensional Schrödinger operator*, in *Perturbation Theory and its Applications in Quantum Mechanics*, pp. 119–128, C. H. Wilcox, editor, John Wiley and Sons, Inc., New York, 1966.
- [20] M. REED and B. SIMON, *Methods of Modern Mathematical physics*, vol. 4, Academic Press, New York, 1978.
- [21] F. S. ROFE-BEKETOV and V. I. KOGAN, *On square-integrable solutions of symmetric systems of differential equations of arbitrary order*, Proc. Roy. Soc. Edin. 74 A(1974), 5–39.
- [22] E. C. TITCHMARSH, *Some theorems on perturbation theory III*, Proc. Roy. Soc. A207 (1951), 321–328.
- [23] E. C. TITCHMARSH, *Some theorems on perturbation theory IV*, Proc. Roy. Soc. A210 (1951), 30–47.
- [24] E. C. TITCHMARSH, *Some theorems on perturbation theory V*, J. Analyse Math. 4 (1954/56), 187–208.
- [25] J. WEIDMAN, *Linear Operators in Hilbert Space*, Graduate Texts in Mathematics, vol. 68, Springer-Verlag, Berlin, 1980.
- [26] D. V. WIDDER, *The Laplace Transform*, Princeton University Press, Princeton, New Jersey, 1946.