

Zeitschrift: Helvetica Physica Acta
Band: 59 (1986)
Heft: 4

Artikel: Rapporteur's report, session (E)
Autor: Steffens, E.
DOI: <https://doi.org/10.5169/seals-115717>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Rapporteur's report, Session (E), E. Steffens:

Cold atomic beam techniques

The discussion following the talks of this session were partly centered around the question to which extent the promises of the cold atomic beam technique which were expressed during the Ann Arbor Workshop [1] five years ago have been fulfilled in practice.

In the relevant temperature range below liquid nitrogen temperature clean surfaces like quartz or teflon do not work any more because of surface recombination. It is believed that surfaces exposed to atomic hydrogen have to be covered by frozen layers of gas or by liquid He films. At very low temperature of about 0.5K and high field of $> 5T$ the magnetic energy $\mu_B^2 B$ becomes large compared to the energy and magnetic trapping of certain substates occurs.

Most of the effort in the last 5 years has been concentrated on the production of cold hydrogen beams which are then state-selected in the usual way (see Session (J)), whereas only recently a Michigan-MIT-CERN-collaboration has started to study magnetic trapping for polarized jet target applications.

1. Free cold hydrogen beams ($5K < T_{acc} < 77K$):

Results from ETH, Kyushu and SIN on beam cooling indicate a gain in density at the ionizer proportional to about $1/\sqrt{T}$, that is a factor of 4 for 20K compared to room temperature beams. Frozen layers of H_2 (below $\sim 20K$) or N_2 prevent recombination. At SIN, stable operation of the atomic beam for more than two weeks has been achieved with a small admixture of N_2 to the hydrogen feed gas.

Beam cooling with accommodator temperatures as low as 5K has been successfully done at BNL (as reported by A. Hershcovitch). Here the "bad" temperature range $T < 70K$ is carefully avoided by feeding the atomic hydrogen through a heated teflon tube to the cold accommodator which is separated by a narrow slit. Intensities in excess of $2 \cdot 10^{18} H^0/sr s$ have been obtained.

T. Niinikoski presented calculations which showed that the density achievable in the focus after the spin selector is limited by gas dynamical phenomena. P. Schmelzbach concluded that these limits are very encouraging because they are two to three orders of magnitude above the present densities.

2. Strong cooling and magnetic trapping

R. Raymond reported on the development undertaken by a Michigan-MIT-CERN-collaboration to produce a slowly pulsed polarized jet target using microwave extraction from a 5 Tesla solenoid. The density aimed at is in excess of $10^{14} H^0/cm^2$. The duty cycle will be in the order of 10% (1/4 sec. every few seconds). The cryostat is being tested and most of the equipment has been obtained. D. Kleppner expressed his confidence that this development will also be useful for d.c. beam.

A very interesting approach to overcome surface recombination at extremely low temperatures has been presented by H. Hess (MIT). Instead of trapping states a+b (the two lower states) in the field maximum at the walls,

he intends to trap in the field minimum in free space, e.g. on the axis of the system. A combination of quadrupole field and mirror coils for radial and axial trapping is required. Wall collisions are only necessary for thermalizing, but not for confining the gas. This development might lead to spin polarized ^{14}H -gas stored in relatively weak trapping fields with densities of $10^{14}/\text{cm}^3$.