

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 59 (1986)  
**Heft:** 4

**Artikel:** Secretary's report, session (J)  
**Autor:** Huttel, E.  
**DOI:** <https://doi.org/10.5169/seals-115741>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Secretary's report, Session (J), E. Huttel:

1. Report W. Gruebler

P. Schiemenz: What is the acceptance of your acellerator compared to the emittance of the atomic beam ?

W. Gruebler: The acceptance of the acellerator has to be compared with the emittance of the ionized beam which depends only on the properties of the ionizer.

2. Report J. Lemaire

S. Jaccard: How did find the quantity of the buffer gas needed ?

J. Lemaire: By optimization, the exact values are not known.

P. Schmelzbach (comment): At our source the amount of the buffer gas is 1-2 % N<sub>2</sub>.

P. Schmelzbach: What is the optimum atomic-beam temperature for your sextupol magnet ?

A. Hershcovitch (comment): The reason for the fact that you do not optain velocities lower than 200 m/sec might be a contact of the plasma in the dissociator to the nozzle.

3. Report W. Kubischa

A. Hershcovitch: How did you measure the dissociation degree ?

W. Kubischa: The ratio of atoms and molecules was measured with a quadrupole mass spectrometer.

4. Discussion

S. Jaccard: Has any one what to say about low-energy polarimeter ?

J. Arvieux: A polarimeter [1] based on the  $^2\text{H}(\vec{d}, p) ^3\text{H}$ . reaction [2] is used for both vector and tensor-polarized deuterons. The polarimeter could now be calibrated by an absolute minimum in the T<sub>20</sub> analyzing power of the  $\vec{d} + p \rightarrow ^3\text{He} + \pi^0$  reaction ( $E_d = 650$  MeV,  $\theta_{CM} = 180^\circ$ ) [3].

G. Clausnitzer: At Gießen a polarimeter [4] based on the  $\text{Li}(\vec{p}, ^3\text{He}) ^4\text{He}$  is used for monitoring the proton polarization and a polarimeter [5] based on the  $^3\text{He}(\vec{d}, p) ^4\text{He}$  reaction is used for tensor-polarized deuterons. We have now remeasured the vector analyzing power of the  $^2\text{H}(\vec{d}, p) ^3\text{H}$  reaction [6], which shows sufficient analyzing power down to 30 keV. A deuterated para-polyphenyl layer on a thin carbon foil is used as target. A measurement of the tensor analyzing power for the same reaction will be performed soon.

J. Alessi: A polarimeter [7] based on the measurement of circular polarized Lyman  $\alpha$  light is used at Brookhaven. The low energy beam passes through a thin carbon foil, leaving fractions of the beam in the  $^2\text{P}$  state. The circular polarization of the decaying photons are used as monitor.

[1] A. Boudard et al., Rapport CEA-N-2330 (1983) 204

[2] B.P. Ad'yasevich et al., Sov. J. Nucl. Phys. 33 (3) (1981) 313 and references given there

[3] J. Arvieux priv. communication

[4] G. Hermann, Diplomarbeit, Gießen

[5] G. Reiter, Diplomarbeit, Gießen

[6] E. Pfaff et al., 6th Int. Symp. on Pol. Phen. on Nucl. Phys. Osaka (1985)

[7] K.P. Schüler et al., 6th Int. Symp. on Pol. Phen. on Nucl. Phys. Osaka (1985)