

Zeitschrift: Helvetica Physica Acta
Band: 59 (1986)
Heft: 4

Artikel: Rapporteur's report, session (F)
Autor: Heeringa, W.
DOI: <https://doi.org/10.5169/seals-115753>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Rapporteur's report, session (F), W. Heeringa:

The first talk in this session was given by Glättli, Saclay, about polarizing nuclei of elements heavier than hydrogen with dynamic methods. The list comprised over 20 nuclides, which have been polarized mainly for measurements of pseudo magnetic moments or for nuclear magnetic ordering studies. The problem of creating free electrons in the various substances was discussed. In this field still a lot of uncertainties and surprises occur. The difficulties inherent to NMR measurements for determining the polarization were briefly discussed. Other methods were also mentioned, upon a question from the audience, namely measuring the nuclear magnetization with a SQUID or employing spin dependent properties of nuclear reactions, e.g. transmission by polarized neutrons.

After this, Arvieux, Saclay, came up with a short note advertising the use of polarized nuclei in heavy-ion nucleus-nucleus collisions with deformed nuclei at high energies. Higher densities can be reached during these collisions when both nuclei are oriented along the beam direction.

The next talk was by Konter, SIN, about brute force polarization. He gave a resumé about the requirements and the methods available to achieve brute-force polarization. The most recent development is the employment of enhanced nuclear refrigeration, which yields a high cooling power below 10 mK. The largest samples and highest polarizations are produced by the solid-state physics community to reach temperatures in the μK-region via nuclear demagnetization. After the talk some questions were asked about nuclear orientation thermometry with ^{60}Co .

The final talk in this session was by Vandeplassche, Leuven, about nuclear polarization by pick-up of polarized electrons. The aim of this work is to study the decay of radioactive isotopes with such short life-times, that they cannot be polarized after on-line implantation, because the spin-lattice relaxation time is longer than the decay time. The solution is to polarize the nuclei before implantation by pick-up of polarized electrons from nickel single crystals. The present problem is that the divergence of the beam after electron pick-up is too large. In the discussion there were questions about the usefulness of this principle to make a polarized source. At present up to 15 nA of polarized beam can be achieved. This can be increased by using more or larger Ni crystals. The divergence problem remains, and anyway one would have to re-ionize in order to make a polarized source.