

Zeitschrift: Helvetica Physica Acta
Band: 62 (1989)
Heft: 4

Artikel: Equation of hydrostatic equilibrium and temperature dependent gravitational constant
Autor: Massa, Corrado
DOI: <https://doi.org/10.5169/seals-116040>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Equation of hydrostatic equilibrium and temperature dependent gravitational constant

By Corrado Massa

Via Fratelli Manfredi 55 42100–Reggio Emilia Italy

(21. XII. 1988 revised, 7.II.1989)

Abstract. The Oppenheimer–Volkoff (shortly OV) equation of hydrostatic equilibrium describes the balance between gravitational force and pressure gradient in a self-gravitating perfect fluid. In the following, a generalized form of the OV equation is obtained by the assumption that the gravitational constant G is temperature-dependent according to the law $G = G_0(1 - bT^2)^{-1}$ where b is a positive constant. Such a law is required by a variety of gauge theories used in the unification program. A direct consequence of the generalized OV equation is the existence of an upper bound $T < (2/b)^{1/2}$ on the temperature of any self-gravitating radiation field.

A variety of gauge theories used in the unification program imply a temperature-dependent gravitational constant G given by:

$$G = G_0(1 - bT^2)^{-1} \quad (1)$$

where T is the temperature, G_0 is the zero temperature value of G , very close to the currently observed value $6.67 \cdot 10^{-8} \text{ cm}^{-3} \text{ g}^{-1} \text{ s}^{-2}$, and b is a positive constant whose numerical value depends on the details of the theory concerned [1–3]. The simplest way to fit the law (1) on the framework of Einstein's gravitational theory is to write the gravitational field equations [1, 4]

$$R^{ik} - (\frac{1}{2})g^{ik}R = 8\pi GT^{ik} \quad (2)$$

where $G = G(T)$ according to equation (1). R^{ik} is the Ricci tensor, g_{ik} is the fundamental tensor, $R = g^{ik}R_{ik}$ is the spacetime curvature scalar, and T^{ik} is the energy tensor. In this paper, latin indices run from 0 to 3, greek indices run from 1 to 3; commas denote partial derivative and semicolons denote covariant derivative with respect to the spacetime coordinates x^i ; dots denote differentiation with respect to x^0 , and apices denote differentiation with respect to x^1 .

By equation (2) and by the Bianchi identity $R_{i;k}^k = (\frac{1}{2})R_{,i}$ we have $(GT^{ik})_{;k} = 0$, namely

$$\dot{G}T^{00} + Gt^0 = 0 \quad (3)$$

$$G'T^{11} + Gt^1 = 0 \quad (3a)$$

where $t^i = T_{;k}^{ik}$.

Consider now a self-gravitating mass described by the energy tensor of a perfect fluid:

$$T^{ik} = (\rho + P)U^i U^k - Pg^{ik} \quad (4)$$

where ρ is the rest frame mass density and P the scalar pressure; U^i is the 4-velocity of the local rest frame of the fluid.

Consider a static, spherically symmetric field; thus:

$$U^\beta = 0, \quad ds^2 = e^\nu(c dt)^2 - e^\lambda dr^2 - d\Omega^2 \quad (5)$$

where ds^2 is the spacetime line-element, ν and λ are functions of r only, and $d\Omega^2$ means $(r d\vartheta)^2 + (r \sin \vartheta d\varphi)^2$ where $(ct, r, \vartheta, \varphi) = x^i$ are the Schwarzschild coordinates (r is the radial coordinate).

The normalization of 4-velocity $U^i U_i = 1$ determines $U^0 = e^{-\nu/2}$, and, by (4)(5), $T^{00} = \rho e^{-\nu}$, $T_0^0 = \rho$, $T^{11} = Pe^{-\lambda}$. For $\dot{G} = 0$, equation (3) gives $t^0 = 0$ (neither creation nor destruction of matter) and we have to consider equation (3b) only. Equation (3b) with $T^{00} = \rho e^{-\nu}$ and $T^{11} = Pe^{-\lambda}$ gives:

$$(G'/G) + P' + (\frac{1}{2})(\rho + P)\nu' = 0 \quad (6)$$

The (00) component of the field equation (2) in the proper reference frames of fluid elements is;

$$(1/r^2)(d/dr)[r(1 - e^{-\lambda})] = 8\pi G\rho,$$

which integrated gives;

$$\left(\frac{1}{4\pi}\right)m(r) = \int_0^r G\rho r^2 dr \quad (7)$$

where $2m(r) = r(1 - e^{-\lambda})$.

The (11) component is

$$-(1/r^2) + (1/r^2)e^{-\lambda} + (1/r)e^{-\lambda}\nu' = 8\pi GP,$$

that gives:

$$(\frac{1}{2})\nu' = (m + 4\pi PG r^3)(r^2 - 2mr)^{-1} \quad (8)$$

Equations (1) (8) lead to;

$$P' = -(\rho + P)(m + 4\pi PG r^3)(r^2 - 2mr)^{-1} - PG'/G \quad (9)$$

with $m = m(r)$ given by equation (7). Equation (9), with $G = G_0 = \text{constant}$, reduces to the usual OV equation for a perfect fluid [5-7]. Consider now equation (1), which gives $G'/G = (T'/T)2bT^2(1 - bT^2)^{-1}$. If the fluid is a radiation field with the equation of state $\rho = 3P = aT^4$ (a = the radiation constant) then equation (9) reads:

$$AT'/T + [Bm + (4\pi/3)aT^4 r^3 G_0](BC)^{-1} = 0, \quad (10)$$

where;

$$A = 1 + (\frac{1}{2})bT^2 B^{-1}, \quad (10a)$$

$$B = 1 - bT^2, \quad (10b)$$

$$C = r^2 - 2mr. \quad (10c)$$

A direct consequence of equation (10) is the existence of an upper bound on the temperature T of the fluid. To see this fact, assume $T > 1/\sqrt{b}$ at some point r_0 inside the fluid, namely $B < 0$. For equation (1), the equation (7) gives $m(r_0) < 0$, viz. (for equation (10c)) $C > 0$, and equation (10) leads to

$$AT' > 0 \quad (11)$$

T' is certainly negative. Indeed, if $T' > 0$, then $A > 0$, and (for equations (10,a,b) and for $B < 0$) $T > (2/b)^{1/2}$; all values between $(1/b)^{1/2}$ and $(2/b)^{1/2}$ would be forbidden, and T could not be continuous.

Put $T' < 0$ into equation (11) and obtain $A < 0$, viz. $T < (2/b)^{1/2}$ that is an upper bound on T .

A more severe bound, namely $T < (1/b)^{1/2}$, can be obtained via the quite different procedure described in Reference (1). In the special case of the Hawking black hole temperature, the bound $T < (1/b)^{1/2}$ can be derived by the very simple procedure of Reference (2). My thanks to an anonymous referee of this Review which brought Reference (3) to my attention.

REFERENCES

- [1] C. MASSA, Helv. Phys. Acta (1989) to be published, and references cited therein.
- [2] P. C. W. DAVIES, Phys. Lett. *101B* 399 (1981).
- [3] P. C. W. DAVIES, "Temperature-Dependent G and Black Hole Thermodynamics", in the Proceedings of the 2nd Moscow Quantum Gravity Seminar (ed. Markov & West, Plenum 1984).
- [4] A. K. RAYCHAUDHURI, and B. BAGCHI, Phys. Lett. *124B*, 168 (1983).
- [5] J. R. OPPENHEIMER, and G. VOLKOFF, Phys. Rev. *55*, 374 (1939).
- [6] C. W. MISNER, K. S. THORNE, and J. A. WHEELER, Gravitation (W. H. Freeman and Company, San Francisco 1973) Chapter 23, Section 5, p. 605.
- [7] R. D. SORKIN, R. M. WALD, and Z. Z. JIU, Gen. Rel. Grav. *13*, 1127 (1981).