

Zeitschrift: Helvetica Physica Acta
Band: 62 (1989)
Heft: 6-7

Artikel: Limit of zero-width conduction band for the extended Falicov-Kimball model
Autor: Marinaro, M. / Noce, C. / Romano, A.
DOI: <https://doi.org/10.5169/seals-116090>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LIMIT OF ZERO-WIDTH CONDUCTION BAND
FOR THE EXTENDED FALICOV-KIMBALL MODEL

M. Marinaro, C. Noce, A. Romano

Dipartimento di Fisica Teorica e Unità GNSM e CISM
Università di Salerno, I-84081 Baronissi (SA), Italy

Abstract: The extended Falicov-Kimball model has been exactly solved in the limit of zero-width conduction band. Preliminary results about the occurrence of both first order and second order metal-insulator transitions are presented.

1. Introduction

The transition metal and rare earth compounds exhibit a wide variety of anomalous electronic and magnetic properties, which arise from the interaction between highly correlated and itinerant electrons. For example, under variation of external conditions (temperature, pressure, alloying) these compounds undergo first order as well as second order transitions from insulator (or semiconductor) to metal. The model proposed by Falicov and Kimball [1] and the extended version of it (EFK) [2,3] seem to be well suited to explain the behaviour of these systems.


2. Model and results

The single site EFK Hamiltonian is

$$H = \sum_k \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \sum_\sigma \epsilon f_\sigma^\dagger f_\sigma + U f_\uparrow^\dagger f_\uparrow f_\downarrow^\dagger f_\downarrow + G \sum_\sigma f_\sigma^\dagger f_\sigma c_{\sigma'}^\dagger c_{\sigma'} + V \sum_\sigma (c_\sigma^\dagger f_\sigma + f_\sigma^\dagger c_\sigma)$$

where standard second quantization notation is used. In the limit of vanishing conduction bandwidth the model has finite degrees of freedom and thus can be exactly solved. In a previous paper [4] the n-point Green functions have been calculated by using the path integral formalism. From two-point Green functions for localized

electrons we have calculated the number of f-electrons per site. It is reported in figure as a function of ϵ for $T=70^\circ\text{K}$, $U=2$ eV $V=0.2$ eV, and different values of G . Due to the well known linear relation between pressure and ϵ , this plot shows the dependence of the valence of the f-electrons from the pressure. We see that the



electronic transition is driven by the strength of the Coulomb interaction between c and f-electrons: increasing the value of G from 0 to 1.5 eV the transition passes continuously from second order to first order.

We stress that in our approach the effect of G , V and U are all included in a non-perturbative way. In this sense our results are more general than the ones reported in ref.[3] where the terms proportional to V and G are treated in mean field approximation and the limit of infinite U is considered.

Besides, we point out that the atomic limit considered here does not qualitatively affect the results about the order of the transition, as noted for example in ref.[3], and it still retains the essential features of the intermediate valence properties of some real systems [5]. More details will be presented elsewhere.

4. References

- [1] L.M.Falicov, J.C.Kimball, Phys. Rev. Lett. **22**, 997 (1969).
- [2] M.Avignon, S.K.Ghatak, Solid State Commun. **16**, 1243 (1975).
- [3] C.E.T.Gonçalves da Silva, L.M.Falicov, Solid State Commun. **17**, 1521 (1975).
- [4] C.Noce, A.Romano, J. Phys.: Cond. Matt., in press.
- [5] J.M.Coey, S.K.Ghatak, M.Avignon, F.Holtzberg, Phys. Rev. **B14**, 3744 (1976); S.M.Shapiro, J.D.Axe, R.J.Birgeneau, J.M.Lawrence, R.D.Parks, Phys. Rev. **B16**, 2225 (1977).