

Zeitschrift: Helvetica Physica Acta
Band: 62 (1989)
Heft: 6-7

Artikel: Magnetic correlations and antiferromagnetic order in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$
Autor: Borsa, F. / Corti, M. / Rega, T.
DOI: <https://doi.org/10.5169/seals-116101>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

MAGNETIC CORRELATIONS AND ANTIFERROMAGNETIC ORDER IN $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$

F.Borsa, M.Corti, T.Reggi, A.Rigamonti and J.Ziolo

Department of Physics A.Volta, Pavia (ITALY)

Abstract: From ^{139}La and ^{63}Cu NQR relaxation it is found that the Cu^{++} long-range in-plane magnetic correlation is strongly limited by mobile charge defects introduced by the Sr doping. Both NQR relaxation and specific heat measurements indicate that at low temperature the antiferromagnetic order is restored at $T_N(x)$ as a consequence of a freezing of the mobile defects.

The ^{139}La and ^{63}Cu spin-spin and spin-lattice relaxation measurements for x ranging from 0 to $x=0.3$ and in the temperature range 1.6–450 K, when analyzed in terms of the possible relaxation mechanisms allow one to conclude that over most of the x and T range the relaxation is controlled by the $\text{Cu}^{++} - \text{Cu}^{++}$ in plane (2D) magnetic correlation length ξ (1).

An x and T dependence of ξ , which is unconventional with respect to the behavior of 2D doped Heisenberg paramagnets is deduced through standard relationships of the relaxation rate T_1^{-1} vs ξ . On the basis of an interpretative model whereby the degree of magnetic correlation that would result from the strength of the $\text{Cu}^{++} - \text{Cu}^{++}$ exchange coupling is controlled by an effective correlation time, one derives

$$1) \quad \xi(x, T) = \frac{1}{\sqrt{n(x, T)}} = \frac{1}{\sqrt{x}} e^{b/2xT} \quad (\text{in lattice units})$$

where $b/2x$ is an effective activation energy with $b \simeq 0.9$ K that has been deduced by the low T behavior of T_1^{-1} for various x (see Fig.1).

When the magnetic order is restored at low temperature as a consequence of the decrease of the concentration of the mobile defects $n(x, T)$ (see Eq.1), one observes a peak in (T_1^{-1}) and a small anomaly in the specific heat $C_p(T)$. The maximum value $(T_1^{-1})_{\text{max}}$ is about the same for all x although $T_N(x)$ ranges from ~ 300 to ~ 2 K. This is taken as an indication that the ordering is triggered by a 2D correlation length ξ_{2D}^* which is about the same at $T \leq T_N$, regardless the x value. This unexpected result could imply an interplane coupling J' which sharply decreases with x or which is not in competition with the thermal fluctuations.

No information can be derived from the NQR data on the character of the 3D order, i.e. antiferromagnetic vs spin-glass. However the anomalies in C_p around T_N are in favour of a 3D antiferromagnetic state. In Fig.2 we show the phase diagram for our samples as deduced from a variety of measurements.





Fig.1 - (a) Temperature behaviours of the ^{139}La relaxation rates for different x in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$. (b) Evaluation of the activation energy for T_1 ; note that according to the interpretative model $T_1^{-1} \propto n^{-1} = \frac{1}{2} \left(\frac{1}{T} - \frac{1}{T_N} \right)$

Fig.2 Phase diagram for $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4-y}$ as deduced in the present work ($\diamond, \square, \circ$) and published data ($x, \bullet, \Delta$)



References

1. F.Borsa, M.Corti, T.Regia, A.Rigamonti, Nuovo Cimento D (in press)