

Zeitschrift: Helvetica Physica Acta
Band: 62 (1989)
Heft: 6-7

Artikel: Dynamics of fluxons in field-cooled high- T_c superconductors
Autor: Ciccarello, I. / Fazio, C. / Guccione, M.
DOI: <https://doi.org/10.5169/seals-116103>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 04.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

DYNAMICS OF FLUXONS IN FIELD-COOLED HIGH- T_c SUPERCONDUCTORS

I. Ciccarello, C. Fazio, M. Guccione and M. Li Vigni, Istituto di Fisica dell'Università, via Archirafi 36, 90123 Palermo, Italy.


Abstract: Evidence is given that for increasing and decreasing applied fields critical states are developed in field-cooled high- T_c ceramic superconductors. In samples in the critical state a microwave magnetic field parallel to the applied field induces a magnetization with odd as well as even Fourier components.

The interaction of microwave fields with field-cooled (FC) single-phase ceramic $\text{YBa}_2\text{Cu}_3\text{O}_7$ has been studied by investigating the second-harmonic (SH) response to the driving field.

The experiments have been performed by exposing the samples to a dc field H_0 and an ac field $H(\omega)$ at an angle θ with respect to H_0 . The SH signal radiated by the samples has been investigated. It depends on both H_0 and θ . The signal is phase coherent. The phase as a function of H_0 and θ has also been investigated.

Fig.1a shows the SH signal intensity as a function of H_0 in the range of fields 1200-1800 Oe. The intensity is independent of H_0 as long as H_0 is increased (or decreased) steadily. Minima are observed when the magnetic field sweep is reversed, independently of the values of H_0 in which the inversions are operated. Fig.1b shows the phase of the SH signal as a function of H_0 . The phase, at a given value of H_0 , may differ by 180° depending on the way H_0 has been attained, at increasing or decreasing fields. On increasing H_0 the phase remains constant as long as H_0 is increased, from D to A. It remains still constant at the same value when H_0 is decreased from A to B, while a 180° variation is observed on crossing B. If the field is further decreased the phase remains constant at the new value until point D is reached.

The results are consistent with the assumption that a critical state



is developed in FC Y-Ba-Cu-O by an increasing (or decreasing) applied field. Supposing valid the Bean's assumption of constant J_c in superconductors⁽¹⁾, because of the rigidity of the fluxon lattice the response of the sample, in a critical state, to a microwave field is expected to be uneven during a period of the wave: for increasing fields the induction flux is essentially influenced only during the negative semiperiod, so that a magnetization with both odd and even Fourier components results⁽²⁾.

According to this picture a theory has been developed⁽²⁾

which accounts quite well for i) SH signal intensity independent of H_0 for constant field sweep; ii) minima at the inversion points of the field sweep; iii) 180° phase variation at those values of H_0 in which the critical state is removed (e.g., points B and D of Fig.1); iv) angular dependence of both intensity and phase of the SH signal.

References

- (1) C. P. Bean, Rev. Mod. Phys. **36**, 31 (1964).
- (2) I. Ciccarello, C. Fazio, M. Guccione and M. Li Vigni, Physica C, submitted.