

<b>Zeitschrift:</b>	Helvetica Physica Acta
<b>Band:</b>	64 (1991)
<b>Heft:</b>	5
<b>Artikel:</b>	A remark on anisotropic superconducting states
<b>Autor:</b>	Feldmann, Joel / Knörrer, Horst / Trubowitz, Eugene
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-116318">https://doi.org/10.5169/seals-116318</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A Remark on Anisotropic Superconducting States

Joel Feldman\*  
Department of Mathematics  
University of British Columbia  
Vancouver, B.C. V6T 1Y4  
CANADA

Horst Knörrer  
Eugene Trubowitz  
Mathematik  
ETH-Zentrum  
CH-8092 Zürich  
SWITZERLAND

(23. III. 1991)

### Abstract

We show that, in three dimensions, there are no nontrivial, isotropic, unitary solutions of the gap equation for angular momentum greater than one, while in two dimensions they exist in all angular momentum sectors.

---

\* Research supported in part by the Natural Science and Engineering Research Council of Canada

Consider the many Fermion system in three dimensions characterized by the effective potential

$$\begin{aligned} \mathcal{G}(\psi^\epsilon, \bar{\psi}^\epsilon) &= \log \frac{1}{Z} \int e^{-\lambda \mathcal{V}(\psi + \psi^\epsilon, \bar{\psi} + \bar{\psi}^\epsilon)} d\mu_C(\psi, \bar{\psi}), \\ \mathcal{V}(\psi, \bar{\psi}) &= \frac{1}{2} \sum_{a_i \in \{\uparrow, \downarrow\}} \int \prod_{i=1}^4 \frac{d^4 k_i}{(2\pi)^4} (2\pi)^4 \delta(k_1 + k_2 - k_3 - k_4) \delta_{a_1, a_3} \delta_{a_2, a_4} \\ &\quad \langle k_1, k_2 | V | k_3, k_4 \rangle \bar{\psi}(k_1, a_1) \bar{\psi}(k_2, a_2) \psi(k_4, a_4) \psi(k_3, a_3), \end{aligned}$$

where  $d\mu_C(\psi, \bar{\psi})$  is the fermionic Gaussian measure in the Grassmann variables

$$\{\psi(\xi), \bar{\psi}(\xi) | \xi = (\tau, \mathbf{x}, \sigma), \tau \in \mathbf{R}, \mathbf{x} \in \mathbf{R}^3, \sigma \in \{\uparrow, \downarrow\}\}$$

with covariance

$$\begin{aligned} C(\xi_1, \xi_2) &= \langle \psi(\xi_1) \bar{\psi}(\xi_2) \rangle \\ &= \delta_{\sigma_1, \sigma_2} \int \frac{d^{d+1} k}{(2\pi)^{d+1}} \frac{e^{i \langle k, \xi_1 - \xi_2 \rangle_-}}{ik_0 - e(\mathbf{k})} \\ \langle k, (\tau, \mathbf{x}) \rangle_- &= -k_0 \tau + \mathbf{k} \cdot \mathbf{x}, \quad \mathbf{k} = (k_0, \mathbf{k}) \\ e(\mathbf{k}) &= \frac{\mathbf{k}^2}{2m} - \mu. \end{aligned}$$

and where the two-body interaction  $\langle k_1, k_2 | V | k_3, k_4 \rangle$  is rotation invariant. That is

$$\langle Rk_1, Rk_2 | V | Rk_3, Rk_4 \rangle = \langle k_1, k_2 | V | k_3, k_4 \rangle$$

for any element  $R$  of  $SO(3)$  acting on spatial components. The chemical potential  $\mu$  in  $e(\mathbf{k})$  determines the electron density of the model.

The infrared behaviour of this model is determined (see [FT]) by a running coupling “constant”  $F^{(h)}(t', s')$ ,  $h \leq 0$  where at scale  $h$  the momentum  $k$  is restricted to a shell  $M^h$  away from the Fermi surface  $e(\mathbf{k}) = 0$  and  $t' = (0, \frac{t}{|t|} k_F)$  projects  $t$  onto the Fermi surface. Initially

$$F^{(0)}(t', s') = -\lambda \langle t', -t' | V | s', -s' \rangle.$$

The kernel  $F^{(h)}(t', s')$  defines an operator on  $L^2(k_F S^2)$ .

By rotation invariance the operator  $F^{(h)}$  commutes with the action of  $SO(3)$ . Therefore the eigenspaces of  $F^{(h)}$  coincide with the  $SO(3)$  irreducible invariant subspaces

of  $L^2(k_F S^2)$ . Recall that the space  $H^n$ , obtained by restricting homogeneous harmonic polynomials of degree  $n$  to  $S^2$ , is a  $2n + 1$  dimensional  $SO(3)$  irreducible invariant subspace of  $L^2(k_F S^2)$  and that

$$L^2(k_F S^2) = \bigoplus_{n \geq 0} H^n.$$

It follows that

$$F^{(h)}(t', s') = \sum_{n \geq 0} \lambda_n^{(h)} \pi_n(t', s')$$

where  $\pi_n$  is the orthogonal projection onto  $H^n$  and  $\lambda_n, n \geq 0$  is the spectrum of  $F^{(h)}$ . Here,  $\pi_n(t', s') = (2n + 1)k_F^{-2-n} P_n(\langle t', s' \rangle)$  where  $P_n$  is the Legendre polynomial of degree  $n$ .

It is widely believed that any (sufficiently weak) interaction  $\langle k_1, k_2 | V | k_3, k_4 \rangle$  flows, after, say,  $h$  steps, to an effective interaction  $F^{(h)}$  that is dominated by a single attractive angular momentum sector  $\lambda_\ell^{(h)} > 0$  (see [KL]). The infrared behaviour is then likely to be determined by the corresponding BCS model with gap equation

$$\Delta(\mathbf{p}) = \frac{1}{2} \int_{|\epsilon(\mathbf{q})| \leq \epsilon} \frac{d^3 \mathbf{q}}{(2\pi)^3} \lambda_\ell^{(h)} \pi_n(\mathbf{p}', \mathbf{q}') \Delta(\mathbf{q}) \frac{1}{E(\mathbf{q})} \tanh\left(\frac{1}{2}\beta E(\mathbf{q})\right). \quad (1)$$

Here,

$$\Delta(\mathbf{p}) = (\Delta_{\sigma, \sigma'}(\mathbf{p}))_{\sigma, \sigma' \in \{\uparrow, \downarrow\}}$$

is a  $2 \times 2$  matrix satisfying

$$\Delta(\mathbf{p}) = -\Delta(-\mathbf{p})^T$$

and

$$E(\mathbf{q})^2 = e(\mathbf{q})^2 + \Delta(\mathbf{q})^* \Delta(\mathbf{q}).$$

The expression  $\frac{1}{E(\mathbf{q})} \tanh\left(\frac{1}{2}\beta E(\mathbf{q})\right)$  is unambiguously defined by expanding  $\frac{1}{\sqrt{x}} \tanh\left(\frac{1}{2}\beta\sqrt{x}\right)$  as a power series in  $x$ . For a derivation of (1) see [AB], [BW].

Every solution of (1) is of the form

$$\Delta(\mathbf{p}) = (Y_{\sigma, \sigma'}(\mathbf{p})), \quad Y_{\sigma, \sigma'} \in H_\ell.$$

The simplest solutions are unitary and isotropic. A solution is unitary when

$$\Delta(\mathbf{p})^* \Delta(\mathbf{p}) = |d(\mathbf{p})|^2 I$$

and isotropic when  $d(\mathbf{p})$  is a constant. In this case the quasiparticle dispersion relation  $(e(\mathbf{q})^2 + |d|^2)^{\frac{1}{2}}$  is isotropic and has a gap  $|d|$  determined by

$$1 = \frac{1}{2} \int_{|e(\mathbf{q})| \leq \epsilon} \frac{d^3 \mathbf{q}}{(2\pi)^3} \lambda_\ell^{(h)} (e(\mathbf{q})^2 + |d|^2)^{-\frac{1}{2}} \tanh \left[ \frac{1}{2} \beta (e(\mathbf{q})^2 + |d|^2)^{\frac{1}{2}} \right] \quad (2)$$

when  $d \neq 0$ . Intuitively, they have the best chance of being stable.

There are two important examples of isotropic, unitary solutions. For  $\ell = 0$  there is the BCS model

$$\Delta = \begin{bmatrix} 0 & d \\ -d & 0 \end{bmatrix}$$

for phononic superconductivity. Balian and Werthamer discovered, in the  $\ell = 1$  sector, the solution

$$\Delta = d \begin{bmatrix} -\mathbf{p}_1 + i\mathbf{p}_2 & \mathbf{p}_3 \\ \mathbf{p}_3 & \mathbf{p}_1 + i\mathbf{p}_2 \end{bmatrix}, \quad \mathbf{p}_1^2 + \mathbf{p}_2^2 + \mathbf{p}_3^2 = k_F^2$$

which describes the B phase of He<sup>3</sup>.

Theorem *There are no nontrivial, isotropic, unitary solutions of (1) for  $\ell \geq 2$ .*

One therefore expects that solutions will have nodes for  $\ell \geq 2$  making the flow harder to control. Such nodes are observed in the A phase of He<sup>3</sup> and in the  $\ell = 2$  theory of heavy fermionic superconductivity. Nodes also appear in the gap function for systems with cubic symmetry. See, for example, [VG].

The proof of Theorem 1 follows immediately from the

Lemma *Let  $f, g \in H_\ell$  satisfy  $f\bar{f} + g\bar{g} = 1$  on  $S^2$ . Then,  $\ell = 0, 1$ .*

Proof Let  $P_\ell$ ,  $\ell \geq 0$ , be the homogeneous polynomials of degree  $\ell$  on  $\mathbf{R}^3$  with SO(3) invariant inner product

$$\langle f, g \rangle := f \left( \frac{\partial}{\partial k_1}, \frac{\partial}{\partial k_2}, \frac{\partial}{\partial k_3} \right) \bar{g}.$$

As usual  $H_\ell$  is identified with  $H_\ell^*$  by the SO(3) equivariant isomorphism

$$f \mapsto \langle \cdot, \bar{f} \rangle.$$

We shall show that under the hypothesis of the lemma

$$U = f \otimes \bar{f} + \bar{f} \otimes f + g \otimes \bar{g} + \bar{g} \otimes g$$

is the (unique up to scalars)  $\text{SO}(3)$  invariant element of  $H_\ell \otimes H_\ell$ . It follows that the homomorphism

$$U \in H_\ell \otimes H_\ell \cong H_\ell \otimes H_\ell^* \cong \text{Hom}(H_\ell, H_\ell)$$

commutes with  $\text{SO}(3)$  and is of rank at most four. Moreover, by Schur's Lemma,  $U$  is an isomorphism since  $H_\ell$  is irreducible. Consequently,  $2\ell + 1 \leq 4$ .

Consider the  $\text{SO}(3)$  equivariant multiplication map

$$H_\ell \otimes_s H_\ell \xrightarrow{M} P_{2\ell}$$

$$\sum c_j \phi_j \otimes \psi_j \longmapsto \sum_j c_j \phi_j \psi_j.$$

Observe that

$$\dim H_\ell \otimes_s H_\ell = 2\ell + 1 + \frac{(2\ell + 1)(2\ell)}{2} = \binom{2\ell + 2}{2} = \dim P_{2\ell}$$

and

$$MU = 2|k|^{2\ell}.$$

If  $M$  is surjective it is an isomorphism and  $U$  is invariant.

The projection of

$$M \left( (k_1 + ik_2)^\ell \otimes_s (k_1 - ik_2)^\ell \right) = (k_1^2 + k_2^2)^\ell$$

onto the irreducible subspace  $|k|^{2(\ell-m)} H_{2m}$  of  $P_{2\ell}$  is nonzero because

$$\begin{aligned} & \left\langle |k|^{2(\ell-m)} (k_1 + ik_3)^{2m}, (k_1^2 + k_2^2)^\ell \right\rangle \\ &= \left( \frac{\partial}{\partial k_1} + i \frac{\partial}{\partial k_3} \right)^{2m} \Delta^{\ell-m} (k_1^2 + k_2^2)^\ell \\ &= \prod_{j=0}^{\ell-m-1} 4(\ell-j)^2 \left( \frac{\partial}{\partial k_1} + i \frac{\partial}{\partial k_3} \right)^{2m} (k_1^2 + k_2^2)^m \\ &\neq 0. \end{aligned}$$

Recall that every invariant subspace of  $P_{2\ell}$  is of the form

$$\bigoplus_{j_i} |k|^{2j_i} H_{2(\ell-j_i)}$$

with  $0 \leq j_1 < j_2 \dots < j_r \leq \ell$  and in particular

$$P_{2\ell} = \bigoplus_{j=0}^{\ell} |k|^{2j} H_{2(\ell-j)}.$$

Finally the image of  $M$  is invariant and therefore all of  $P_{2\ell}$ . ■

We observe that in two dimensions there are unitary isotropic solutions of the gap equation for every angular momentum. For example,

$$\Delta(\mathbf{p}) = d \begin{bmatrix} \cos \ell\theta & \sin \ell\theta \\ \sin \ell\theta & -\cos \ell\theta \end{bmatrix}$$

when  $\ell$  is odd and

$$\Delta(\mathbf{p}) = d \begin{bmatrix} 0 & e^{i\ell\theta} \\ -e^{i\ell\theta} & 0 \end{bmatrix}$$

when  $\ell$  is even. Here,  $\mathbf{p} = |\mathbf{p}|(\cos \theta, \sin \theta)$ .

#### References

- [AB] P.W. Anderson and W.F. Brinkman, Theory of Anisotropic Superfluidity in "Basic Notions of Condensed Matter Physics" by P.W. Anderson, Benjamin/Cummings, Menlo Park (1984).
- [BW] R. Balian and N.R. Werthamer, Superconductivity with Pairs in a Relative p Wave, Phys. Rev. 131, 1553-1564 (1963).
- [FT] J. Feldman and E. Trubowitz, The Flow of an Electron-Phonon System to the Superconducting State, to appear in Helvetica Physica Acta.
- [KL] W. Kohn and J.M. Luttinger, New Mechanism for Superconductivity, Phys. Rev. Lett. 15, 524-526 (1965).
- [VG] G.E. Volovik and L.P. Gor'kov, An Unusual Superconductivity in UBe<sub>13</sub>, JETP Lett. 39, 674-677 (1984).