

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Multifractality and scaling in disordered 2-D Landau levels
Autor: Huckestein, B. / Schweitzer, L.
DOI: <https://doi.org/10.5169/seals-116415>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Multifractality and Scaling in Disordered 2-D Landau Levels

B. Huckestein and L. Schweitzer

Physikalisch-Technische Bundesanstalt,
Bundesallee 100, W-3300 Braunschweig, F. R. Germany

Abstract. The structure of the eigenstates of disordered 2-D systems of size $M \times M$ in strong perpendicular magnetic fields is investigated. A multifractal behaviour is found in the centres of the Landau bands, which is described in terms of the $f(\alpha(q))$ vs. $\alpha(q)$ spectrum. The eigenstates $\psi_{E,M}(\mathbf{r})$ and the equilibrium current density $\mathbf{j}_{E,M}(\mathbf{r})$ show similar $f(\alpha(q))$ spectra which supports the view of a universal one-parameter scaling behaviour.

Introduction

Electronic states in disordered two-dimensional systems, subject to a strong perpendicular magnetic field, exhibit complex self similar structures on length scales l , which are larger than both the magnetic length and the correlation length of the potential, but are smaller than the system size M [1, 2, 3]. These states appear at energies near the centres E_n of the Landau bands, where the localization length $\xi(E)$ diverges as $|E - E_n|^{-\nu}$, with $\nu = 2.35$ [4, 5, 6]. To characterize the multifractal states, the $f(\alpha(q))$ vs. $\alpha(q)$ spectrum has been calculated [3, 6], which can be fitted to a parabola, $f(\alpha) = D_0 - (\alpha - \alpha_0)^2/(4(\alpha_0 - D_0))$. This reflects the log-normal distribution of the box-probabilities $p_i(l, E, M) = \sum_{\mathbf{r} \in \Omega_i(l)} |\psi_{E,M}(\mathbf{r})|^2$. With $D_0 = 2$, we find for the lattice model $\alpha_0 = 2.29 \pm 0.02$, which is in accordance with the result for the continuum model [3].

In this contribution a multifractal analysis of the equilibrium current density $\mathbf{j}_{E,M}(\mathbf{r})$ is presented. This quantity shows a similar behaviour as found for the eigenstates. Thus, a universal $f(\alpha(q))$ -spectrum for a variety of physical observables seems likely.

Model and Method

A one band tight binding model on a simple cubic lattice with periodic boundary conditions is considered to describe non-interacting electrons in disordered 2-D systems in strong perpendicular magnetic fields. The site energies are independent random numbers distributed with probability $1/W$ over the interval $[-W/2, W/2]$ and the magnetic field B enters the phase factors of the transfer matrix elements. A Lanczos algorithm is applied to calculate the eigenvalues and eigenvectors of the corresponding sparse hermitian matrices.

The $f(\alpha)$ -spectrum is calculated according to [7] from

$$\alpha(q) = \lim_{\lambda \rightarrow 0} \sum_i \mu_i(q, \lambda) \ln P_i(\lambda) / \ln(\lambda) \quad (1)$$

and

$$f(q) = \lim_{\lambda \rightarrow 0} \sum_i \mu_i(q, \lambda) \ln \mu_i(q, \lambda) / \ln(\lambda), \quad (2)$$

with $\mu_i(q, \lambda) = P_i^q(\lambda) / \sum_j P_j^q(\lambda)$ and $\lambda = l/M$.

For the investigation of the equilibrium current density of an eigenstate with energy E , $\mathbf{j}_{E,M}(\mathbf{r}) = \sum_{\mathbf{r}'} \langle E | \mathbf{r}' \rangle \langle \mathbf{r}' | \mathbf{j} | \mathbf{r} \rangle \langle \mathbf{r} | E \rangle$, a normalized box-probability was defined as $P_i(\lambda) = |\mathbf{j}_i(\lambda)| / \sum_k |\mathbf{j}_k(\lambda)|$ [3], because the current density is, in contrast to the eigenstates, not normalized, and $\mathbf{j}_i(\lambda) = \sum_{\mathbf{r} \in \Omega_i(\lambda)} \mathbf{j}_{E,M}(\mathbf{r})$.




Fig. 1: a) The density $|\psi_{E,M}(\mathbf{r})|^2$ of the most extended state in the centre of the lowest Landau band. b) $f(\alpha)$ -spectrum of the current density of the state shown on the left. Crosses give the errors in $f(q)$ and $\alpha(q)$ for $q = 2, 1, -1, -2$ (from left to right).

Results and Discussion

In fig. 1a the modulus of the most extended state, $|\psi_{E,M}(\mathbf{r})|^2$, of the lowest Landau band for a system of size $M = 150a$, is shown as a function of the position. The magnetic field strength is equivalent to $1/5$ flux-quanta h/e per unit lattice cell a^2 . A strongly peaked structure of the wavefunction is observed. However, this must not be regarded as a sign of localization, but reflects the fact that the logarithm of the density $|\psi_{E,M}(\mathbf{r})|^2$ is normally distributed [3]. The $f(\alpha(q))$ -spectrum of the corresponding equilibrium current density $\mathbf{j}_{E,M}(\mathbf{r})$ is plotted in fig. 1b together with a parabolic fit from which a value of $\alpha_0 = 2.31 \pm 0.03$ is obtained. Therefore, the $f(\alpha(q))$ -spectrum of the current density is, within the current uncertainty, the same as for the eigenstates [6].

References

- [1] H. Aoki, Phys. Rev. B **33**, 7310 (1986).
- [2] B. Huckestein and L. Schweitzer, in *High Magnetic Fields in Semiconductor Physics III*, edited by G. Landwehr (Springer Series in Solid State Sciences, (in print), 1991).
- [3] W. Pook and M. Janßen, Z. Phys. B **82**, 295 (1991).
- [4] G. V. Mil'nikov and I. M. Sokolov, JETP Lett. **48**, 536 (1988).
- [5] B. Huckestein and B. Kramer, Phys. Rev. Lett. **64**, 1437 (1990).
- [6] B. Huckestein, B. Kramer, and L. Schweitzer , in *Proceedings of the Ninth International Conference on the Electronic Properties of Two-Dimensional Systems*, (Surface Sciences, (in print), 1991).
- [7] A. Chhabra and R. V. Jensen, Phys. Rev. Lett. **62**, 1327 (1989).