

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Nonperturbatively renormalizable quantum field theories in 2+1 dimensions
Autor: Kovner, A. / Rosenstein, B. / Gat, G.
DOI: <https://doi.org/10.5169/seals-116459>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Nonperturbatively Renormalizable Quantum Field Theories in 2+1 Dimensions

A.Kovner, B. Rosenstein and G. Gat

Physics Department
University of British Columbia
Vancouver, B.C.
Canada, V6T 2A6

Abstract. Recently several 2+1 dimensional models were found which are non-renormalizable in weak coupling expansion but renormalizable in $1/N$. We show that the IR fixed points of Yukawa theories and scalar QED_3 (Landau-Ginzburg) are some of these theories. The critical 4-Fermi interaction model is the IR stable fixed point of Yukawa theories while the CP^{N-1} model is the IR fixed point of N-component scalar QED_3 .

As is well known, there is a close relation between renormalizable relativistic quantum field theories (QFT) in 2+1 dimensions and second order phase transitions in 3D. For (rotationally invariant) statistical systems universal critical properties are determined by an infrared (IR) fixed point of the corresponding QFT. For example, critical exponents of Heisenberg (anti)ferromagnets are described by the $O(3)$ symmetric ϕ_3^4 [1] theory

$$S_E[\phi] = \frac{1}{2}(\partial\phi_i)^2 - \frac{\mu^2}{2}\phi_i^2 + \frac{\lambda}{4!}\phi_i^4. \quad (1)$$

The IR fixed point of ϕ^4 is the conformally invariant critical σ -model. Note that although the nonlinear σ -model is non renormalizable in weak coupling perturbation theory it is renormalizable nonperturbatively.

Recently several new classes of nonperturbatively renormalizable QFT's in 2+1 dimensions were found using the $1/N$ expansion. These include four-Fermi interaction models like the Z_2 chiral invariant Gross-Neveu (GN) model [2]

$$\mathcal{L} = \bar{\psi}_i \not{\partial} \psi_i + \frac{g^2}{2N} (\bar{\psi}_i \psi_i)^2. \quad (2)$$

We show that the critical GN model is the IR fixed point of the superrenormalizable Yukawa theories

$$\mathcal{L} = \frac{1}{2}(\partial\sigma)^2 + \bar{\psi}_i \not{\partial} \psi_i + \lambda_0 \sigma \bar{\psi}_i \psi_i + \frac{m_0^2}{2} \sigma^2 \quad (3)$$

and therefore determines its universality class. At the phase transition point the chiral Z_2 breaks down spontaneously. The renormalized coupling of the Yukawa model is determined by (we use notations of [1])

$$\sqrt{m}\lambda = \Gamma_{1,2}(p_1 = 0, p_2 = 0, p_3 = 0) \quad (4)$$

where m is the boson's mass, $\Gamma_b^2(m) = 0$. To fix the field's normalizations we choose to impose $\frac{\partial \Gamma_b^2}{\partial p^2} \Big|_{p^2=m^2} = 1$, $\frac{\partial \Gamma_b^2}{\partial p^2} \Big|_{p^2=0} = 1$. To leading order in $1/N$ the corresponding Callan-Symanzick β function is

$$\beta(\lambda) = -\frac{1}{2}\lambda + \frac{1}{16}\lambda^3 \quad (5)$$

It has a finite IR fixed point at $\lambda^* = 2\sqrt{2}$. The connection between the bare and renormalized couplings is determined by the RG equation

$$\left(1 + \beta(\lambda)\frac{\partial}{\partial\lambda}\right) \frac{\lambda_0^2}{m} = 0 \quad (6)$$

The solution is

$$\lambda_0(\lambda) = \sqrt{m}\lambda \exp\left\{-\frac{1}{2}\int_0^\lambda \left[\frac{1}{\beta(x)} + \frac{1}{x}\right] dx\right\} \quad (7)$$

As $\lambda \rightarrow \lambda^*$, the bare coupling λ_0 diverges $\lambda_0 \rightarrow \infty$. In this limit the Yukawa action becomes the GN action eq.(2).

This is seen explicitly by calculating the dynamical dimension of the kinetic term for σ : $[(\partial\sigma)^2] = 4$ [3]. It becomes irrelevant for $d = 3$. On the other hand some irrelevant operators at weak coupling become relevant or marginal near the fixed point. The dimension four $(\bar{\psi}\psi)^2$ operator acquires scaling dimension two and thus becomes relevant , while the dimension six operator $(\bar{\psi}\psi)^3$ has in the leading order in $1/N$ scaling dimension three and thus becomes marginal. In this case more careful analysis is needed. The next to leading order in $1/N$ shows that in fact it becomes relevant. Similar phenomenon occurs in abelian gauge theories . In scalar QED the IR fixed point is the nonperturbatively renormalizable CP^{N-1} model [4]. It is interesting to note that the vortex operator V that creates Nielsen-Olesen vortex [5] becomes relevant at $N < 8$.

To summarize we found that recently discovered nonperturbatively renormalizable 2+1 dimensional QFT contain IR fixed points describing new universality class. It is interesting to know whether these universality classes are realized in real materials.

The work of A.K and B.R was supported by NSERC of Canada.

References

- [1] C. Itzykson and J.M. Drouffe, *Statistical Field Theory*, (Cambridge University Press, 1991) vol. I
- [2] B. Rosenstein, B.J. Warr and S.H. Park *Phys.Rev.Lett.* **62**, 1433 (1989) ; B. Rosenstein, B.J. Warr and S.H. Park *Phys.Rep.* **205**, 59 (1991) and references therein ; C. de Calan et al, *Phys.Rev.Lett* **66**, 3233 (1991)
- [3] Y. Kikukawa and K. Yamawaki, *Phys.Lett B* **234**, 497 (1990) ; G. Gat et al, *Phys.Lett. B* **240**, 158 (1990)
- [4] A. Kovner and B. Rosenstein *Phys.Lett B* **261**, 97 (1991)
- [5] A. Kovner and B. Rosenstein *Phys.Lett B* **261**, 104 (1991)