

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Slave Boson approach to magnetic order in the Hubbard model
Autor: Frésard, Raymond / Wölfle, Peter
DOI: <https://doi.org/10.5169/seals-116465>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Slave Boson Approach to Magnetic Order in the Hubbard Model

Raymond Fr  sard and Peter W  lfle

**Institut für Theorie der Kondensierten Materie, Universität Karlsruhe,
D-W-7500 Karlsruhe, Germany**

Abstract. Using a saddlepoint approximation of a spin-rotation-invariant form of the slave boson representation introduced by Kotliar and Ruckenstein we determine for the Hubbard model on the square lattice the phase diagram when spiral magnetic states are taken into consideration. We display the spiral wave-vector and discuss its temperature dependence.

Introduction

It is fairly well established that there is antiferromagnetic long range order in the ground state of the single band Hubbard model on a two dimensional square lattice at half-filling [1,2]. Relatively little is known about the magnetic state of the model away from half-filling. In this work we report numerical results of a slave boson theory supporting the view that spiral states are good candidates for the ground state of the Hubbard model near half-filling and at intermediate to large interaction strength U .

The method

Our analytical treatment of the Hubbard model is based on the spin-rotation-invariant form [3] of the slave boson representation introduced by Kotliar and Ruckenstein [4]. We consider a saddle-point approximation corresponding to the site-dependent magnetization $\vec{M}_i = M \hat{n}_i$, where $\hat{n}_i = (\cos \vec{Q} \cdot \vec{R}_i, \sin \vec{Q} \cdot \vec{R}_i, 0)$ forms a spiral structure. The free energy takes the form:

$$F = -T \sum_{\vec{k},\sigma} \ln(1 + e^{-\frac{E_{\vec{k},\sigma}}{T}}) + U d^2 - 2\beta_0(p_o^2 + p^2 + d^2) - 4p_0 p \beta + \alpha(1 - e^2 - 2p_0^2 - 2p^2 - d^2)$$

with the quasiparticle energies:

$$E_{\vec{k},\sigma} = (z_0^2 + z^2) \left(\frac{t_k + t_{k-Q}}{2} \right) + \beta_0 - \mu_0 \pm \left[\left((z_0^2 - z^2) \left(\frac{t_k - t_{k-Q}}{2} \right) \right)^2 + (z_0 z(t_k + t_{k-Q}) + \beta)^2 \right]^{\frac{1}{2}}$$

where e^2 , $p_0^2 + \sigma p^2$, d^2 represent respectively the number of empty, spin σ singly occupied, and doubly occupied sites, and α , β_0 and β the Lagrange multipliers enforcing the constraints; they are determined by minimizing F . For details about this representation we refer to [5,6].

Results

We have solved numerically the saddlepoint equations on a 98x98 lattice. The results are summarized in the (U, δ) phase diagram, fig. 1. At $T = 0$ we see that the Antiferromagnetic (AF) state takes place at half-filling only, for any value of U . Upon doping the system immediately enters a spiral phase characterised by $\vec{Q} = (Q_x, Q_z)$ (see below). In this part of the phase diagram the Fermi surface consists of one "hole pocket" centered at $(\pi/2, \pi/2)$. Moreover we obtain that the magnetic gap 2β is severely reduced from $\sim U$ at half-filling (for large U) down to $\sim t$ which implies that the band width is getting close to $8t$, contrary to Hartree-Fock results, but in agreement with exact diagonalisation results, for instance for two holes in the $t-J$ model [7]. When the doping is increased the system undergoes a first-order phase transition towards another spiral phase characterised by $\vec{Q} = (Q_x, \pi)$. In this phase the magnetization is decreasing linearly with the doping up to a second-order phase transition to the Paramagnetic state. Fig. 2 is displaying Q_z in slave boson (SB) and Hartree-Fock (H-F) approaches as func-

tions of the density for $U = 8t$ (the jump indicates the transition from (Q_x, Q_z) to (Q_x, π)). The deviation from π for small doping is given by $\delta U/t$, its large U behaviour for both SB and H-F. In SB Q_x then saturates to ~ 2.2 , thus showing no ferromagnetism, in agreement with [4].




Fig. 1




Fig. 2

We also investigated the temperature dependence of \vec{Q} and obtained as ground state the AF state over a finite range of doping given by $\sim 2T/U$.

Conclusion

Within the spin-rotation invariant slave boson mean-field approach [3,4] we showed a phase diagram up to rather large U which exhibits AF, P and 2 different spiral states which extends over a large density domain. We obtain that the temperature is stabilizing the AF state with respect to the spiral one.

We thank M. Dzierzawa for providing his Hartree-Fock data and for many stimulating discussions. One of us (RF) thanks the Swiss National Science Foundation for financial support. This project has been supported by ESPRIT project MESH-3041.

References

- [1] Liang S., Doucot B. and Anderson P. W., Phys. Rev. Lett. **61** (1988) 365 ;
- [2] Trivedi N. and Ceperley D., Phys. Rev. **B40** (1989) 2737.
- [3] Li T., Wölfle P. and Hirschfeld P., Phys. Rev. **B40** (1990) 6817.
- [4] Kotliar G. and Ruckenstein A., Phys. Rev. Lett. **57** (1986) 1362.
- [5] Wölfle P. and Li T., Z. Phys. B - Condensed Matter **78** (1990) 45.
- [6] Frésard R., Dzierzawa M. and Wölfle P., Europhys. Lett. **15**, 325-330 (1991).
- [7] Stephan, W. and Horsch P., Phys. Rev. Lett. **66** (1991) 2258.