

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Finite-size effects in the negative U Hubbard model
Autor: Bormann, D. / Schneider, T. / Frick, M.
DOI: <https://doi.org/10.5169/seals-116471>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Finite-Size Effects in the Negative U Hubbard Model

D. Bormann^{a,b}, T. Schneider^a and M. Frick^{a,c} ¹

^a IBM Research Division, Zurich Research Laboratory, CH-8803 Rüschlikon

^b Institute for Theoretical Physics, University of Heidelberg, D-6900 Heidelberg

^c Institute for Theoretical Physics, University of Groningen, NL-9700 AV Groningen

Abstract. Using the BCS approach and Quantum Monte Carlo (QMC) simulations, we investigate the size dependence of the pairing correlations in the weak and intermediate coupling regimes. Pronounced finite size effects appear as long as the average spacing of the single-particle levels is larger than or comparable to the characteristic energy scale (superconducting gap). The irregularity of the size dependencies are traced back to variations in the occupation of the highest non-empty energy levels with the system size. We propose means to reduce the size dependence and its irregularity.

Introduction

The occurrence of superconductivity in models for strongly correlated fermions is a topical issue. Numerical methods have been developed to study these models without further approximations in small systems. Studying finite-size effects, we concentrate on the negative U Hubbard model,

$$H = -t \sum_{\langle i,j \rangle \alpha} (c_{i\alpha}^\dagger c_{j\alpha} + \text{h.c.}) - U \sum_i n_{i\uparrow} n_{i\downarrow} \quad (1)$$

as a generic system exhibiting superconductivity. Because the physical properties of a fermion system strongly depend on the density of states (DOS) at the Fermi level, the shell structure in the DOS on the finite lattice has drastic consequences [1,2]. These effects are significant as long as the average level spacing exceeds the energy scale of the superconducting gap.

Comparison between BCS and QMC

We performed Projector QMC simulations [3] on a two-dimensional lattice to obtain ground state properties and compared the (exact) results with a BCS treatment of (1) at fixed particle number (FBCS [4]). The corresponding component of the variational BCS wave function is projected out, before performing the variation of the parameters in the wave function. The variational equations are then solved numerically.

In Fig.(1), the results of the two methods for the Cooper pair vertex correlation functions

$$\chi_\nu = \frac{1}{N^2} \sum_{kl} f_k^\nu f_l^{\nu*} (\langle c_{k\uparrow}^\dagger c_{-k\downarrow}^\dagger c_{-l\downarrow} c_{l\uparrow} \rangle - \langle c_{k\uparrow}^\dagger c_{l\uparrow} \rangle \langle c_{-k\downarrow}^\dagger c_{-l\downarrow} \rangle). \quad (2)$$

are compared in the weak-coupling regime. The perfect agreement with the PQMC data shows that the FBCS wave function serves as a good variational wave function for the ground state on the finite lattice. The FBCS method improves the BCS results. Discrepancies at half-filling can be traced back to enhanced charge-density wave correlations. In the intermediate coupling regime, the FBCS scheme overestimates the superconducting correlations, which are reduced due to quantum fluctuations [5].

¹Part of the work has been financed by the stichting FOM, which is supported by the NWO.

Finite-size effects

Fig.(1) shows the pronounced effects of the DOS shell structure on χ_0 . Open shell configuration exhibit strongly enhanced superconducting correlations, which can be understood as a DOS effect. Moreover, we find that the symmetry of the Fermi surface is reflected in the symmetry of the Cooper pair. The fluctuations between open and closed shell configuration get reduced with increasing coupling strength and band filling. However, they complicate any extrapolation to the thermodynamic limit, especially for weak coupling. Finite-size effects are prominent as long as the average level spacing exceeds the energy scale of the superconducting gap. In the weak-coupling regime, this criterion yields as a lower bound on the lattice sizes, where these shell effects disappear,

$$N > \frac{gW}{U} \exp\left(\frac{W}{U}\right), \quad (3)$$

where $W = 4Dt$ denotes the band width on a D -dimensional lattice and g the average level degeneracy. However, the feasibility of an extrapolation is improved, if only closed shell configurations are used, as shown in fig.(2). The number of these configurations is enlarged by considering tilted clusters on the square lattice.

References

- [1] D.Bormann, T.Schneider and M.Frick, *Europhys.Lett.* 14, 101 (1991)
- [2] D.Bormann, T.Schneider and M.Frick, submitted to *Z.Phys.B* (1991)
- [3] S.E.Koonin, G.Sugiyama and H.Friederich in *Proc.Intl.Symp., Bad Honnef*, Springer, Heidelberg (1982)
- [4] P.Ring and P.Schuck, *The Nuclear Many-Body Problem*, Springer, Heidelberg (1980)
- [5] R.T.Scalettar, E.Y.Loh, J.E.Gubernatis, A.Moreo, S.R.White, D.J.Scalapino, R.L.Sugar and E.Dagotto, *Phys.Rev.Lett.* 62, 1407 (1989)




Figure 1. χ_0 ($f_k^0 = 1$) as a function of the band filling $\langle n \rangle$. 4×4 lattice, $U/t = 1$. Comparison of BCS ($\diamond$), FBCS ($\bullet$) and PQMC ($\blacksquare$) data.




Figure 2. $\sqrt{\chi_0}$ ($f_k^0 = 1$) as a function of the inverse linear lattice size $1/L$. $U/t = 1$, $\langle n \rangle = 0.75$. FBCS data. The infinite lattice BCS result is indicated.