

Zeitschrift:	Helvetica Physica Acta
Band:	65 (1992)
Heft:	2-3
Artikel:	Lyapunov exponents of the Schrödinger equation with quasi-periodic potential on a strip
Autor:	Goldsheid, I.Ya. / Sorets, E.
DOI:	https://doi.org/10.5169/seals-116475

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Lyapunov Exponents of the Schrödinger Equation with Quasi-periodic Potential on a Strip

I. Ya. Goldsheid,^{1)*} E. Sorets²⁾

¹⁾Fakultät für Mathematik
Ruhr-Universität-Bochum (FRG)
SFB-237 Bochum-Essen-Düsseldorf

²⁾Parametric Technology Corporation
128 Technology Drive
Waltham, MA 02138

Abstract. We describe conditions under which all the nonnegative Lyapunov exponents of the quasi-periodic difference Schrödinger equation on a strip are strictly positive.

We shall study the Lyapunov exponents of the difference equation:

$$-y_{n+1} + Q_n y_n - y_{n-1} = 0 \quad -\infty < n < +\infty \quad (1)$$

where $y_n \in \mathbb{R}^m$ and Q_n is a symmetric $m \times m$ matrix whose off-diagonal elements do not depend on n , and the diagonal elements are quasi-periodic functions

$$q_{ni}(\theta) = \lambda f_i(e^{2\pi i(\theta+n\alpha)}) - E$$

with $f_i(z)$ analytic on $\mathcal{A} \equiv \{z \mid r < |z| < 1/r\}$, taking values in $[-1, 1]$ for $|z| = 1$, λ is a (large) parameter called coupling constant, E is the energy, and α is any irrational number. Without loss of generality we shall assume that $\max_{1 \leq i \leq m} \sup_{|z|=1} f_i(z) = 1$ and $\min_{1 \leq i \leq m} \inf_{|z|=1} f_i(z) = -1$.

Equation (1) becomes equivalent to the finite-difference Schrödinger equation when the off-diagonal elements are chosen properly. For example, the case $q_{ij} = -1$ for $|i-j| = 1$ and $q_{ij} = 0$ for $|i-j| > 1$ corresponds to Schrödinger operator on the strip $\mathbb{Z} \times \{1, \dots, m\}$. Equation (1) can be written in the form

$$\begin{pmatrix} y_{n+1} \\ y_n \end{pmatrix} = \begin{pmatrix} Q_n & -I \\ I & 0 \end{pmatrix} \begin{pmatrix} y_n \\ y_{n-1} \end{pmatrix} \equiv A_n \begin{pmatrix} y_n \\ y_{n-1} \end{pmatrix} \quad (2)$$

and, thus, the asymptotic behavior of the solutions of equation (1) is determined by the asymptotic behavior of the product $S(n) := A_n \cdots A_1$.

Various problems in solid-state physics give rise to different classes of matrices Q_n . These classes are characterized by the level of randomness. The case of independent random Q_n was studied in [GM], where it was shown that under certain algebraic conditions on the support of the corresponding measure in the space of symmetric matrices all the

* On leave from Math. Institute, Academy of Sciences USSR, 450057 Ufa, USSR

Lyapunov exponents are different and, therefore, the smallest non-negative exponent is, in fact, positive. When Q_n 's are non-deterministic at least some of the Lyapunov exponents are strictly positive [KS,S,K]. On the other hand, if Q_n 's form a periodic sequence, it is easy to show that the "interesting" Lyapunov exponents vanish.

The case of quasi-periodic potentials exhibits mixed behavior. If the coupling constant λ is small and α is poorly approximated by rationals, it is known [BLT] that at least on part of the spectrum the Lyapunov exponent is zero when $m = 1$. When λ is large and $m = 1$ the Lyapunov exponent is positive [Si,CS,FSW,SS].

Here we study the case of $m > 1$ and large λ . We prove [GS] that all non-negative Lyapunov exponents are positive. The method we use is an extension of the one used in [SS], which, in turn, grew out of M. Herman's proof of positivity of the Lyapunov exponent in the case of $m = 1$, $f(z) = z + 1/z$, and $\lambda > 1$ [H].

To describe our result let us consider the following decomposition of $S(n)$:

$$S(n) = U(n)D(n)V(n)$$

where $U, V \in O(n)$, and $D(n) = \text{diag}(d_1^{(n)}, \dots, d_{2m}^{(n)})$ with $d_1 \geq d_2 \geq \dots \geq d_{2m} \geq 0$. Since $A_n \in Sp(m, \mathbb{R})$ for all n , we have $d_k = d_{2m-k+1}^{-1}$. The k -th Lyapunov exponent γ_k is defined by

$$\gamma_k := \lim_{n \rightarrow \infty} \frac{1}{n} \log d_k(n). \quad (3)$$

Clearly, $\gamma_k = -\gamma_{2m-k+1}$, and $\gamma_1 \geq \gamma_2 \geq \dots \geq \gamma_m \geq 0$. Existence of the limit in (3) for almost every θ and its independence of θ are guaranteed by the Subadditive Ergodic Theorem [Ki] and ergodicity of the underlying dynamical system $\theta \mapsto \theta + \alpha$.

Our main result is the following

Theorem. *There exists λ_0 such that for all $\lambda > \lambda_0$ and all E , there exists a set $\Omega(E) \subset [0, 1]$ of Lebesgue measure 1, such that*

$$\gamma_m(E) = \gamma_m(E, \theta) > 0 \quad \forall \theta \in \Omega(E).$$

This, together with Oseledec's Multiplicative Ergodic Theorem [O,GM] implies existence of m solutions of equation (1), which decay exponentially as $n \rightarrow +\infty$ and grow exponentially as $n \rightarrow -\infty$, and m other solutions of equation (1), which decay exponentially as $n \rightarrow -\infty$ and grow exponentially as $n \rightarrow +\infty$. These $2m$ solutions are linearly independent and form a basis of all solutions of equation (1). There are two important consequences of this fact.

- 1) The Green's function of H decays exponentially for almost every energy E .
- 2) The spectrum of the operator H defined by the left hand side of equation (1) is singular.

Remarks. We should point out that the spectrum of H can actually be purely singular continuous if α is a Liouville number [CFKS].

The requirement that λ is large cannot be avoided, for when λ is small KAM theory is applicable and there is absolutely continuous spectrum [BLT].

We strongly use the non-triviality of f_i 's in our proof. This condition is necessary, for there are examples in which the presence of constant f_i 's leads to appearance of zero Lyapunov exponents.

References

- [BLT] Bellisard J., Lima R., Testard D.: *A Metal-Insulator Transition for the Almost Mathieu Model*, Comm. Math. Phys., **88**, 207–234 (1983)
- [CFKS] Cycon H., Froese, R., Kirsch W., Simon B.: *Schrödinger operators, with application to quantum mechanics and global geometry*. Berlin, Heidelberg, New York: Springer 1987
- [CS] Chulaevsky V., Sinai Ya.: *Anderson localization for multi-frequency quasi-periodic potentials in one dimension*, Comm. Math. Phys., **125**, 91–112 (1989)
- [FSW] Fröhlich J., Spencer T., Wittwer P.: *Localization for a class of one-dimensional quasi-periodic potentials*, Comm. Math. Phys., **132**, 5–25 (1990)
- [GS] Goldsheid I.Ya., Sorets E.: *Lyapunov exponents of the Schrödinger equation with quasi-periodic potential on a strip*, to appear in Comm. Math. Phys. (1991)
- [GM] Goldsheid I.Ya., Margulis G.A.: *Lyapunov indices of products of random matrices*, Uspekhi Mat. Nauk **44**:5, 13–60 (1989). Translation in Russian Math. Surveys **44**:5, 11–71 (1989)
- [H] Herman M.: *Une méthode pour minorer les exposants de Lyapunov et quelques exemples montrant le caractère local d'un Théorème d'Arnold et de Moser sur le tore en dimension 2*, Commun. Math. Helv., **58**, 453–502 (1983)
- [Ki] Kingman J.F.C.: Subadditive processes. Springer Lecture Notes in Math. **539** 1976
- [K] Kotani S.: *Ljapunov indices determine absolutely continuous spectrum of stationary random one-dimensional Schrödinger operators*, Proc. of Taniguchi Symp. SA. Kata-ta, 225–247 (1982)
- [KS] Kotani S., Simon, B.: *Stochastic Schrödinger Operators and Jacobi matrices on the strip*, Comm. Math. Phys. **119**, 403–429 (1988).
- [O] Oseledeć V.I.: *A multiplicative ergodic theorem, Characteristic Lyapunov exponents of dynamical systems*, Trudy Moskov. Mat. Obshch. **19**, 179–210 (1968) translation in Trans. Moscow Math. soc. **19**, 197–231 (1968)
- [S] Simon B.: *Kotani theory for one-dimensional stochastic Jacobi matrices*, Commun. Math. Phys. **89**, 227–234 (1983).
- [Si] Sinai Ya.: *Anderson localization for one-dimensional difference Schrödinger operator with quasi-periodic potential*, J. Stat. Phys., **46**, 861–918 (1987)
- [SS] Sorets E., Spencer T. : *Positive Lyapunov exponents for Schrödinger operator with quasi-periodic potentials*, to appear in Comm. Math. Phys. (1991).