

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Singular dynamics of a fermion coupled to the planar antiferromagnet
Autor: Ramšak, Anton / Prelovšek, Peter
DOI: <https://doi.org/10.5169/seals-116476>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Singular Dynamics of a Fermion Coupled to the Planar Antiferromagnet

Anton Ramšak and Peter Prelovšek
 J. Stefan Institute, University of Ljubljana,
 61111 Ljubljana, Slovenia/Yugoslavia

Abstract. A model for a single fermion coupled to the planar antiferromagnetic spin background via the local spin - fermion coupling is studied. Within the perturbation theory as well as the selfconsistent calculation of the quasiparticle spectra the fermion self energy shows a nonanalytical behaviour $\Sigma''(\omega > 0) \propto \omega^\alpha$, $\alpha \leq 1$. The original Hamiltonian is approximated with an effective single level model with a linear coupling to soft bosons. Exact results of the latter model again indicate on the orthogonality catastrophe in this system.

In order to understand the character of quasiparticles (QP) in strongly correlated systems, a number of authors have so far considered the problem of a single charge carrier introduced by doping the planar antiferromagnetic (AFM) insulator [1]. We consider in this paper the spin-hole (fermion) model which represents well the qualitative features of more complete two-band Hubbard model for the CuO₂ layers as well as the *t-J* model in superconducting oxides [2]. One of the most important qualitative questions is the correct description of the low frequency excitation spectra, connected with the existence of the QP peak [3]. This question is clearly related also to the possible breakdown of the Landau Fermi liquid description at arbitrary doping, as deduced from experiments [4].

We investigate the low energy features of a simplified version of the spin-hole model [2], where we retain the Heisenberg interaction among the localized spins (*J*), fermion hopping between two sublattices (neighbouring sites) A and B (*t*) and the local spin-fermion coupling (*V*),

$$H = -t \sum_s (c_{As}^\dagger c_{Bs} + c_{Bs}^\dagger c_{As}) + J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j + V (\mathbf{s}_A \cdot \mathbf{S}_A + \mathbf{s}_B \cdot \mathbf{S}_B).$$

We perform the expansion of spin operators \mathbf{S}_i into AFM linear spin waves and use a suitable transformation of fermion operators $c_{A(B),\pm\frac{1}{2}}$ into $\tilde{c}_{1(2),\pm}$ to get

$$H_{\text{mag}} = \epsilon \sum_{i=1}^2 (\tilde{n}_{i+} - \tilde{n}_{i-}) + \sqrt{2}J \sum_{q \sim 0} |\mathbf{q}| \alpha_q^\dagger \alpha_q + \frac{32^{-1/4}V}{\sqrt{N}} \sum_{\substack{i=1 \\ q \sim 0}}^2 \left(|\mathbf{q}|^{-1/2} \tilde{c}_{i+}^\dagger \tilde{c}_{i-} \alpha_q^\dagger + \text{H.c.} \right) + \sum_{q \sim (\pi, \pi)} \dots,$$

where $\epsilon = (t^2 + \frac{1}{16}V^2)^{1/2}$ and for the low energy excitations only the most important long-wavelength part is presented.

Fermion self-energy $\Sigma(\omega)$ corresponding to H_{mag} can be studied using the perturbation expansion. Within the lowest non-trivial order the result is similar as in Ref. 3, namely $\Sigma(\omega) \propto V^4 \epsilon^{-2} J^{-2} \omega \log(-\omega)$. Terms of higher order in *V* are also singular, $\propto \log^n(-\omega)$, $n = 1, 2, \dots$, and are therefore difficult to deal with. On the other hand contributions to $\Sigma(\omega)$ can be partly summed

up on a self-consistent (SC) way as in Ref. 3 yielding a non-analytical result $\Sigma(\omega) \propto \sqrt{-\omega}$ for $|\omega|/J \ll 1$.

These results can to some extent be improved by approximating the original (two level) Hamiltonian H_{mag} with a simpler one, which mimics two magnon processes among the lower energy ($-\epsilon$) states $\tilde{c}_{1(2),-}$ within H_{mag} . We thus introduce an effective linear interaction between the fermion (c) and magnons (β_q)

$$H_{\text{eff}} = J \sum_{q \sim 0} |\mathbf{q}| \beta_q^\dagger \beta_q + \frac{V^2}{\epsilon \sqrt{N}} \sum_{q \sim 0} c^\dagger c (\beta_q^\dagger + \beta_{-\mathbf{q}}).$$

Most divergent terms of $\Sigma(\omega)$ within H_{mag} show the same form of singularities as calculated using H_{eff} , e.g. the order $2n$ within H_{mag} corresponds to the order n within the perturbation expansion in V etc.

Solution for the fermion Green's function within H_{eff} can be obtained exactly using a canonical transformation [5]. The result is

$$G(\omega) = -i \int_0^\infty e^{\alpha h(t)} e^{i\omega t} dt, \quad h(t) = - \int_0^{q_m} \frac{1 - e^{-iJ_q t}}{q} dq,$$

and $2\pi\alpha = V^4(\epsilon J)^{-2}$. For small values $\omega/J > 0$ and $\alpha < 1$ it is easy to see that $\text{Im}G(\omega) \propto \omega^{\alpha-1}$ and $\Sigma''(\omega) \propto \omega^{1-\alpha}$, while for arbitrary values of the parameters the solution has to be obtained numerically (see Figure). To conclude we point out that at least within the effective model the QP peak does not exist. The nonanalytical self-energy shows the similarity with the orthogonality catastrophe [4].




Figure: Spectral function $A(\omega) = -\frac{1}{\pi} \text{Im}G(\omega)$ vs. ω/J for various values of α .

References

- [1] C. L. Kane, P. A. Lee and N. Read, Phys. Rev. B **39**, 6880 (1989) and references therein.
- [2] P. Prelovšek, Phys. Lett. A **126**, 287 (1988); A. Ramšak and P. Prelovšek, Phys. Rev. B **40**, 2239 (1989).
- [3] A. Ramšak and P. Prelovšek, Phys. Rev. B **42**, 10415 (1990).
- [4] C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams and A. E. Ruckenstein, Phys. Rev. Lett. **63**, 1996 (1989); P. W. Anderson, Phys. Rev. Lett. **64**, 1839 (1990).
- [5] I. G. Lang and Yu. A. Firsov, Zh. Eksp. Teor. Fiz. **43**, 1843 (1962).