

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Antiferromagnets in 2 + dimensions
Autor: Trimper, Steffen
DOI: <https://doi.org/10.5169/seals-116477>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ANTIFERROMAGNETS IN $2 + \varepsilon$ DIMENSIONS

Steffen Trimper

Fachbereich Physik der Martin-Luther-Universität, Halle
 D- 4020 Halle, Federal Republic of Germany

ABSTRACT. Using a perturbative approach with a modified Wick theorem for a spin 1/2 antiferromagnet we find the excitation spectra, an effective exchange interaction and the staggered magnetization for the square lattice at $T = 0$. In the framework of the non-linear sigma model we discuss the dimension shift, the influence of anisotropic and helical terms. Anisotropy leads always to a reduction of the quantum disordered fixed point. The results of large n expansion complete the conclusions of a renormalization group approach.

INTRODUCTION

The discovery of high temperature superconductors has led to renew efforts to understand quantum antiferromagnets /1/. The behaviour of an antiferromagnet depends on the value of the underlying spin. Since there is no general theory which covers the whole region between $s = 1/2$ and large spin we consider the two limiting cases in detail using a perturbative approach for the $s = 1/2$ system and the renormalization group approach for a continuum model which seems to be valid only for large spin /2/.

PERTURBATION THEORY FOR SPIN 1/2

The Heisenberg Hamiltonian can be expressed in terms of Pauli-operators

$$H = J(0) \sum_i b_i^+ b_i + J(0) \sum_j a_j^+ a_j + \sum_{i,j} J_{ij} (b_i a_j + a_j^+ b_i^+ - 2 b_i^+ b_i a_j^+ a_j) \quad (1)$$

The operators a and b act in different sublattices. They obey the commutation relation $[b_i, b_j^+] = \delta_{ij} (1 - 2 b_i^+ b_i)$ (2)

which prevent the application of the conventional Wick theorem. We have applied

a modified Wick theorem /3/ to calculate the temperature Green's function

$$G_{lm}(\tau_1 - \tau_m) = - \begin{pmatrix} \langle T_\tau (b_1 b_m^+) \rangle & \langle T_\tau (b_1 a_m) \rangle \\ \langle T_\tau (a_1^+ b_m^+) \rangle & \langle T_\tau (a_1 a_m^+) \rangle \end{pmatrix} \quad (3)$$

The non-diagonal part gives rise to a renormalized interaction $J_{\text{eff}}(k)$ summing up a whole class of diagrams. J_{eff} diverges for $d=1$ and $T=0$ whereas for finite temperatures the divergency appears for $d \leq 2$.

The commutation rule (2) leads also to additional diagrams (kinematic interaction) which contribute to the staggered magnetization M_s . We get $M_s = 0.381$ (perturbational approach, second order) or $M_s = 0.359$ (self-consistent summation) for $d = 2$ and at $T = 0$ in agreement with recent results /1/.

NON-LINEAR SIGMA MODEL

In the large spin limit one may start with a Lagrangian in terms of a unit vector field \underline{n} the average of which is related to the staggered magnetization. Alternatively we derive the Lagrangian using the semi-macroscopic equation of motion for the two relevant variables \underline{n} and \underline{l} related to the generator for spin rotations. It results /2/

$$S = \rho_s / 2 \hbar \int d\tau d^d x \left\{ (\nabla \underline{n})^2 + w^{-2} \dot{\underline{n}}^2 \right\} \quad \underline{n}^2 = 1 \quad (4)$$

Using Polyakov's renormalization group procedure /4/ we find the flow equations for the parameters $y = g \coth x$, $x = g/2t$ with $t = T/\rho_s$ and $g = h w/\rho_s$, and additional to /2/ also for the magnetization, the correlation length and the correlation function. For $d=2$ there occur a fixed point g^* above it quantum fluctuations destroy the conventional Néel order. If the action (4) is completed by an anisotropic term the quantum disordered fixed point is reduced. We have also considered the influence of a term proportional to $\underline{n} \cdot \text{curl } \underline{n}$ which favours a helical order. The value of the fixed point g^* increases. The restriction $\underline{n}^2 = 1$ can be lifted by introducing a Lagrange multiplier. Such an approach can be used in a $1/n$ expansion (n number of components).

REFERENCES

- /1/ E. Manousakis, Rev. Mod. Phys. 63, 1, 1991.
- /2/ S. Chakravarty, B. Halperin, D. Nelson, Phys. Rev. Lett. 60, 1057, 1988.
- /3/ A. Kühnel, J. Phys. C 2, 704, 1969.
- /4/ A. M. Polyakov, Phys. Lett. B 59, 79, 1975.