

Zeitschrift:	Helvetica Physica Acta
Band:	65 (1992)
Heft:	2-3
Artikel:	Low temperature properties of two dimensional frustrated quantum antiferromagnets
Autor:	Azaria, P. / Delamotte, B. / Mouhanna, D.
DOI:	https://doi.org/10.5169/seals-116482

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Low temperature properties of two dimensional frustrated quantum antiferromagnets

P. Azaria

Laboratoire de Physique Théorique des liquides, Université Pierre et Marie Curie
4 Place Jussieu, 75005 Paris, France.

B. Delamotte and D. Mouhanna

Laboratoire de Physique Théorique et Hautes Energies, Université Paris 7
2 Place Jussieu, 75005 Paris, France.

Abstract. We extend Chakravarty, Halperin and Nelson's analysis of low temperature $2D$ quantum antiferromagnets to systems which have classical canted ground states, such as helical or triangular lattice systems. In particular, we calculate the quantum correlation length ξ at two loop order and argue that the measure of ξ could be a test of the existence of long range order in these models.

Partly motivated by the study of superconductivity, many investigations have been recently devoted to the nature of the order in $D = 2$ quantum frustrated antiferromagnets. Most of these works assume a disordered phase at $T = 0$ and propose the existence of non trivial ground states. However, numerical simulations as well as analytical studies suggest that Néel order persists, even for spin $1/2$, in many canted systems[1,2,3]. We propose to test the nature of the order at $T = 0$ in frustrated canted systems by measuring the correlation length at low temperature. The comparison between experimental results and a theoretical calculation based upon the assumption of Néel order at low temperature, was already successfully performed by Chakravarty, Halperin and Nelson for collinear magnets[4]. We have calculated the quantum correlation length at two loop order for a wide class of $2D$ canted systems having zero net magnetization, such as the triangular and helical models. In contrast with the collinear case, the $O(3)$ rotation group is completely broken in the classical ground state of canted systems so that we expect three spin-waves in the quantum case, if the long range order is not destabilized by the quantum fluctuations. The hydrodynamical theory describing the long distance physics of frustrated systems with zero net magnetization, predicts a linear spectrum with a priori different velocities for these spin-waves[5]. The simplest model that reproduces the spectrum and the interaction among spin waves is the quantum non-linear sigma (NL σ) model. When two spin-waves have the same velocity (triangular, helical models, etc...), the symmetry is not only $O(3)$, but $O(3) \otimes O(2)$, where the $O(2)$ indicates that two spin-waves are equivalent[6]. The quantum NL σ model is thus $O(3) \otimes O(2)/O(2)$ and its action is[7]:

$$S = -\frac{1}{2} \int_0^\beta d\tau \int d^2x \left[\text{Tr} \left(P_0 \left(g^{-1} \partial_0 g \right)^2 + P_\perp \left(g^{-1} \partial_i g \right)^2 \right) \right] \quad (1)$$

where $g(x, \tau) \in SO(3)$, $i = 1, 2$, β is the inverse temperature and P_0, P_\perp are two diagonal matrices containing the 4 coupling constants, i.e. the 2 spin-waves velocities and the 2 spin-stiffnesses. Since the three spin-waves do not have the same velocities, i.e. P_0 is not proportional to P_\perp , the model is not Lorentz invariant. Nevertheless, we have been able to obtain the one loop recursion relations for β and the 4 parameters in eq.(1)[6]. These equations show that a non trivial, Lorentz and $O(3) \otimes O(3)/O(3) = O(4)/O(3)$ symmetric fixed point exists, at $T = 0$. The hypersurface associated to this fixed point divides the coupling constant space into a disordered phase for small S and a Néel ordered phase for large S . There also exists a whole fixed hypersurface at $T = 0$ and

$S = \infty$. This surface controls the long distance physics of the spin-waves in the Néel phase since the renormalization group flow drives the system towards it. After a sufficient number of iterations, $e^{l^*} \sim T^{-1}$, the quantum fluctuations become negligible. We end up with an effective quasi classical $2D$, $O(3) \otimes O(2)/O(2)$ NL σ model with effective parameters $P_0(l^*), P_\perp(l^*)$. The whole effect of quantum fluctuations is to renormalize the coupling constants. The quantum correlation length can then be calculated from the classical one, in which P_0, P_\perp are replaced by $P_0(l^*)$ and $P_\perp(l^*)$. We have carefully calculated at one loop order the effective parameters $P_0(l^*), P_\perp(l^*)$ by integrating the quantum fluctuations in eq.(1). We have also calculated at two loop order the classical correlation length from which follows the quantum one. We find finally, for the quantum correlation length in the Pauli-Villars regularization scheme[6a]:

$$\xi = C_\xi^{PV} \frac{\hbar c_1}{k_B T} \sqrt{\frac{2(1-\alpha)}{\rho_1}} \exp \left(\left[(1-\alpha)G(\alpha) - G(0) \right] \left[-\frac{1}{8} - \frac{1-\alpha}{16\alpha} \left(1 + \frac{\log(c_3/c_1)^2}{1 - (c_3/c_1)^2} \right) \right] \right) \exp \left(2\pi\rho_1 G(\alpha)/2k_B T \right) \quad (2)$$

where:

$$\begin{aligned} G(\alpha) &= 2 + 2(1-\alpha)(\arg \tanh \sqrt{\alpha})/\sqrt{\alpha} ; \quad 0 \leq \alpha < 1 \\ G(\alpha) &= 2 + 2(1-\alpha)(\arctan \sqrt{-\alpha})/\sqrt{-\alpha} ; \quad \alpha \leq 1 \end{aligned} \quad (3)$$

and where C_ξ^{PV} is a constant, ρ_1 and ρ_3 are spin-stiffnesses, $\alpha = 1 - \rho_3/\rho_1$ and c_1 and c_3 are spin-wave velocities. This is the general expression for the correlation length of frustrated quantum antiferromagnets with zero net magnetization, when two spin-waves have the same velocity. It contains no adjustable parameter. As argued by Chakravarty et al.[4], the constants c_1, c_3, ρ_1 and α are phenomenological input parameters that can be obtained, for instance, by a spin-wave calculation. Let us finally emphasize that our expression for ξ can be, in principle, experimentally tested on ^3He adsorbed on graphite substrate[8], since this system is thought to be modelized by a Heisenberg $S = 1/2$ model on the triangular lattice, with interactions either ferromagnetic or antiferromagnetic, depending on coverage.

References

- [1] T. Jolicoeur, E. Dagotto, E. Gagliano and S. Bacci, *Phys. Rev. B*, 42 (1990) 4800.
- [2] D. A. Huse and V. Helser, *Phys. Rev. Lett.*, 60 (1988) 2531.
- [3] C. Lhuillier, B. Bernu, P. Azaria and L. Pierre *In preparation*.
- [4] S. Chakravarty, B. I. Halperin and D. R. Nelson, *Phys. Rev. B*, 39 (1989) 2344.
- [5] B. I. Halperin and W. M. Saslow, *Phys. Rev. B*, 16 (1977) 2154.
- [6] a) P. Azaria, B. Delamotte and T. Jolicoeur, *Phys. Rev. Lett.*, 64 (1990) 3175;
b) *J. Appl. Phys.*, 69 (1991) 6170 and c) *preprint PAR/91-44*.
- [7] T. Dombre and N. Read, *Phys. Rev. B*, 39 (1989) 6797.
- [8] H. Godfrin, R. E. Rapp and D. D. Osheroff, *Physica A*, 163 (1990) 101.